arXiv:math/0310414v1 [math.SG] 26 Oct 2003

HOMOLOGICAL MIRROR SYMMETRY FOR THE QUARTIC SURFACE

PAUL SEIDEL

Warning. This preprint is in finished form, and should be readable by itself. However, the argument uses certain general properties of Fukaya categories, the proofs of which will appear in [61] (under preparation). I apologize for this unsatisfactory state of affairs.

Contents

1. Introduction 1
2. A_∞-categories 6
3. Deformation theory 12
4. Group actions 24
5. Coherent sheaves 29
6. Symplectic terminology 33
7. Monodromy and negativity 43
8. Fukaya categories 58
9. Computations in Fukaya categories 74
10. The algebras Q_4 and Q_{64} 83
11. Counting polygons 94

References 120

1. Introduction

This paper deals with a special case of Kontsevich's "homological mirror symmetry" conjecture [36]. Before formulating the precise statement, it is necessary to introduce the relevant coefficient rings.

Notation 1.1. *Let* $\Lambda_{\mathbb{N}} = \mathbb{C}[[q]]$ *be the ring of formal power series in one variable. We denote by* $\Lambda_{\mathbb{Z}}$ *its quotient field, obtained by formally inverting* q*. By adjoining roots* $q^{1/d}$ *of all orders to* $\Lambda_{\mathbb{Z}}$*, one obtains its algebraic closure* $\Lambda_{\mathbb{Q}}$*, known in symplectic geometry as the rational Novikov field. Elements of this*

Date: This copy LaTeXed: February 1, 2008.

field are formal series

$$f(q) = \sum_m a_m q^m$$

where m *runs over all numbers in* $(1/d)\mathbb{Z} \subset \mathbb{Q}$ *for some* $d \geq 1$ *which depends on* f*, and* $a_m \in \mathbb{C}$ *vanishes for all sufficiently negative* m*. Geometrically, one should think of* $\Lambda_{\mathbb{N}}$ *as a small disc, of* $\Lambda_{\mathbb{Z}}$ *as the punctured disc, and of* $\Lambda_{\mathbb{Q}}$ *as its universal cover. Consider the semigroup* $End(\Lambda_{\mathbb{N}})$ *of* q*-adically continuous* $\mathbb{C}$*-algebra endomorphisms of* $\Lambda_{\mathbb{N}}$*. More concretely, these are substitutions*

$$\psi^* : q \longmapsto \psi(q)$$

with $\psi \in q\mathbb{C}[[q]]$*. Each nonzero* ψ *extends to an endomorphism of the quotient field, and the* q*-adically continuous Galois group of* $\Lambda_{\mathbb{Z}}/\mathbb{C}$ *is the group of invertibles* $End(\Lambda_{\mathbb{N}})^\times$*, consisting of those* ψ *with* $\psi'(0) \neq 0$*. Moreover, the Galois group of* $\Lambda_{\mathbb{Q}}/\Lambda_{\mathbb{Z}}$ *is the profinite group* $\hat{\mathbb{Z}}$*, whose topological generator* $\hat{1}$ *takes* $q^{1/d}$ *to* $e^{2\pi i/d}q^{1/d}$*.*

On the symplectic side of mirror symmetry, take any smooth quartic surface $X_0 \subset \mathbb{C}\mathrm{P}^3$, with its standard symplectic structure. To this we associate a triangulated category linear over $\Lambda_{\mathbb{Q}}$, the split-closed derived Fukaya category $D^\pi\mathcal{F}(X_0)$, defined using Lagrangian submanifolds of X_0 and pseudo-holomorphic curves with boundary on them. The use of formal power series to take into account the area of pseudo-holomorphic curves is a familiar device. $\Lambda_{\mathbb{Q}}$ appears because we allow only a certain kind of Lagrangian submanifolds, namely the rational ones (there is also another restriction, vanishing of Maslov classes, which is responsible for making the Fukaya category $\mathbb{Z}$-graded).

On the complex side, we start with the quartic surface in $\mathbb{P}^3_{\Lambda_{\mathbb{Q}}}$ defined by

$$y_0y_1y_2y_3 + q(y_0^4 + y_1^4 + y_2^4 + y_3^4) = 0. \tag{1.1}$$

Note that here q is a “constant” (an element of the ground field). The group $\Gamma_{16} = \{[diag(a_0, a_1, a_2, a_3)] \;:\; a_k^4 = 1,\; a_0a_1a_2a_3 = 1\} \subset PSL(V)$, $\Gamma_{16} \cong \mathbb{Z}/4 \times \mathbb{Z}/4$, acts on this surface in the obvious way. We denote by Z_q^* the unique minimal resolution of the quotient orbifold. Non-algebraic-geometers may wish to remind themselves that $\Lambda_{\mathbb{Q}}$ is an algebraically closed field of characteristic zero, so that all the geometric constructions like minimal resolutions go through as in the familiar case of $\mathbb{C}$. In fact, Z_q^* could already be defined over the smaller field $\Lambda_{\mathbb{Z}}$, since the fixed points of the action that need to be resolved are $\Lambda_{\mathbb{Z}}$-points. The associated category is the bounded derived category of coherent sheaves, $D^bCoh(Z_q^*)$, which is also a triangulated category linear over $\Lambda_{\mathbb{Q}}$.

Theorem 1.2. *There is a* $\psi \in End(\Lambda_{\mathbb{N}})^\times$ *and an equivalence of triangulated categories,*

$$D^\pi\mathcal{F}(X_0) \cong \hat{\psi}^* D^bCoh(Z_q^*).$$

Here $\hat{\psi}$ is a lift of ψ to an automorphism of $\Lambda_{\mathbb{Q}}$, which we use to change the $\Lambda_{\mathbb{Q}}$-module structure of $D^bCoh(Z_q^*)$, by letting $\hat{\psi}^*f$ act instead of f. The theorem says that the outcome of this "base change" is equivalent to $D^\pi\mathcal{F}(X_0)$. The choice of lift $\hat{\psi}$ is irrelevant because $D^bCoh(Z_q^*)$ carries a $\hat{\mathbb{Z}}$-action (and so does $D^\pi\mathcal{F}(X_0)$ by [17]).

The function ψ which occurs in the theorem is unknown at present (or rather, we cannot verify that it agrees with the standard "mirror map"). Nevertheless, certain consequences can already be drawn from the result as given. For instance, let $\mathcal{M}^*$ be the classifying space (or moduli stack, depending on the reader's preference) for smooth $K3$ surfaces equipped with an ample cohomology class A whose square is $A \cdot A = 4$, and with a choice of nonzero holomorphic two-form. There is a canonical homomorphism $\pi_1(\mathcal{M}^*) \rightarrow \pi_0(Aut(X_0))$, which is defined by symplectic monodromy of families of $K3$ surfaces equipped with Kähler forms in the class A. On the other hand, since $D^\pi\mathcal{F}(X_0)$ is a symplectic invariant, we have a map $\pi_0(Aut(X_0)) \rightarrow Aut(D^\pi\mathcal{F}(X_0)/\Lambda_{\mathbb{Q}})/[2]$, where now Aut on the right hand side is the group of isomorphism classes of $\Lambda_{\mathbb{Q}}$-linear autoequivalences, and we divide by the subgroup generated by even translations. Actually, the monodromy maps obtained from loops in $\mathcal{M}^*$ are naturally graded, so the shift ambiguity can be removed, and we get a homomorphism

$$\pi_1(\mathcal{M}^*) \longrightarrow Aut(D^bCoh(Z_q^*)/\Lambda_{\mathbb{Q}}). \tag{1.2}$$

This had been postulated by several people, starting with Kontsevich himself. Interestingly, even though both $\mathcal{M}^*$ and $D^bCoh(Z_q^*)$ are objects from algebraic geometry, there seems to be no definition of the map which avoids symplectic geometry and mirror symmetry. Previous work on Dehn twists [62] shows that (1.2) is highly nontrivial, but the exact kernel and cokernel remain unknown. Another direct implication of Theorem 1.2 is the equality of certain simpler invariants, such as the Grothendieck groups

$$K_0(D^\pi\mathcal{F}(X_0)) \cong K_0(D^bCoh(Z_q^*)) = K_0(Z_q^*).$$

Every object of the Fukaya category (in particular, every oriented Lagrangian sphere in X_0) determines a class in $K_0(D^\pi\mathcal{F}(X_0))$, and $\pi_0(Aut(X_0))$ acts linearly on this group. On the other hand $K_0(Z_q^*) \otimes \mathbb{Q} \cong CH_*(Z_q^*) \otimes \mathbb{Q}$ by Grothendieck's theorem, so it contains $CH_0(Z_q^*) \otimes \mathbb{Q}$ which is quite an interesting object.

Having stated our result, it is maybe time to review briefly some other work on homological mirror symmetry. The first case to be considered was that of elliptic curves, where Polishchuk and Zaslow [50] proved a result roughly similar to Theorem 1.2, with an explicit mirror map ψ but not taking into account the triangulated structure (subsequent work on Massey products [49] largely makes up for this deficiency). There are many beautiful and deep results on abelian varieties [19, 37, 14, 46], but none of them comes close to describing the entire Fukaya category. A common thread of all these papers is to establish an explicit bijection between objects on both sides (Lagrangian submanifolds and sheaves). The prevailing view is that this correspondence should be set up as a Fourier-Mukai type transform associated to a Strominger-Yau-Zaslow torus fibration. This is the approach of [37], in the situation where the SYZ fibration has no singularities, and where the Lagrangian submanifolds are supposed to be transverse to the fibres. Both restrictions are serious ones from the point of symplectic topology: work on removing them has started, see in particular [19], but it may be some time before a complete thery emerges.

Actually, this entire circle of ideas is of little importance for our proof (although the SYZ conjecture does serve as a source of intuition in some places). We rely instead on techniques from symplectic topology and homological algebra, which are maybe closer to the spirit of Kontsevich's original paper. The computation on the algebraic geometry side relies on a classical idea of Beilinson, together with a Massey product computation which the author learned from [13]. On the symplectic side, where the overwhelming majority of the work is invested, there are three steps, namely (slightly simplified, and in reverse logical order):

- First, we prove that $D^\pi\mathcal{F}(X_0)$ can be entirely reconstructed from the full A_∞-subcategory of $\mathcal{F}(X_0)$ consisting of a particular set of 64 Lagrangian two-spheres, which are vanishing cycles for the standard Fermat pencil (the choice of these cycles is best explained by physics considerations, compare for instance [31]). The necessary "split-generation" argument resembles the ones in [61], with some added geometric input concerning the monodromy around the singular fibre at infinity.

- By construction, all 64 vanishing cycles lie in an affine Zariski-open subset $M_0 \subset X_0$. Following a proposal from [59] the relation between the Fukaya categories $\mathcal{F}(M_0)$ and $\mathcal{F}(X_0)$ can be formulated in terms of the deformation theory of A_∞-structures. This recourse to abstract deformation theory is the reason why we cannot determine ψ explicitly.

- Having reduced our problem to a finite collection of Lagrangian two-spheres in an affine $K3$ surface, we now apply a general dimensional induction argument from [61] to compute the relevant subcategory of $\mathcal{F}(M_0)$. This starts with an application of Picard-Lefschetz theory in one dimension lower than before, fibering our $K3$ surface by a pencil of curves.

The last-mentioned argument involves a certain amount of computation, of a combinatorial (drawing loops on surfaces, and counting polygons) and homological (determining morphisms between certain twisted complexes in A_∞-categories) nature. Although the algorithm is simple, the computations themselves are not particularly enlightening, and in fact a computer was used to do parts of them. The reason why this step in the proof is less transparent is that, unlike the previous two, it has no known meaning in terms of the mirror. This is partially due to the fact that the fibres of our pencil are curves of genus 3, hence outside the scope of mirror symmetry.

In principle, the approach taken here is fairly general. Suppose for instance that one wanted to adapt it to the quintic threefold case. The deformation theoretic part actually becomes easier since, unlike the situation for $K3$s, there are no noncommutative deformations of the mirror variety. On the other hand, the definition of the Fukaya category is much more complicated [18], and one would have to generalize the twist exact sequence [60] accordingly, but these are on the whole technical issues. There are also some elementary geometric complications, which arise where we have relied on intuition provided by the SYZ conjecture. The final obstacle is in the computational part, where the procedure remains precisely the same but the complexity rises by a factor of at least 10. This raises serious practical concerns, but it may still be feasible, or one can try to find a more conceptual way of carrying it out (ideas of Moishezon might prove helpful, see the comments and computations in [5]).

The paper is structured as follows. In the first two parts we assemble the basic algebraic (Sections 2–5) and geometric (Sections 6–9) techniques. Much of this is not really new, but the application to mirror symmetry often results a change of perspective, which means that it is hard to find the theorems stated elsewhere in the form in which we need them. The level of generality varies somewhat: we have tried to include generalizations which were reasonably easy or natural, and to exclude those which are either irrelevant to homological mirror symmetry, or which would require a significant amount of additional techniques or terminology. The last part of the paper (Sections 10–11) contains the argument which is really specific to the quartic surface, including figures and tables which list the results of the computational step. The aim of that was to give enough data so that the reader desirous of verifying the computation independently could break up that process into manageable steps.

Acknowledgments. This paper could not have been written without the assistance of Michael Douglas, who showed me the “quiver presentation” of the derived category of coherent sheaves which appears as the algebras Q_4 and Q_{64} in this paper, and also the crucial Massey product formula (Lemma 10.8) from [13]. Maxim Kontsevich helped me in many ways, one of them being the Hochschild cohomology computations for semidirect products (Lemma 4.2). Simon Donaldson introduced me to the notion of matching cycle, on which the main dimensional induction procedrure is based. To all of them, thanks! ETH Zürich invited me to give a lecture course on Fukaya categories, and provided a responsive audience on which parts of this paper were tested. The following computer programs have been used: SINGULAR for elimination theory, MATHEMATICA for visualization, and MAPLE for the homological algebra part.

2. A_∞-CATEGORIES

This section is designed to serve as quick reference for the basic algebraic terminology, for the benefit of readers with a geometry background. The main construction is that of the bounded derived category of a A_∞-category, following Kontsevich [36]. We then consider the idempotent (Karoubi, or split-closed) completion, and the corresponding notion of split-generators. A brief look at twist functors and exceptional collections completes the tour. Some references for more in-depth treatments of various matters touched on here are [35, 34, 40, 55, 18, 16, 61].

(2a) A non-unital A_∞-category $\mathcal{C}$ consists of a set (all our categories are small) of objects $Ob\,\mathcal{C}$, together with a graded $\mathbb{C}$-vector space $hom_{\mathcal{C}}(X_0,X_1)$ for any two objects, and multilinear maps

$$\mu^d_{\mathcal{C}} : hom_{\mathcal{C}}(X_{d-1},X_d) \otimes \cdots \otimes hom_{\mathcal{C}}(X_0,X_1) \longrightarrow hom_{\mathcal{C}}(X_0,X_d)[2-d] \tag{2.1}$$

for all $d \geq 1$ and all $(d+1)$-tuples of objects $(X_0,\dots,X_d)$, satisfying the A_∞-associativity equations

$$\sum_{e,i} (-1)^* \mu^{d-e+1}_{\mathcal{C}}(a_d,\dots,a_{i+e+1},\mu^e_{\mathcal{C}}(a_{i+e},\dots,a_{i+1}),a_i,\dots,a_1) = 0, \tag{2.2}$$

where $* = |a_1| + \cdots + |a_i| - i$ (the decreasing numbering of the a_k is the result of trying to bring together various standard conventions). The underlying (non-unital) cohomological category $H(\mathcal{C})$ has the same objects as $\mathcal{C}$, with

$$Hom^*_{H(\mathcal{C})}(X_0,X_1) = H^*(hom_{\mathcal{C}}(X_0,X_1),\mu^1_{\mathcal{C}})$$

and composition induced by the cochain level map $a_2a_1 = (-1)^{|a_1|}\mu^2_{\mathcal{C}}(a_2,a_1)$. There is also the full subcategory $H^0(\mathcal{C})$ which retains only the morphisms of degree zero. One says that $\mathcal{C}$ is *c-unital* (cohomologically unital) if $H^0(\mathcal{C})$ has identity morphisms. It is true, but by no means obvious, that this is a good class of A_∞-categories to work with: the proofs of many properties rely on the fact that c-unitality turns out to be equivalent to the less flexible but more visibly well-behaved notion of strict unit [40, 61]. *From now on, all A_∞-categories are assumed to be c-unital.* Similarly, when considering A_∞-functors $\mathcal{F} : \mathcal{C} \to \mathcal{D}$, we always require that $H(\mathcal{F}) : H(\mathcal{C}) \to H(\mathcal{D})$ be unital (carries identity morphisms to identity morphisms). $\mathcal{F}$ is called a quasi-equivalence if $H(\mathcal{F})$ is an equivalence. One can prove that then, there is a $\mathcal{G} : \mathcal{D} \to \mathcal{C}$ such that $H(\mathcal{G})$ is an inverse equivalence to $H(\mathcal{F})$.

Remark 2.1. *A linear category with one object is the same as an associative algebra, and similarly, an A_∞-category with one object is an A_∞-algebra. Slightly more generally, given a category C with only finitely many objects $X_1,\dots,X_m$, one can form the* total morphism algebra

$$A = \bigoplus_{j,k} Hom_C(X_j,X_k)$$

which is linear over the semisimple ring $R_m = \mathbb{C} \oplus \cdots \oplus \mathbb{C}$ *(*m *summands). Again, the same holds for* A_∞*-categories versus* R_m*-linear* A_∞*-algebras, and we will use that frequently. Our notation will be that* $u_1, \dots, u_m$ *are the orthogonal idempotents in* R_m*, so* $u_k A u_j = Hom_C(X_j, X_k)$*.*

Remark 2.2. *For concreteness, we stick to the ground field* $\mathbb{C}$ *for our exposition of the basic algebraic theory, but any other field would do just as well (later on though, we will make use of deformation theoretic ideas which work best in characteristic* 0*). Defining* A_∞*-categories over a ring is a more difficult issue, and the only simple solution is to assume that all the hom spaces are projective modules. That of course holds automatically for the semisimple rings* R_m*. The only other ring of interest to us is* $\Lambda_{\mathbb{N}}$*, which requires special treatment because of its topology. Recall that for a* $\Lambda_{\mathbb{N}}$*-module* V_q *which is* q*-adically complete, the following conditions are equivalent: (1)* V_q *is torsion-free, which means that multiplication by* q *is an injective endomorphism; (2)* V_q *is topologically free, which means that projection* $V_q \to V$ *to the vector space* $V = V_q \otimes_{\Lambda_q} \mathbb{C} = V_q/qV_q$ *can be lifted to a topological isomorphism* $V_q \to V \hat{\otimes} \Lambda_{\mathbb{N}}$*. A* topological A_∞-algebra *over* $\Lambda_{\mathbb{N}}$ *consists of a graded* $\Lambda_{\mathbb{N}}$*-module* $\mathcal{A}_q$ *such that each degree* k *piece is complete and torsion-free. If* $H(\mathcal{A}_q)$ *is torsion-free, it is automatically also complete.*

(2b) The twisted complex construction embeds $\mathcal{C}$ into a larger A_∞-category which is closed under (appropriately understood) mapping cones. This can be broken into two steps:

(1) The additive enlargement $\Sigma\mathcal{C}$ has objects which are formal sums

$$X = \bigoplus_{f \in F} X_f[\sigma_f],$$

where F is some finite set, $X_f \in Ob\,\mathcal{C}$, and $\sigma_f \in \mathbb{Z}$. The morphisms between any two objects are

$$hom_{\Sigma\mathcal{C}}\Big(\bigoplus_{f\in F} X_f[\sigma_f], \bigoplus_{g \in G} Y_g[\tau_g]\Big) = \bigoplus_{f,g} hom_{\mathcal{C}}(X_f, Y_g)[\tau_g - \sigma_f],$$

and the compositions are defined using those of $\mathcal{C}$ and the obvious "matrix multiplication" rule, with additional signs put in as follows: for $a_1 \in hom(X_0[\sigma_0], X_1[\sigma_1]), \dots, a_d \in hom(X_{d-1}[\sigma_{d-1}], X_d[\sigma_d])$,

$$\mu^d_{\Sigma\mathcal{C}}(a_d, \dots, a_1) = (-1)^{\sigma_0} \mu^d_{\mathcal{C}}(a_d, \dots, a_1).$$

(2) A twisted complex is a pair (C, δ) consisting of $C \in Ob\,\Sigma\mathcal{C}$ and $\delta \in hom^1(C, C)$, with the "upper triangularity" property that the indexing set F for $C = \bigoplus_{f \in F} C_f$ can be ordered in such a way that all components δ_{fg} with $f \geq g$ are zero, and subject to the generalized Maurer-Cartan equation

$$\mu^1_{\Sigma\mathcal{C}}(\delta) + \mu^2_{\Sigma\mathcal{C}}(\delta, \delta) + \cdots = 0.$$

Upper triangularity says that this is a finite sum, hence the condition makes sense. We define an A_∞-category $Tw\mathcal{C}$ of which the (C,δ) are the objects. The spaces of morphisms are the same as for $\Sigma\mathcal{C}$, but composition involves the δs: for $a_k \in hom((C_{k-1},\delta_{k-1}),(C_k,\delta_k))$,

$$\mu^d_{Tw\mathcal{C}}(a_d,\dots,a_1) = \sum_{j_0,\dots,j_d\geq 0} \mu^d_{\Sigma\mathcal{C}}(\overbrace{\delta_d,\dots,\delta_d}^{j_d}, a_d, \overbrace{\delta_{d-1},\dots,\delta_{d-1}}^{j_{d-1}},\dots).$$

The c-unitality of $\mathcal{C}$ implies that of $Tw\mathcal{C}$. The underlying cohomological category in degree zero, which means we retain only the H^0 of the morphisms spaces, is (somewhat improperly) called the *bounded derived category* $D^b(\mathcal{C}) = H^0(Tw\mathcal{C})$. Given two twisted complexes (C_k,δ_k), $k=0,1$, and a morphism $a \in hom^0(C_0,C_1)$ such that $\mu^1_{Tw\mathcal{C}}(a) = 0$, one can define the mapping cone as usual:

$$Cone(a) = \left\{C_0 \xrightarrow{a} C_1\right\} \overset{\text{def}}{=} \left(C_0[1]\oplus C_1, \begin{pmatrix} \delta_0 & 0 \\ a & \delta_1 \end{pmatrix}\right) \in Ob\,Tw\,\mathcal{C}.$$

The isomorphism class of this in the derived category depends only on the cohomology class $\alpha = [a] \in Hom_{D^b(\mathcal{C})}(C_0,C_1)$, hence we will sometimes write sloppily $Cone(\alpha)$. Declare *exact triangles* in the derived category to be those which are isomorphic to the standard mapping cone triangles $C_0 \to C_1 \to Cone(a) \to C_0[1]$. This makes $D^b(\mathcal{C})$ into a triangulated category. It contains $H^0(\mathcal{C})$ as a full subcategory, and the objects of that subcategory *generate* $D^b(\mathcal{C})$ in the usual triangulated sense, that is, any object of $D^b(\mathcal{C})$ can be obtained from objects of $H^0(\mathcal{C})$ by successive mapping cones and isomorphism. However, this does not mean that $H^0(\mathcal{C})$ determines $D^b(\mathcal{C})$. For instance, triangulated categories have intrinsically defined Massey product operations [20], and in our case these involve the higher order $\mu^d_{\mathcal{C}}$.

The idempotent completion of an arbitrary linear category has objects which are pairs (X,p), where X is an object of the original category and $p \in Hom(X,X)$ an idempotent endomorphism. One thinks of this as the "direct summand $im(p)$", and so $Hom((X_0,p_0),(X_1,p_1)) = p_1 Hom(X_0,X_1)p_0$. It is proved in [6] that the idempotent completion of a triangulated category is again triangulated, in a unique way which makes the inclusion $X \mapsto (X, id_X)$ into an exact functor. The result of applying this construction to the bounded derived category of an A_∞-category is called the *split-closed derived category* $D^\pi(\mathcal{C})$. Suppose that we have a set of objects in an idempotent complete triangulated category; these are called *split-generators* if one can obtain any object from them by successive mapping cones, splitting off direct summands, and isomorphism.

Here are some obvious but useful facts about derived categories:

Lemma 2.3. *If two A_∞-categories are quasi-equivalent, their derived categories D^b and D^π are equivalent triangulated categories.*

Proof. Any A_∞-functor $\mathcal{F} : \mathcal{C} \to \mathcal{D}$ admits a canonical extension $Tw\mathcal{F} : Tw\mathcal{C} \to Tw\mathcal{D}$, and $D^b\mathcal{F} = H^0(Tw\mathcal{F}) : D^b(\mathcal{C}) \to D^b(\mathcal{D})$ is exact (it takes cones to cones). Suppose that $\mathcal{F}$ is a quasi-equivalence, and consider the subcategory of $D^b(\mathcal{C})$ consisting of those objects X such that $D^b\mathcal{F}$ maps $Hom^*(X,Y)$ isomorphically to $Hom^*(D^b\mathcal{F}(X), \mathcal{F}(Y))$ for all $Y \in Ob\,H^0(\mathcal{C})$. By assumption this subcategory contains $H^0(\mathcal{C})$, and it is a standard exercise to show that it is triangulated, so it is the whole of $D^b(\mathcal{C})$. Repetition of the argument proves that $D^b\mathcal{F}$ is full and faithful. Its image is a triangulated subcategory of $D^b(\mathcal{D})$ which, up to isomorphisms, contains all objects of $H^0(\mathcal{D})$. Since these are generators for $D^b(\mathcal{D})$, we conclude that $D^b(\mathcal{F})$ is an equivalence. The corresponding statement for D^π follows immediately. □

Lemma 2.4. *Let $\mathcal{C}' \subset \mathcal{C}$ a full A_∞-subcategory such that the set of all objects of $\mathcal{C}'$ split-generates $D^\pi(\mathcal{C})$. Then $D^\pi(\mathcal{C}') \cong D^\pi(\mathcal{C})$ as triangulated categories.*

Proof. Inclusion induces a full and faithful functor $D^\pi(\mathcal{C}') \to D^\pi(\mathcal{C})$, the image of which contains the objects of $H^0(\mathcal{C}')$ and is closed under cones and idempotent splittings. □

Combining the two previous Lemmas, we have

Lemma 2.5. *Let $\mathcal{C}, \mathcal{D}$ be two A_∞-categories. Suppose that they contain full A_∞-subcategories $\mathcal{C}' \subset \mathcal{C}, \mathcal{D}' \subset \mathcal{D}$ which are quasi-equivalent to each other. If the objects of $\mathcal{C}', \mathcal{D}'$ split-generate $D^\pi(\mathcal{C})$ respectively $D^\pi(\mathcal{D})$, then these two derived categories are equivalent.* □

(2c) Let C be a $\mathbb{C}$-linear triangulated category, and X, Y two objects such that $Hom^*_C(X,Y) = \bigoplus_{k\in\mathbb{Z}} Hom_C(X, Y[k])$ is finite-dimensional. Then one can form two new objects $T_X(Y)$, $T^\vee_Y(X)$ which fit into exact triangles

$$T_X(Y)[-1] \to Hom^*_C(X,Y) \otimes X \xrightarrow{ev} Y \to T_X(Y),$$

$$T^\vee_Y(X) \to X \xrightarrow{ev^\vee} Hom^*_C(X,Y)^\vee \otimes Y \to T^\vee_Y(X)[1].$$

The notation means that we choose a homogeneous basis $\{f_k\}$ of $Hom^*(X,Y)$, set $Hom^*(X,Y) \otimes X = \bigoplus_k X[-deg(f_k)]$, and take the evaluation map $ev = \bigoplus_k f_k[-k]$. The dual map $ev^\vee$ is defined in the same way. These algebraic *twist* and *untwist* operations are well-defined up to non-unique isomorphism, and the isomorphism type of the new objects depends only on those of X and Y.

The construction becomes somewhat more functorial in the case where $C = D^b(\mathcal{C})$ is the bounded derived category of an A_∞-category $\mathcal{C}$ such that (1) the cochain level morphism space $hom_{Tw\mathcal{C}}(X,Y)$ is itself finite-dimensional, and (2) X has a strict cochain level identity morphism e_X. One can then form the tensor product $hom(X,Y) \otimes X \in Ob\,Tw(\mathcal{C})$, which is a finite direct sum of shifted copies of X with a differential $\mu^1_{hom(X,Y)} \otimes e_X$, and $T_X(Y)$ can be defined as a twisted complex:

$$T_X(Y) = \{hom^*(X,Y) \otimes X \longrightarrow Y\}. \tag{2.3}$$

In fact, if the finite-dimensionality condition holds for all Y, one can realize T_X by an A_∞-endofunctor of $Tw\mathcal{C}$, inducing an exact functor on $D^b(\mathcal{C})$. Similarly, if (1) holds and (2') Y has a strict cochain level identity morphism e_Y, then $T_Y^\vee(X) = \{X \rightarrow hom^*(X,Y)^\vee \otimes Y\}$ will do. Fukaya categories are not in general strictly unital, so the previous observation does not apply directly to them. This can be remedied, but instead of entering into a full discussion, we just record the technical statement that will be used later on:

Lemma 2.6. *Let $\mathcal{C}$ be a c-unital A_∞-category whose hom spaces are finite-dimensional, and X, Y, Z objects of it. Any pair*

$$k \in hom^0_{\mathcal{C}}(Y,Z), \quad h : hom_{\mathcal{C}}(X,Y) \longrightarrow hom_{\mathcal{C}}(X,Z)[-1]$$

satisfying

$$\mu^1_{\mathcal{C}}(k) = 0, \quad \mu^1_{\mathcal{C}} h(a) + h(\mu^1_{\mathcal{C}} a) + \mu^2_{\mathcal{C}}(k,a) = 0 \tag{2.4}$$

for all $a \in hom_{\mathcal{C}}(X,Y)$, determines a morphism $T_X(Y) \rightarrow Z$ in $D^b(\mathcal{C})$. This is an isomorphism if for all objects W of $\mathcal{C}$, the following complex of vector spaces is acyclic:

$$\begin{aligned} &(hom(X,Y) \otimes hom(W,X))[2] \oplus hom(W,Y)[1] \oplus hom(W,Z), \\ &\partial(a_2 \otimes a_1, b, c) = \big(\mu^1_{\mathcal{C}}(a_2) \otimes a_1 + (-1)^{|a_2|-1} a_2 \otimes \mu^1_{\mathcal{C}}(a_1), \mu^1_{\mathcal{C}}(b) \\ &\qquad + \mu^2_{\mathcal{C}}(a_2,a_1), -\mu^1_{\mathcal{C}}(c) + \mu^2_{\mathcal{C}}(k,b) + \mu^3_{\mathcal{C}}(k,a_2,a_1) + \mu^2(h(a_2),a_1)\big). \end{aligned} \tag{2.5}$$

Proof. Suppose first that $\mathcal{C}$ is strictly unital. Then (k,h) do indeed define a μ^1-closed cochain level morphism from the object $T_X(Y)$ as defined in (2.3) to Z. Composition with this morphism is a map of complexes of vector spaces

$$hom_{Tw\mathcal{C}}(W, T_X(Y)) \longrightarrow hom_{\mathcal{C}}(W,Z)$$

and (2.5) is the mapping cone of it, so the assumption that this is acyclic means that our morphism induces isomorphisms

$$Hom^*_{D^b(\mathcal{C})}(W, T_X(Y)) \cong Hom^*_{D^b(\mathcal{C})}(W,Z)$$

for all $W \in Ob\,\mathcal{C}$. A straightforward long exact sequence argument extends this to objects W in the derived category, and then the Yoneda Lemma implies that $T_X(Y) \cong Z$.

In the c-unital case, one argues as follows. There is a differential graded category $\mathcal{C}^\vee = \mathcal{F}un(\mathcal{C}, Ch)$ of c-unital A_∞-functors from $\mathcal{C}$ to the dg category Ch of finite-dimensional chain complexes over $\mathbb{C}$, and a natural A_∞-functor $\mathcal{C} \rightarrow \mathcal{C}^\vee$, which on cohomology is full and faithful. This extends to an A_∞-functor $Tw\mathcal{C} \rightarrow \mathcal{C}^\vee$, which induces a full embedding of triangulated categories $D^b(\mathcal{C}) \hookrightarrow H^0(\mathcal{C}^\vee)$. Since $\mathcal{C}^\vee$ is strictly unital, one can define the object $T_X(Y)$ in $H^0(\mathcal{C}^\vee)$ as in (2.3); of course, this will be non-canonically isomorphic to the object of the same name in the subcategory $D^b(\mathcal{C})$. By unravelling the definitions one sees that the pair (h,k) gives a morphism $T_X(Y) \to Z$ in $H^0(\mathcal{C}^\vee)$; one proves as before that this is an isomorphism. $\square$

Lemma 2.7. *Let C be a $\mathbb{C}$-linear idempotent closed triangulated category, such that $Hom^*_C(X,Y)$ is finite-dimensional for all X, Y. Take a collection of objects $X_1, \dots, X_m$ and another object Y, and consider the sequence of objects $Y_0 = Y$, $Y_{d+1} = T_{X_1} \dots T_{X_m}(Y_d)$. Suppose there is a $d > 0$ such that $Hom_C(Y, Y_d) = 0$ (that is, there are no nontrivial morphisms of degree zero). Then Y lies in the idempotent closed triangulated subcategory of C split-generated by $X_1, \dots, X_m$.*

Proof. By definition, we have exact triangles

$$\begin{tikzcd}
T_{X_{i+1}} \dots T_{X_m}(Y_j) \arrow[rr] & & T_{X_i} \dots T_{X_m}(Y_j) \arrow[dl, "{[1]}"] \\
& Hom^*_C(X_i, T_{X_{i+1}} \dots T_{X_m}(Y_j)) \otimes X_i \arrow[ul, "ev"] &
\end{tikzcd} \tag{2.6}$$

The octahedral axiom allows us to conflate these, for all i and all $j < k$, into a single exact triangle

$$\begin{tikzcd}
Y \arrow[rr] & & Y_d \arrow[dl, "{[1]}"] \\
& R_d \arrow[ul] &
\end{tikzcd} \tag{2.7}$$

where R_d is an object built by taking repeated mapping cones from the objects in the same positions in (2.6), hence which lies in the triangulated subcategory generated by $X_1, \dots, X_m$. The assumption says that the $\to$ in (2.7) must be zero, hence $R_d \cong Y \oplus Y_d[-1]$. $\square$

Our applications of this Lemma will be along the following line: if $C = D^\pi(\mathcal{C})$ and there are fixed $X_1, \dots, X_m \in Ob\, H^0(\mathcal{C})$ such that any $Y \in Ob\, H^0(\mathcal{C})$ has the property required above, then the X_k are split-generators for $D^\pi(\mathcal{C})$.

(2d) Let $C^\to$ be a linear graded category with only finitely many objects $(X_1, \dots, X_m)$, ordered in a fixed way. We say that $C^\to$ is *directed* if

$$Hom^*_{C^\to}(X_i, X_j) = \begin{cases} 0 & i > j, \\ \mathbb{C} \cdot id_{X_i} & i = j, \\ \text{finite-dimensional} & i < j. \end{cases} \tag{2.8}$$

If C is a linear $\mathbb{C}$-graded category with finite-dimensional Hom^*'s, and $\Gamma = (X_1, \dots, X_m)$ an ordered finite family of nonzero objects in it, we can define the associated directed subcategory to be the subcategory $C^\to \subset C$ with the same objects, which is directed and satisfies $Hom^*_{C^\to}(X_i, X_j) = Hom^*_C(X_i, X_j)$ for $i < j$. This is a full subcategory iff the X_i form an *exceptional collection*, which means that their morphisms in C already satisfy the conditions of (2.8).

Similarly, let $\mathcal{C}^{\rightarrow}$ be an A_∞-category with objects $(X_1, \dots, X_m)$. We call it directed if the *hom*-spaces satisfy the same conditions as in (2.8), with the additional condition that there are generators e_i of $hom(X_i, X_i)$ which are units in the strict sense. Given a finite collection of objects $(X_1, \dots, X_m)$ in an arbitrary A_∞-category $\mathcal{C}$ with finite-dimensional *hom*'s, one can define an associated directed A_∞-category $\mathcal{C}^{\rightarrow}$ by requiring that $hom_{\mathcal{C}^{\rightarrow}}(X_i, X_j) = hom_{\mathcal{C}}(X_i, X_j)$ for $i < j$, and $\mu^d_{\mathcal{C}^{\rightarrow}}(a_d, \dots, a_1) = \mu^d_{\mathcal{C}}(a_d, \dots, a_1)$ for all $a_k \in hom(X_{i_{k-1}}, X_{i_k})$ with $i_0 < \dots < i_d$. This is not really an A_∞-subcategory of $\mathcal{C}$ (unless that happens to have strict units), but after some more work one can still construct an A_∞-functor $\mathcal{C}^{\rightarrow} \rightarrow \mathcal{C}$ which maps the $hom(X_i, X_j)$, $i < j$, trivially. We will therefore tolerate a slight abuse of terminology, and call $\mathcal{C}^{\rightarrow}$ the *directed A_∞-subcategory* associated to $(X_1, \dots, X_m)$.

The main interest in exceptional collections and directed categories comes from the notion of mutations. Occasional glimpses of this theory will be visible later in the paper, in particular in Section 9, but it is not strictly necessary for our purpose, so we will just refer the interested reader to [55, 35, 58].

3. Deformation theory

We need to put together some material (well-known in the right circles) from the classification theory of A_∞-structures. Using the Homological Perturbation Lemma [26], the "moduli space" of quasi-isomorphism types of A_∞-algebras with fixed cohomology algebra can be written as the quotient of an infinite-dimensional space of solutions to the A_∞-equations, by the action of an infinite-dimensional "gauge" group. There is a canonical base point, represented by the trivial A_∞-structure (the one where the higher order composition maps are all zero). The formal "Zariski tangent space" at this point is a product of suitable Hochschild cohomology groups of the cohomology algebra. Moreover, the "moduli space" carries a $\mathbb{C}$-action which contracts it to the base point. As noted by Kadeishvili [32], this implies that the "tangent space" already carries a lot of information about the "moduli space". A more straightforward argument along the same lines applies to the classification of infinitesimal deformations of a fixed possibly nontrivial A_∞-structure, using a suitable generalization of Hochschild cohomology [21]. We will recall these and some similar ideas, aiming in each case for uniqueness results with easily verifiable criteria, tailored to the specific computations later on. The proofs are exercises in abstract deformation theory à la Deligne [22], and we postpone them to the end.

(3a) Let A be a graded algebra. Consider the set $\mathfrak{A}(A)$ of those A_∞-algebras $\mathcal{A}$ whose underlying graded vector space is A, and where the first two composition maps are

$$\mu^1_{\mathcal{A}} = 0, \quad \mu^2_{\mathcal{A}}(a_2, a_1) = (-1)^{|a_1|} a_2 a_1. \tag{3.1}$$

This means that the cohomology algebra is $H(\mathcal{A}) = A$ itself. $\mathcal{A}, \mathcal{A}' \in \mathfrak{A}(A)$ are considered equivalent if there is an A_∞-homomorphism $\mathcal{G} : \mathcal{A} \to \mathcal{A}'$ whose underlying linear map is $\mathcal{G}^1 = id_A$. After writing out the relevant equations, of which the first one is

$$\begin{aligned} \mu^3_{\mathcal{A}'}(a_3, a_2, a_1) &= \mu^3_{\mathcal{A}}(a_3, a_2, a_1) \\ &\quad + \mathcal{G}^2(a_3, \mu^2_{\mathcal{A}}(a_2, a_1)) + (-1)^{|a_1|-1}\mathcal{G}^2(\mu^2_{\mathcal{A}}(a_3, a_2), a_1) \\ &\quad - \mu^2_{\mathcal{A}}(a_3, \mathcal{G}^2(a_2, a_1)) - \mu^2_{\mathcal{A}}(\mathcal{G}^2(a_3, a_2), a_1), \end{aligned} \tag{3.2}$$

one sees that $\mathcal{A}$ and $\mathcal{G}$ determine $\mathcal{A}'$ completely, and that there are no constraints on $\mathcal{G}$. In other words, one can define a group $\mathfrak{G}(A)$ of *gauge transformations* which are arbitrary sequences of multilinear maps $\{\mathcal{G}^1 = id_A, \mathcal{G}^2 : A \otimes A \to A[-1], \dots\}$, and then (3.2) and its successors define an action of $\mathfrak{G}(A)$ on $\mathfrak{A}(A)$, whose quotient is the set of equivalence classes (this becomes entirely transparent in a geometric formulation, where it is the action of formal noncommutative diffeomorphisms on integrable odd vector fields). Additionally, there is a canonical action of the multiplicative semigroup $\mathbb{C}$ on $\mathfrak{A}(A)$, by multiplying μ^d with ϵ^{d-2}, which we denote by $\mathcal{A} \mapsto \epsilon^*\mathcal{A}$. For $\epsilon \neq 0$, it is still true that $\mathcal{A}$ is isomorphic to $\epsilon^*\mathcal{A}$, even though not by a gauge transformation: the isomorphism is multiplication by ϵ^k on the degree k component. The $\mathbb{C}$-action obviously descends to $\mathfrak{A}(A)/\mathfrak{G}(A)$.

The main justification for imposing conditions (3.1) is the Homological Perturbation Lemma, which says the following. Let $\mathcal{B}$ be an A_∞-algebra with an isomorphism $F : A \to H(\mathcal{B})$. Then one can find an A_∞-structure $\mathcal{A}$ on A satisfying (3.1) and an A_∞-homomorphism $\mathcal{F} : \mathcal{A} \to \mathcal{B}$ such that $H(\mathcal{F}) = F$. This establishes a bijection

$$\frac{\mathfrak{A}(A)}{\mathfrak{G}(A)} \cong \frac{\{\text{pairs } (\mathcal{B}, F : A \to H(\mathcal{B}))\}}{\{A_\infty\text{-quasi-isomorphisms compatible with } F\}}.$$

Remark 3.1. *The construction of $\mathcal{A}$ is explicit: take maps*

$$\pi : \mathcal{B} \to A, \quad \lambda : A \to \mathcal{B}, \quad h : \mathcal{B} \to \mathcal{B}[-1] \tag{3.3}$$

such that $\pi\mu^1_{\mathcal{B}} = 0$, $\mu^1_{\mathcal{B}}\lambda = 0$ (with the map induced by λ on cohomology being F), $\pi\lambda = id_A$, $\lambda\pi = id_{\mathcal{B}} + \mu^1_{\mathcal{B}}h + h\mu^1_{\mathcal{B}}$. Then $\mu^d_{\mathcal{A}}$ is a sum over planar trees with d leaves and one root. The contribution from each tree involves only the $\mu^k_{\mathcal{B}}$ for k at most d, and the auxiliary data (3.3) *(this particular formulation is taken from* [37]*). For instance, writing $b_k = \lambda(a_k)$ one has*

$$\begin{aligned} \mu^4_{\mathcal{A}}(a_4, a_3, a_2, a_1) &= \pi\mu^4_{\mathcal{B}}(b_4, b_3, b_2, b_1) + \\ &\pi\mu^2_{\mathcal{B}}(h\mu^3_{\mathcal{B}}(b_4, b_3, b_2), b_1) + \pi\mu^2_{\mathcal{B}}(b_4, h\mu^3_{\mathcal{B}}(b_3, b_2, b_1)) + \\ &\pi\mu^3_{\mathcal{B}}(h\mu^2_{\mathcal{B}}(b_4, b_3), b_2, b_1) + \pi\mu^3_{\mathcal{B}}(b_4, h\mu^2_{\mathcal{B}}(b_3, b_2), b_1) + \end{aligned} \tag{3.4}$$

$$\pi\mu^3_{\mathcal{B}}(b_4, b_3, h\mu^2_{\mathcal{B}}(b_2, b_1)) + \pi\mu^2_{\mathcal{B}}(h\mu^2_{\mathcal{B}}(b_4, b_3), h\mu^2_{\mathcal{B}}(b_2, b_1)) +$$
$$\pi\mu^2_{\mathcal{B}}(h\mu^2_{\mathcal{B}}(h\mu^2_{\mathcal{B}}(b_4, b_3), b_2), b_1) + \pi\mu^2_{\mathcal{B}}(h\mu^2_{\mathcal{B}}(b_4, h\mu^2_{\mathcal{B}}(b_3, b_2)), b_1) +$$
$$\pi\mu^2_{\mathcal{B}}(b_4, h\mu^2_{\mathcal{B}}(h\mu^2_{\mathcal{B}}(b_3, b_2), b_1)) + \pi\mu^2_{\mathcal{B}}(b_4, h\mu^2_{\mathcal{B}}(b_3, h\mu^2_{\mathcal{B}}(b_2, b_1))).$$

The Hochschild cohomology of A is a bigraded vector space. In our slightly unusual notation, elements of $HH^{s+t}(A,A)^t$ are represented by cocycles $\tau \in CC^{s+t}(A,A)^t$, which are multilinear maps $\tau : A^{\otimes s} \rightarrow A$ of degree t. Suppose that for $\mathcal{A} \in \mathfrak{A}(A)$, the compositions $\mu^s_{\mathcal{A}}$ vanish for $2 < s < d$. Then $\mu^d_{\mathcal{A}}$ is a Hochschild cocycle; its class

$$m^d_{\mathcal{A}} = [\mu^d_{\mathcal{A}}] \in HH^2(A,A)^{2-d},$$

the order d *deformation class* of $\mathcal{A}$, forms the obstruction to making $\mu^d_{\mathcal{A}}$ trivial by a gauge transformation which is itself equal to the identity to all orders $< d-1$. In particular, if all the obstruction groups $HH^2(A,A)^{2-s}$, $s > 2$, are zero, $\mathfrak{A}(A)/\mathfrak{G}(A)$ reduces to a point, which means that each A_∞-algebra structure on A can be trivialized by a gauge transformation (this is the *intrinsically formal* situation from [27, 32]). A slightly more general result allows the total obstruction group to be one-dimensional:

Lemma 3.2. *Suppose that there is some $d > 2$ such that*

$$\begin{cases} HH^2(A,A)^{2-d} = \mathbb{C}, \\ HH^2(A,A)^{2-s} = 0 \quad \text{for all } s > 2,\ s \neq d. \end{cases}$$

Suppose also that there exists an $\mathcal{A} \in \mathfrak{A}(A)$ such that $\mu^s_{\mathcal{A}} = 0$ for $2 < s < d$, and whose deformation class $m^d_{\mathcal{A}}$ is nonzero. Then any $\mathcal{A}' \in \mathfrak{A}(A)$ is equivalent to $\epsilon^*\mathcal{A}$ for some $\epsilon \in \mathbb{C}$, which means the $\mathbb{C}$-orbit of $[\mathcal{A}]$ is the whole of $\mathfrak{A}(A)/\mathfrak{G}(A)$.

We say that $\mathcal{A}$ is *versal*. All other problems that we will consider have the same structure: a set of equivalence classes described as a quotient, often involving the appeal to some kind of Perturbation Lemma; an additional group of reparametrizations acting on the quotient; a suitable version of Hochschild cohomology, and notion of deformation class; and ultimately, results expressing intrinsic formality or versality.

Remark 3.3. *Lemma 3.2 can be somewhat generalized, even though that does not fit in as well with our general philosophy as the original statement. The generalization goes as follows: suppose that $HH^2(A,A)^{2-s} = 0$ for all $s > 2$ except $s = d$, with $HH^2(A,A)^{2-d}$ arbitrary. Then every point in $\mathfrak{A}(A)/\mathfrak{G}(A)$ has a representative $\mathcal{A}$ with $\mu^s_{\mathcal{A}} = 0$ for $2 < s < d$; and if two such representatives $\mathcal{A}, \mathcal{A}'$ have the same order d deformation class, they are equivalent.*

(3b) Let $\mathcal{A}$ be an A_∞-algebra. A *one-parameter deformation* of $\mathcal{A}$ is a topological A_∞-algebra $\mathcal{A}_q$ over $\Lambda_{\mathbb{N}}$, whose underlying graded vector space is $\mathcal{A}\hat{\otimes}\Lambda_{\mathbb{N}}$, and whose composition maps reduce to those of $\mathcal{A}$ if one sets $q = 0$. More concretely,

$$\mu^d_{\mathcal{A}_q} = \mu^d_{\mathcal{A}} + q\mu^d_{\mathcal{A}_q,1} + q^2\mu^d_{\mathcal{A}_q,2} + \cdots = \mu^d_{\mathcal{A}} + O(q),$$

where each coefficient is a $\mathbb{C}$-linear map $\mathcal{A}^{\otimes d} \to \mathcal{A}[2-d]$. Two one-parameter deformations are equivalent if there is a $\Lambda_{\mathbb{N}}$-linear A_∞-homomorphism $\mathcal{G}_q : \mathcal{A}_q \to \mathcal{A}'_q$ between them of the form

$$\mathcal{G}^d_q = \begin{cases} id + O(q) & d = 1, \\ O(q) & d > 1. \end{cases} \tag{3.5}$$

Let $\mathfrak{A}_q(\mathcal{A})$ be the set of all one-parameter deformations, and $\mathfrak{G}_q(\mathcal{A})$ the group of sequences of multilinear maps $\{\mathcal{G}^1_q, \mathcal{G}^2_q, \dots\}$ on $\mathcal{A}\hat{\otimes}\Lambda_{\mathbb{N}}$ which satisfy (3.5). Then $\mathfrak{A}_q(\mathcal{A})/\mathfrak{G}_q(\mathcal{A})$ is the set of equivalence classes. The semigroup $End(\Lambda_{\mathbb{N}})$ acts on $\mathfrak{A}_q(\mathcal{A})$ by reparametrizations $q \mapsto \psi(q)$; we denote the action by $\mathcal{A}_q \mapsto \psi^*\mathcal{A}_q$. In this context, versality therefore means that the $End(\Lambda_{\mathbb{N}})$-orbit of some equivalence class $[\mathcal{A}_q]$ is the whole of $\mathfrak{A}_q(\mathcal{A})/\mathfrak{G}_q(\mathcal{A})$.

Remark 3.4. *We will also occasionally use terminology from finite order deformation theory. An order d infinitesimal deformation of $\mathcal{A}$ would be an A_∞-algebra $\mathcal{A}_q$ as before, except that the ground ring is now $\mathbb{C}[[q]]/q^{d+1}$. The notion of equivalence is obvious. Of course, a one-parameter deformation can be truncated to any desired finite order in q.*

Hochschild cohomology generalizes to A_∞-algebras [21] as a graded (no longer bigraded) space $HH^*(\mathcal{A}, \mathcal{A})$. Consider $CC^d(\mathcal{A}, \mathcal{A}) = \prod_{s+t=d} Hom_{\mathbb{C}}(\mathcal{A}^{\otimes s}, \mathcal{A}[t])$ with the Gerstenhaber product

$$(\sigma \circ \tau)^d(a_d, \dots, a_1) = \sum_{i,j} (-1)^{\sharp} \sigma^{d-j+1}(a_d, \dots, \tau^j(a_{i+j}, \dots, a_{i+1}), \dots, a_1), \tag{3.6}$$

where $\sharp = (|\tau| - 1)(|a_1| + \cdots + |a_i| - i)$, and the corresponding Lie bracket $[\sigma, \tau] = \sigma \circ \tau - (-1)^{(|\sigma|-1)(|\tau|-1)}\tau \circ \sigma$. As $\mu^*_{\mathcal{A}}$ itself lies in $CC^2(\mathcal{A}, \mathcal{A})$ and satisfies $\mu^*_{\mathcal{A}} \circ \mu^*_{\mathcal{A}} = 0$ by definition, one can define a differential $\delta\tau = [\mu^*_{\mathcal{A}}, \tau]$, and this is the A_∞ version of the Hochschild differential. Filtration by length yields a spectral sequence with

$$E_2^{s,t} = HH^{s+t}(A, A)^t$$

for $A = H(\mathcal{A})$, and this converges to $HH^*(\mathcal{A}, \mathcal{A})$ under suitable technical assumptions (for instance, if the nonzero terms in the E_2 plane are concentrated in a region cut out by $t \leq c_1$, $t \geq c_2 - c_3 s$ for c_1, c_2 arbitrary and $c_3 < 1$). If $\mu^d_{\mathcal{A}}$, $d \geq 3$, is the first nonzero higher order composition, then the first potentially nontrivial differential in the spectral sequence is

$$\delta_{d-1} = [m^d_{\mathcal{A}}, \cdot] : E_2^{s,t} \longrightarrow E_2^{s+d-1,t+2-d} \tag{3.7}$$

where $[\cdot, \cdot]$ is the Gerstenhaber bracket on $HH^{*+1}(A, A)$, and $m^d_{\mathcal{A}}$ is the order d deformation class. Actually, the spectral sequence is a quasi-isomorphism invariant, so that one can study it under the assumption (3.1). Then, either

$\mathcal{A}$ is equivalent to the trivial A_∞-structure on A and the spectral sequence degenerates; or else, after using gauge transformations to make as many higher order terms zero as possible, one encounters a nontrivial deformation class $m^d_{\mathcal{A}}$ for some $d > 2$. In that case, taking the Euler element $\tau \in HH^1(A,A)^0$ given by the derivation which is k times the identity on A^k, one finds that

$$[m^d_{\mathcal{A}}, \tau] = (d-2) m^d_{\mathcal{A}}. \tag{3.8}$$

By (3.7), δ_{d-1} is nonzero and kills $m^d_{\mathcal{A}} \in E^{d,2-d}_{d-1} = HH^2(A,A)^{2-d}$.

Having come this far by straightforward adaptation of the previously used notions, we encounter a slight snag, which is that the deformation problem governed by $HH^*(\mathcal{A},\mathcal{A})$ reaches beyond the class of A_∞-algebras into that of "curved" or "obstructed" ones. To exclude these additional deformations, one can proceed as follows. Let $\mathcal{A}$ be an A_∞-algebra with $\mu^1_{\mathcal{A}} = 0$. The *truncated Hochschild cohomology* is a graded vector space $HH^*(\mathcal{A},\mathcal{A})^{\leq 0}$, built from the subcomplex $CC^*(\mathcal{A},\mathcal{A})^{\leq 0}$ of Hochschild cochains $\tau : \mathcal{A}^{\otimes s} \rightarrow \mathcal{A}[t]$ with $t \leq 0$. There is a spectral sequence leading to it as before, where now

$$E_2^{s,t} = \begin{cases} HH^{s+t}(A,A)^t & t \leq 0, \\ 0 & \text{otherwise.} \end{cases} \tag{3.9}$$

All remarks made above carry over to this modification, including (3.8) because the Euler element lies in (3.9). For the application to deformation theory, consider the subset $\mathfrak{A}_q(\mathcal{A})^{\leq 0} \subset \mathfrak{A}_q(\mathcal{A})$ of those one-parameter deformations $\mathcal{A}_q$ during which the differential remains trivial, $\mu^1_{\mathcal{A}_q} = 0$. This still carries actions of $\mathfrak{G}_q(\mathcal{A})$ and of $End(\Lambda_{\mathbb{N}})$. The order q term of a deformation of this kind defines a primary deformation class

$$m^{\leq 0}_{\mathcal{A}_q} = [\mu^*_{\mathcal{A}_q,1}] \in HH^2(\mathcal{A},\mathcal{A})^{\leq 0},$$

which is the obstruction to transforming $\mathcal{A}_q$ into a deformation of order q^2 (or in other words, making the associated first order deformation trivial). The desired versality result is:

Lemma 3.5. *Suppose that $HH^2(\mathcal{A},\mathcal{A})^{\leq 0} \cong \mathbb{C}$. Let $\mathcal{A}_q$ be a one-parameter deformation with $\mu^1_{\mathcal{A}_q} = 0$, whose primary deformation class is nonzero. Then $\mathcal{A}_q$ is versal in $\mathfrak{A}_q(\mathcal{A})^{\leq 0}/\mathfrak{G}_q(\mathcal{A})$.*

Take an arbitrary A_∞-algebra $\mathcal{B}$, and use the Perturbation Lemma to find an $\mathcal{A}$ with $\mu^1_{\mathcal{A}} = 0$ and a quasi-isomorphism $\mathcal{F} : \mathcal{A} \rightarrow \mathcal{B}$. Suppose that $\mathcal{B}_q$ is a one-parameter deformation of $\mathcal{B}$ such that $H(\mathcal{B}_q)$ is torsion-free. Then $H(\mathcal{B}_q) \cong H(\mathcal{B}) \hat{\otimes} \Lambda_{\mathbb{N}}$ non-canonically, and a straightforward extension of the Perturbation Lemma allows one to construct a one-parameter deformation $\mathcal{A}_q$ of $\mathcal{A}$ with vanishing differential, and an $\mathcal{F}_q : \mathcal{A}_q \rightarrow \mathcal{B}_q$ which deforms $\mathcal{F}$. This in fact yields a bijection

$$\frac{\mathfrak{A}_q(\mathcal{A})^{\leq 0}}{\mathfrak{G}_q(\mathcal{A})} \cong \frac{\{\mathcal{B}_q \in \mathfrak{A}_q(\mathcal{B}), H(\mathcal{B}_q) \text{ torsion-free}\}}{\mathfrak{G}_q(\mathcal{B})}. \tag{3.10}$$

(3c) Lemma 3.2 is essentially sufficient for our needs, but Lemma 3.5 is not, since $HH^2(\mathcal{A},\mathcal{A})^{\leq 0}$ turns out to be too large in our example. This means that we need to restrict the class of allowed deformations further, so as to whittle down the dimension of the relevant "moduli space" to one. The solution adopted here (largely motivated by the desire to minimize work on the symplectic geometry side later on) is to consider A_∞-algebras equipped with an automorphism, and deformations of those. In fact, to turn this into a standard deformation problem, we replace the automorphism by its graph, which is an A_∞-bimodule.

Some background (see [66] for a more detailed exposition): to any A_∞-algebra $\mathcal{A}$ one can associate the dg category $\mathcal{A}$–$\mathcal{A}$ of A_∞-bimodules. An object is a graded vector space $\mathcal{M}$ with operations $\mu^{f,e}_{\mathcal{M}} : \mathcal{A}^{\otimes f} \otimes \mathcal{M} \otimes \mathcal{A}^{\otimes e} \to \mathcal{M}[1-e-f]$, for all $e, f \geq 0$, satisfying certain structural equation, and also subject to a certain c-unitality condition (see below). The cochain space of morphisms $\mathcal{M} \to \mathcal{N}$ of degree k is the space of sequences of multilinear maps $\beta^{f,e} : \mathcal{A}^{\otimes f} \otimes \mathcal{M} \otimes \mathcal{A}^{\otimes e} \to \mathcal{N}[k-e-f]$. Note that $\mu^{*,*}_{\mathcal{M}}$ itself is an element in $\mathit{hom}^1(\mathcal{M},\mathcal{M})$. To give explicit formulae for the dg structure, it is best to start with composition of morphisms, which is somewhat similar to the Gerstenhaber product:

$$\begin{aligned}(\gamma \circ \beta)^{f,e}&(b_f, \dots, b_1, m, a_e, \dots, b_1) = \\ &= \sum_{i,j} (-1)^{\dagger} \gamma^{f-j,i}(b_f, \dots, b_{j+1}, \beta^{j,e-i}(b_j, \dots, m, \dots, a_{i+1}), a_i, \dots, a_1),\end{aligned}$$

where $\dagger = |\beta|(|a_i| + \dots + |a_1| - i)$. We also have a preliminary version of the differential, involving only the A_∞-structure of $\mathcal{A}$:

$$\begin{aligned}(-1)^{|\beta|+1}&(\delta_{\mathrm{pre}}\beta)^{f,e}(b_f, \dots, b_1, m, a_e, \dots, a_1) = \\ &= \sum_{i,j} (-1)^{*} \beta^{f,e-j+1}(b_f, \dots, m, \dots, a_{i+j+1}, \mu^j_{\mathcal{A}}(a_{i+j}, \dots, a_{i+1}), a_i, \dots, a_1) \\ &\quad + \sum_{i,j} (-1)^{\natural} \beta^{f-j+1,e}(b_f, \dots, b_{i+j+1}, \mu^j_{\mathcal{A}}(b_{i+j}, \dots, b_{i+1}), b_i, \dots, m, \dots, a_1),\end{aligned}$$

with $*$ as in (2.2) and $\natural = |a_1| + \dots + |a_e| + |m| + |b_1| + \dots + |b_i| - e - 1 - i$. This satisfies $\delta^2_{\mathrm{pre}} = 0$ and $\delta_{\mathrm{pre}}(\gamma \circ \beta) = \delta_{\mathrm{pre}}(\gamma) \circ \beta + (-1)^{|\gamma|} \gamma \circ \delta_{\mathrm{pre}}(\beta)$. Having that, one can write the structural equations of an A_∞-bimodule as

$$\delta_{\mathrm{pre}}\mu^{*,*} + \mu^{*,*} \circ \mu^{*,*} = 0, \tag{3.11}$$

and the actual differential on $\mathit{hom}(\mathcal{M},\mathcal{N})$ as $\delta\beta = \delta_{\mathrm{pre}}\beta + \mu^{*,*}_{\mathcal{N}} \circ \beta - (-1)^{|\beta|} \beta \circ \mu^{*,*}_{\mathcal{M}}$. The cohomology $M = H(\mathcal{M}, \mu^{0,0}_{\mathcal{M}})$ of any A_∞-bimodule is naturally a graded bimodule over $A = H(\mathcal{A})$, and we require that the unit of A acts as the identity from both sides. By using the length filtration of $\mathit{hom}_{\mathcal{A}\text{–}\mathcal{A}}(\mathcal{M},\mathcal{N})$, one gets a spectral sequence whose E_1 term can be identified with $E_1^{s,t} = Hom_{A-A}(B^{-s}_{A-A}M, N[t])$, where $B^*_{A-A}M$ is the classical bimodule bar resolution. This is compatible with the differential, so that

$$E_2^{s,t} = Ext^s_{A-A}(M, N[t]). \tag{3.12}$$

In particular, if $\mathcal{M}$ or $\mathcal{N}$ are acyclic, then so is $hom_{\mathcal{A}\text{–}\mathcal{A}}(\mathcal{M}, \mathcal{N})$. If we allow ourselves to loosely borrow language from from [64], the fact that the bar resolution is built into the definition of our category ensures that "all A_∞-bimodules are K-projective (and K-injective)". It follows that any quasi-isomorphism $\mathcal{M} \to \mathcal{N}$ has a quasi-inverse, hence is an isomorphism in the category $H^0(\mathcal{A}\text{–}\mathcal{A})$.

There is a "coupled Hochschild cohomology" theory associated to a pair $(\mathcal{A}, \mathcal{M})$ consisting of an A_∞-algebra and an A_∞-bimodule over it. It sits in a long exact sequence

$$\begin{aligned} \cdots \longrightarrow HH^{*-1}(\mathcal{A}, \mathcal{A}) \xrightarrow{d} H^{*-1}(hom_{\mathcal{A}\text{–}\mathcal{A}}(\mathcal{M}, \mathcal{M})) \xrightarrow{a} \\ \longrightarrow HH^*(\mathcal{A} \,\&\, \mathcal{M}, \mathcal{A} \,\&\, \mathcal{M}) \xrightarrow{b} HH^*(\mathcal{A}, \mathcal{A}) \longrightarrow \cdots \end{aligned} \tag{3.13}$$

Later on, when we consider it from the point of view of deformation theory, the meaning of the sequence will be clear: b forgets $\mathcal{M}$ and concentrates on the deformation of $\mathcal{A}$, while the image of a consists of deformations of $\mathcal{M}$ with respect to a fixed $\mathcal{A}$. To define $HH^*(\mathcal{A} \,\&\, \mathcal{M}, \mathcal{A} \,\&\, \mathcal{M})$ one can use the triangular matrix trick, see for instance [25]. Define an A_∞-algebra

$$\mathcal{T} = \begin{pmatrix} \mathcal{T}_{11} & \mathcal{T}_{12} \\ \mathcal{T}_{21} & \mathcal{T}_{22} \end{pmatrix} = \begin{pmatrix} \mathcal{A} & 0 \\ \mathcal{M} & \mathcal{A} \end{pmatrix} \tag{3.14}$$

whose underlying graded vector space is the direct sum of two copies of our A_∞-algebra, $\mathcal{T}_{11} = \mathcal{T}_{22} = \mathcal{A}$, with one copy of the bimodule, $\mathcal{T}_{12} = \mathcal{M}$. The only nonzero compositions are

$$\begin{aligned} &\mu^d_{\mathcal{T}} = \mu^d_{\mathcal{A}} : \mathcal{T}_{kk}^{\otimes d} \longrightarrow \mathcal{T}_{kk}, \qquad k = 1, 2, \\ &\mu^{e+f+1}_{\mathcal{T}} = \mu^{f,e}_{\mathcal{M}} : \mathcal{T}_{22}^{\otimes f} \otimes \mathcal{T}_{21} \otimes \ \mathcal{T}_{11}^{\otimes e} \longrightarrow \mathcal{T}_{21}. \end{aligned}$$

Consider the subcomplex $CC^*(\mathcal{A} \,\&\, \mathcal{M}, \mathcal{A} \,\&\, \mathcal{M}) \subset CC^*(\mathcal{T}, \mathcal{T})$ of cochains of the same form, meaning that they are made up of maps $\mathcal{T}_{kk}^{\otimes d} \to \mathcal{T}_{kk}$, which should be the same for $k = 1, 2$, and maps $\mathcal{T}_{22}^{\otimes f} \otimes \mathcal{T}_{21} \otimes \mathcal{T}_{11}^{\otimes e} \to \mathcal{T}_{21}$. The cochains of the second kind form a subcomplex of $CC^*(\mathcal{A} \,\&\, \mathcal{M}, \mathcal{A} \,\&\, \mathcal{M})$ which is isomorphic to $hom(\mathcal{M}, \mathcal{M})[-1]$, and the quotient complex is just $CC^*(\mathcal{A}, \mathcal{A})$, which makes (3.13) clear.

The bimodules that we will be interested in are graphs $\mathcal{M} = Graph(\mathcal{F})$ of A_∞-homomorphisms $\mathcal{F} : \mathcal{A} \to \mathcal{A}$. The underlying graded vector space of $\mathcal{M}$ is the same as for $\mathcal{A}$, and the operations are

$$\begin{aligned}&\mu_{\mathcal{M}}^{f,e}(b_f,\dots,b_1,m,a_e,\dots,a_1) = \\ &= \sum_{i_1+\dots+i_r=f} \mu_{\mathcal{A}}^{f+r+1}(b_f,\dots,b_1,m,\mathcal{F}^{i_r}(a_e,\dots,a_{e-i_r+1}),\dots,\mathcal{F}^{i_1}(a_{i_1},\dots,a_1)).\end{aligned}$$

LEMMA 3.6. *For a quasi-isomorphism $\mathcal{F}$, one can map $H^*(hom_{\mathcal{A}\text{-}\mathcal{A}}(\mathcal{M},\mathcal{M})) \cong HH^*(\mathcal{A},\mathcal{A})$ in such a way that the differential d in* (3.13) *becomes $id-\mathcal{F}^*$, where $\mathcal{F}^*$ is the automorphism of Hochschild cohomology induced by $\mathcal{F}$.*

Our proof of this will be rather pedestrian, but a more satisfactory treatment requires all the machinery of A_∞-categories of A_∞-functors, for which we have no real use elsewhere. To get started, we need to recall how quasi-isomorphism invariance of Hochschild cohomology for A_∞-algebras is proved. The most obvious scheme does not apply because HH^* is neither co- nor contravariantly functorial, and instead one passes through a generalization, which is the *functor Hochschild cohomology* $HH^*(\mathcal{F},\mathcal{F})$ of an A_∞-homomorphism $\mathcal{F}:\mathcal{A}\to\mathcal{B}$. We will not give the full definition of this, but the underlying cochain complex is $CC^d(\mathcal{F},\mathcal{F}) = \prod_{s+t=d} Hom(\mathcal{A}^{\otimes s},\mathcal{B}[t])$, and the E_2 term of the filtration-by-length spectral sequence is $Ext^s_{A-A}(A,B[t])$, where B is considered as an A-bimodule through pullback by $F = H(\mathcal{F})$. There are natural maps $\mathcal{F}^{*,\mathrm{left}} : HH^*(\mathcal{A},\mathcal{A}) \to HH^*(\mathcal{F},\mathcal{F})$, $\mathcal{F}^{*,\mathrm{right}} : HH^*(\mathcal{B},\mathcal{B}) \to HH^*(\mathcal{F},\mathcal{F})$ defined on the cochain level by

$$\begin{aligned}&(\mathcal{F}^{*,\mathrm{left}}\tau)(a_d,\dots,a_1) = \sum_{i,j}(-1)^{\sharp}\mathcal{F}^{d-j+1}(a_d,\dots,\tau^j(a_{i+j},\dots,a_{i+1}),\dots,a_1),\\ &(\mathcal{F}^{*,\mathrm{right}}\tau)(a_d,\dots,a_1) = \\ &\qquad\qquad = \sum_{i_1+\dots+i_r=d}\tau^r(\mathcal{F}^{i_r}(a_d,\dots,a_{d-i_r+1}),\dots,\mathcal{F}^{i_1}(a_{i_1},\dots,a_1)).\end{aligned}$$

with $\sharp$ as in (3.6). If $\mathcal{F}$ is a quasi-isomorphism, a spectral sequence comparison argument shows that both maps are isomorphisms. One then defines $\mathcal{F}^*$ to be the composition $(\mathcal{F}^{*,\mathrm{left}})^{-1}\circ\mathcal{F}^{*,\mathrm{right}} : HH^*(\mathcal{B},\mathcal{B}) \to HH^*(\mathcal{A},\mathcal{A})$.

Suppose from now on that $\mathcal{B} = \mathcal{A}$, so we are in the situation of Lemma 3.6, and $\mathcal{F}^*$ is the automorphism of Hochschild cohomology induced by $\mathcal{F}$. There is a natural isomorphism $c : HH^*(\mathcal{F},\mathcal{F}) \to H^*(hom_{\mathcal{A}\text{-}\mathcal{A}}(\mathcal{M},\mathcal{M}))$ for $\mathcal{M} = Graph(\mathcal{F})$, given on cochains by

$$\begin{aligned}&(c\tau)^{f,e}(b_f,\dots,b_1,m,a_e,\dots,a_1) = \\ &\quad = \sum_{\substack{k\leq r\\ i_1+\dots+i_r=e}} (-1)^{\flat}\mu_{\mathcal{A}}^{f+1+r}(b_f,\dots,b_1,m,\mathcal{F}^{i_r}(a_e,\dots,a_{e-i_r+1}),\dots,\\ &\qquad\qquad \tau^{i_k}(a_{i_1+\dots+i_k},\dots,a_{i_1+\dots+i_{k-1}+1}),,\dots,\mathcal{F}^{i_1}(a_{i_1},\dots,a_1)).\end{aligned}$$

$\flat = (|\tau|-1)(|a_1|+\dots+|a_{i_{k-1}}|-i_1-\dots-i_{k-1})$. To prove that this is an isomorphism, one considers the E_2 term of the usual spectral sequences, and checks that the action of c on it can be traced back to a comparison map between two different resolutions of the bimodule $M = H(\mathcal{M})$. Finally, the following diagram, where d is from (3.13), commutes:

$$\begin{tikzcd} HH^*(\mathcal{A},\mathcal{A}) \arrow[r, "\mathcal{F}^{*,\mathrm{left}} - \mathcal{F}^{*,\mathrm{right}}"] \arrow[dr, bend right=15, "d"'] & HH^*(\mathcal{F},\mathcal{F}) \arrow[d, "c", "\cong"'] \\ & H^*(hom_{\mathcal{A}\text{-}\mathcal{A}}(\mathcal{M},\mathcal{M})) \end{tikzcd} \tag{3.15}$$

In view of the definition of $\mathcal{F}^*$, this proves Lemma 3.6. Actually (3.15) does not strictly commute on cochains, but the necessary homotopy can be easily provided,

$$(h\tau)(b_f,\dots,m,\dots,a_1) = \\ = \sum_{i_1+\dots+i_r=e} \tau(b_f,\dots,m,\mathcal{F}^{i_r}(a_e,\dots,a_{e-i_r+1}),\dots,\mathcal{F}^{i_1}(a_{i_1},\dots,a_1)).$$

(3d) Take a graded algebra A, and let $\mathcal{A}$ be an A_∞-structure on it as in (3.1). Given a graded A-bimodule M, we want to look at A_∞-bimodule structures $\mathcal{M}$ on it such that the analogue of (3.1) holds,

$$\mu^{0,0}_{\mathcal{M}} = 0, \quad \mu^{1,0}_{\mathcal{M}}(a,m) = (-1)^{|a|}ma, \quad \mu^{0,1}_{\mathcal{M}}(m,a) = (-1)^{|m|}am. \tag{3.16}$$

Lemma 3.7. *Suppose that $Ext^k_{A\otimes A^{op}}(M, M[1-k]) = 0$ for all $k \geq 2$. Then, up to isomorphism, there is at most one $\mathcal{M}$ satisfying* (3.16).

It might seem that there is a straightforward "linear" approach to this: take two A_∞-bimodules $\mathcal{M},\mathcal{M}'$ satisfying (3.16), and look at the spectral sequence (3.12) associated to $hom(\mathcal{M},\mathcal{M}')$. The assumption in the Lemma ensures that $id_M \in E_2^{0,0}$ survives to E_∞, and if the spectral sequence converges, one could use a standard comparison argument to show that the resulting element of $H^0(hom(\mathcal{M},\mathcal{M}'))$ is an isomorphism. However, there seems to be no reason why convergence should hold in general, and our proof will rely instead on the "nonlinear" methods of deformation theory. To complete our discussion, it is maybe useful to say that there is a version of the Perturbation Lemma for A_∞-bimodules, which shows that any such bimodule over an A_∞-algebra $\mathcal{A}$ with $\mu^1_{\mathcal{A}} = 0$ is quasi-isomorphic to one with $\mu^{0,0}_{\mathcal{M}} = 0$, hence which is of the form (3.16). To put this on a more formal footing, consider the dg subcategory $\mathcal{A}\text{–}\mathcal{A}^{\leq 0} \subset \mathcal{A}\text{–}\mathcal{A}$ whose objects are A_∞-bimodules $\mathcal{M}$ with $\mu^{0,0}_{\mathcal{M}} = 0$, and where the morphisms β of degree d must satisfy $\beta^{p,q} = 0$ for $p+q < d$. By definition, the functor induced by inclusion is a full embedding $H^0(\mathcal{A}\text{–}\mathcal{A}^{\leq 0}) \to H^0(\mathcal{A}\text{–}\mathcal{A})$. The bimodule Perturbation Lemma shows that it is an equivalence.

We now turn to the "coupled" deformation problem which is our main interest in this matter. Let A, $\mathcal{A}$ and M be as before, and suppose that we have some fixed A_∞-bimodule structure $\mathcal{M}$ satisfying (3.16). Let $\mathfrak{A}_q(\mathcal{A}\,\&\,\mathcal{M})^{\leq 0}$ be the set of pairs $(\mathcal{A}_q, \mathcal{M}_q)$ where $\mathcal{A}_q$ is a one-parameter deformation of $\mathcal{A}$, and $\mathcal{M}_q$ a corresponding one-parameter deformation of $\mathcal{M}$ (that is, an $\mathcal{A}_q$-bimodule structure on $\mathcal{M}\hat{\otimes}_{\mathbb{C}}\Lambda_{\mathbb{N}}$ having the obvious kind of q-expansion), such that

$$\mu^1_{\mathcal{A}_q} = 0, \quad \mu^{0,0}_{\mathcal{M}_q} = 0. \tag{3.17}$$

The formal gauge group $\mathfrak{G}_q(\mathcal{A}\,\&\,\mathcal{M})$ for this problem can be defined as a suitable subgroup of the corresponding group for deformations of the triangular algebra $\mathcal{T}$. For us, what matters is that gauge equivalence of pairs implies equivalence of the A_∞-deformations $\mathcal{A}_q$, and we will later state the versality result in a correspondingly weak form. Let $HH^*(\mathcal{A}\,\&\,\mathcal{M}, \mathcal{A}\,\&\,\mathcal{M})^{\leq 0}$ be the truncated version of coupled Hochschild cohomology, defined by looking only at cochains $\tau_{11} = \tau_{22} : \mathcal{A}^{\otimes s} \rightarrow \mathcal{A}[t]$, $\tau_{12} : \mathcal{A}^{\otimes f} \otimes \mathcal{M} \otimes \mathcal{A}^{\otimes e} \rightarrow \mathcal{M}[t]$ with $t \leq 0$. This sits in a suitable version of (3.13),

$$\begin{aligned} \dots \longrightarrow HH^{*-1}(\mathcal{A},\mathcal{A})^{\leq 0} \xrightarrow{d} H^{*-1}(hom_{\mathcal{A}\text{–}\mathcal{A}^{\leq 0}}(\mathcal{M},\mathcal{M})) \longrightarrow \\ \rightarrow HH^*(\mathcal{A}\,\&\,\mathcal{M}, \mathcal{A}\,\&\,\mathcal{M})^{\leq 0} \longrightarrow HH^*(\mathcal{A},\mathcal{A})^{\leq 0} \xrightarrow{d} \dots \end{aligned} \tag{3.18}$$

In the case where $\mathcal{M} = Graph(\mathcal{F})$, the obvious variation of Lemma 3.6 applies. Any pair $(\mathcal{A}_q, \mathcal{M}_q) \in \mathfrak{A}_q(\mathcal{A}\,\&\,\mathcal{M})^{\leq 0}$ has a primary deformation class in truncated coupled Hochschild cohomology. Again, the important thing for us is that this maps to the deformation class of $\mathcal{A}_q$ under (3.18).

Lemma 3.8. *Suppose that* $HH^2(\mathcal{A}\,\&\,\mathcal{M}, \mathcal{A}\,\&\,\mathcal{M})^{\leq 0} \cong \mathbb{C}$*, and that there is a deformation* $(\mathcal{A}_q, \mathcal{M}_q)$ *satisfying* (3.17) *with nonzero* $m^{\leq 0}_{\mathcal{A}_q} \in HH^2(\mathcal{A},\mathcal{A})^{\leq 0}$*. Then, for any other pair* $(\mathcal{A}'_q, \mathcal{M}'_q) \in \mathfrak{A}_q(\mathcal{A}\,\&\,\mathcal{M})^{\leq 0}$*, there is a* $\psi \in End(\Lambda_{\mathbb{N}})$ *such that* $\mathcal{A}'_q \cong \psi^*\mathcal{A}_q$ *as a one-parameter deformation of* A_∞*-algebras.*

(3e) The framework of abstract one-parameter deformation theory is as follows. Let $\mathfrak{g}$ be a dg Lie algebra. We consider the space $\mathfrak{A}_q(\mathfrak{g})$ of elements $\alpha_q \in \mathfrak{g}^1 \hat{\otimes}\, q\Lambda_{\mathbb{N}}$ which satisfy the Maurer-Cartan equation

$$\delta\alpha_q + \tfrac{1}{2}[\alpha_q, \alpha_q] = 0. \tag{3.19}$$

In particular, if we expand $\alpha_q = \alpha_{q,1}q + \alpha_{q,2}q^2 + \cdots$, the leading order term defines a primary deformation class $[\alpha_{q,1}] \in H^1(\mathfrak{g})$. An infinitesimal abstract gauge transformation $\gamma_q \in \mathfrak{g}^0\hat{\otimes} q\Lambda_{\mathbb{N}}$ defines a formal vector field X_q on $\mathfrak{A}_q(\mathfrak{g})$, $X_q(\alpha_q) = \delta\gamma_q + [\alpha_q, \gamma_q]$. These vector fields exponentiate to an action of a group $\mathfrak{G}_q(\mathfrak{g})$ on $\mathfrak{A}_q(\mathfrak{g})$, which we denote by $\exp(\gamma_q)(\alpha_q)$. The primary deformation class is an invariant of this action. As usual, there is also an action of $End(\Lambda_{\mathbb{N}})$ on $\mathfrak{A}_q(\mathfrak{g})$ by reparametrizations, hence a notion of versality. The abstract formality and versality theorems are

Lemma 3.9. *If* $H^1(\mathfrak{g}) = 0$*,* $\mathfrak{A}_q(\mathfrak{g})/\mathfrak{G}_q(\mathfrak{g})$ *is a point.*

Lemma 3.10. *Suppose that $H^1(\mathfrak{g}) \cong \mathbb{C}$. Then, every $\alpha_q \in \mathfrak{A}_q(\mathfrak{g})$ whose primary deformation class is nonzero is versal.*

These are both standard results, but for the convenience of the reader we reproduce the conventional induction-by-order proof of the second one. Take an arbitrary $\beta_q \in \mathfrak{A}_q(\mathfrak{g})$. Suppose that for some $d \geq 1$, we have $\psi^d \in End(\Lambda_{\mathbb{N}})$ and $\beta_q^d \in \mathfrak{A}_q(\mathfrak{g})$ gauge equivalent to β_q, such that

$$\beta_q^d - (\psi^d)^* \alpha_q \stackrel{\text{def}}{=} \epsilon_q \in O(q^d). \tag{3.20}$$

Because $\delta\epsilon_q = \frac{1}{2}[(\psi^d)^*\alpha_q + \beta_q^d, \epsilon_q] \in O(q^{d+1})$, the leading order term $\epsilon_{q,d} \in \mathfrak{g}^1$ is closed. By assumption, one can therefore write $\beta_q^d = (\psi^d)^*\alpha_q + \epsilon_q = (\psi^d)^*\alpha_q + cq^d\alpha^{d,1} + q^d\delta b + O(q^{d+1})$ for some $c \in \mathbb{C}$, $b \in \mathfrak{g}^0$. With

$$\psi^{d+1}(q) = \psi^d(q) + cq^d, \quad \beta_q^{d+1} = \exp(-q^d b)(\beta_q^d) \tag{3.21}$$

one finds that (3.20) holds for $d+1$ as well. Since the parameter change and gauge transformation in (3.21) are equal to the identity up to $O(q^d)$, the process converges to $(\psi^\infty)^*\alpha_q = \beta_q^\infty$, with β_q^∞ gauge equivalent to β_q.

When applied to $\mathfrak{g}_{\mathcal{A}} = CC^{*+1}(\mathcal{A}, \mathcal{A})^{\leq 0}$, this yields Lemma 3.3. Similarly, for the coupled case one sets $\mathfrak{g}_{\mathcal{A}\,\&\,\mathcal{M}} = CC^{*+1}(\mathcal{A}\,\&\,\mathcal{M}, \mathcal{A}\,\&\,\mathcal{M})^{\leq 0}$ and obtains Lemma 3.8. The other results stated above, which lack the parameter q, require modifications of the basic framework. Let A be a graded algebra, and $\mathfrak{g}_A = CC^{*+1}(A, A)$ its shifted Hochschild complex with the Gerstenhaber bracket. One can reformulate the bigrading of Hochschild cohomology as a $\mathbb{C}^*$-action on $\mathfrak{g}_A$, which has weight t on $CC^{s+t}(A, A)^t$. We combine this with the $\mathbb{C}^*$-action on $\Lambda_{\mathbb{N}}$ which has weight d on $\mathbb{C}q^d$, and obtain an action on $\mathfrak{g}_A \otimes q\Lambda_{\mathbb{N}}$. Given an A_∞-structure $\mathcal{A}$ on A satisfying (3.1), one can define a one-parameter deformation $\mathcal{A}_q$ of the trivial A_∞-structure by $\mu^d_{\mathcal{A}_q} = q^{d-2}\mu^d_{\mathcal{A}}$, and this is given by a $\mathbb{C}^*$-invariant element of $\mathfrak{g}_A \otimes q\Lambda_{\mathbb{N}}$. The only possible changes of parameter in this context are the homogeneous ones $q \mapsto \epsilon q$, $\epsilon \in \mathbb{C}^*$, and with that in mind the proof of Lemma 3.2 is by a straightforward adaptation of Lemma 3.10. In contrast, Lemma 3.5 requires a more substantial tweaking of the theory, where the basic object one considers is a dg Lie algebra $\mathfrak{g}$ with a complete decreasing filtration $F_0\mathfrak{g} = \mathfrak{g} \supset F_1\mathfrak{g} \supset \dots$ such that $\delta(F_i\mathfrak{g}) \subset F_{i+1}\mathfrak{g}$ and $[F_j\mathfrak{g}, F_k\mathfrak{g}] \subset F_{j+k}\mathfrak{g}$. Take a solution $\alpha \in (F_1\mathfrak{g})^1$ of (3.19), and consider the corresponding deformed differential

$$\delta_\alpha = \delta + [\alpha, \cdot].$$

There is a series of obstruction groups, which are the terms $E_2^{d,1-d}$, $d \geq 2$, in the spectral sequence leading to the cohomology of $(\mathfrak{g}, \delta_\alpha)$. A slightly refined version of the computation in the proof of Lemma 3.10 shows that if all of them vanish, then any other solution of (3.19) with $\beta - \alpha \in F_2\mathfrak{g}$ is equivalent to α by a gauge transformation, obtained by exponentiating the action of $(F_1\mathfrak{g})^0$. For our purpose, one applies this to $\mathfrak{g} = \bigoplus_{e,f} Hom^*(A^{\otimes e} \otimes M \otimes A^{\otimes f}, A)$, with δ and the bracket as suggested by (3.11), and $F_i\mathfrak{g}$ the filtration by length $i = e + f$. The obstruction groups are precisely those which occur in Lemma 3.5.

(3f) The previous discussion generalizes effortlessly to A_∞-categories. For didactic reasons, start with an ordinary category C having finitely many objects $X_1, \dots, X_m$, and let A be its total morphism algebra, which is linear over R_m. Consider Hochschild cochains that preserve this structure, which means that they are R_m-bimodule maps $A \otimes_{R_m} \cdots \otimes_{R_m} A \to A$. By writing this out, one sees that the relevant complex can be generalized to arbitrary categories, leading to a Hochschild cohomology theory $HH^*(C, C)$. The same applies to Hochschild cohomology of A_∞-categories, and all the other notions of deformation theory.

Remark 3.11. *For any category C with finitely many objects, the Hochschild cohomology $HH^*(C, C)$ is the same as $HH^*(A, A)$ of the total morphism algebra. In other words, the inclusion of subcomplex of R_m-bilinear Hochschild cochains into the whole cochain complex is a quasi-isomorphism. The reason is that the relative bar resolution*

$$\cdots \longrightarrow A \otimes_{R_m} A \otimes_{R_m} \otimes A \longrightarrow A \otimes_{R_m} A \longrightarrow A$$

is also a resolution by projective A-bimodules in the non-relative sense. By quasi-isomorphism invariance of abstract deformation theories, this shows that any A_∞-structure $\mathcal{A}$ on A satisfying (3.1) comes, up to gauge equivalence, from an A_∞-category structure $\mathcal{C}$ on C. The previous observation combined with the standard spectral sequence argument also proves that for any finite A_∞-category $\mathcal{C}$ with total morphism A_∞-algebra $\mathcal{A}$, $HH^*(\mathcal{C}, \mathcal{C}) \cong HH^*(\mathcal{A}, \mathcal{A})$, and similarly for the truncated version. This implies that the one-parameter deformation theory of $\mathcal{A}$ as an A_∞-algebra is the same as that of $\mathcal{C}$ as an A_∞-category. On occasion, these facts will allow us to be somewhat sloppy about the distinction between finite categories and their total morphism algebras.

To make the connection with Section 2, we should relate the deformation theory of $\mathcal{C}$ with that of its derived categories. However, this cannot be done in a truly satisfactory way within the restrictions of the present paper, because we have not introduced an A_∞-category underlying $D^\pi(\mathcal{C})$, and so we will limit ourselves to a single rudimentary statement, Let $\mathcal{C}_q$ be a one-parameter deformation of $\mathcal{C}$. Suppose that we have a finite set $(X_1, \dots, X_m)$ of objects of $\mathcal{C}$ which are split-generators for $D^\pi(\mathcal{C})$. Let $\mathcal{A}$ be their total morphism A_∞-algebra, and $\mathcal{A}_q$ its one-parameter deformation induced from $\mathcal{C}_q$.

Lemma 3.12. *Assume that there are $Y_0, Y_1 \in Ob\,\mathcal{C}$ such that $H(hom(Y_0, Y_1))$ changes at first order in the deformation. More precisely, this means that the natural map*

$$H(hom_{\mathcal{C}_q}(Y_0, Y_1) \otimes_{\Lambda_\mathbb{N}} \mathbb{C}[q]/q^2, \mu^1_{\mathcal{C}_q}) \otimes_{\mathbb{C}[q]/q^2} \mathbb{C} \longrightarrow H(hom_{\mathcal{C}}(Y_0, Y_1), \mu^1_{\mathcal{C}}) \tag{3.22}$$

is not an isomorphism. Then $\mathcal{A}_q$ is nontrivial at first order in q.

Proof. Suppose that the statement is false. After an R_m-linear gauge transformation, we can assume that $\mathcal{A}_q$ is a deformation of order q^2. Then the same holds for the induced one-parameter deformation $Tw\mathcal{A}_q$ of $Tw\mathcal{A}$. But the natural A_∞-functor $\mathcal{A}_q \to \mathcal{C}_q$ extends to a $\Lambda_{\mathbb{N}}$-linear A_∞-functor $Tw\mathcal{F}_q : Tw\mathcal{A}_q \to Tw\mathcal{C}_q$, which is full and faithful on cohomology. This means that (3.22) must be an isomorphism for all objects Y_0, Y_1 which are twisted complexes constructed from the X_k. Since that property carries over to direct summands, we have a contradiction. □

4. Group actions

Semidirect products $\Lambda V \rtimes \Gamma$ of an exterior algebra with a finite abelian group appear naturally in the context of homological mirror symmetry for hypersurfaces in projective spaces. This is a consequence of Serre duality and Beilinson's description of $D^bCoh(\mathbb{CP}^n)$ [7]. The Hochschild cohomology of such algebras can be computed by a version of the classical Hochschild-Kostant-Rosenberg (HKR) theorem [29], and this will later form the basis for applying the deformation theory from the previous section. Besides proving that theorem, we also use the opportunity to assemble some elementary facts about semidirect products, both of algebras and A_∞-algebras. The exposition is by no means as general as possible, in particular, we shamelessly restrict ourselves to abelian groups whenever this makes things simpler. Dissatisfied readers may want to turn to [44] and references therein.

(4a) Let A be a graded algebra carrying an action ρ of a finite group Γ. The semidirect product $A \rtimes \Gamma$ is the tensor product $A \otimes \mathbb{C}\Gamma$ with the twisted multiplication

$$(a_2 \otimes \gamma_2)(a_1 \otimes \gamma_1) = a_2\rho(\gamma_2)(a_1) \otimes \gamma_2\gamma_1.$$

The purpose of this construction is to turn the automorphisms $\rho(\Gamma)$ into inner ones: in fact, $(1\otimes\gamma)(a\otimes e)(1\otimes\gamma)^{-1} = \rho(\gamma)(a)\otimes e$. Therefore, it is not surprising that $A\rtimes\Gamma$ depends only on the "outer part" of the group action, in the following sense: suppose that $\tilde\rho$ is another action related to the previous one by

$$\tilde\rho(\gamma) = \iota(\gamma)\rho(\gamma)\iota(\gamma)^{-1}, \tag{4.1}$$

where $\iota : \Gamma \to (A^0)^\times$ is a map satisfying $\iota(\gamma_2\gamma_1) = \iota(\gamma_2)\rho(\gamma_2)(\iota(\gamma_1))$. Then $A\rtimes_\rho\Gamma$ and $A \rtimes_{\tilde\rho} \Gamma$ are isomorphic by $a\otimes\gamma \mapsto a\iota(\gamma)^{-1}\otimes\gamma$. This kind of consideration appears naturally in connection with the following construction: suppose that there is a normal subgroup $\Gamma_0 \subset \Gamma$ such that the projection $\Gamma \to \Gamma/\Gamma_0$ admits a right inverse Σ. Then $\Gamma \cong \Gamma_0\rtimes(\Gamma/\Gamma_0)$, and there is a corresponding isomorphism

$$A \rtimes \Gamma \cong (A \rtimes \Gamma_0) \rtimes (\Gamma/\Gamma_0). \tag{4.2}$$

More precisely, let ρ_Γ be the given Γ-action, and ρ_{Γ_0} its restriction. Through the splitting Σ we have an action ρ_{Γ/Γ_0} of the quotient on $A \rtimes \Gamma_0$:

$$\rho_{\Gamma/\Gamma_0}(\bar\gamma)(a \otimes \gamma_0) = \rho_\Gamma(\Sigma(\bar\gamma))(a) \otimes \Sigma(\bar\gamma)\gamma_0\Sigma(\bar\gamma)^{-1}, \tag{4.3}$$

and these are the three actions involved in (4.2). Different choices of Σ lead to actions (4.3) which differ by inner automorphisms as in (4.1), and the discussion there shows that the right hand side of (4.2) is independent of the choice, as indeed it ought to be.

Any semidirect product $A \rtimes \Gamma$ carries a natural action ρ_{Γ^*} of the group of scalar characters $\Gamma^* = Hom(\Gamma, \mathbb{C}^*)$; $\rho_{\Gamma^*}(\chi)(a \otimes g) = a\chi(g) \otimes g$. Additional automorphisms can be produced from those of A. We have already seen an instance of this in (4.3), and the general construction goes as follows: let U be an automorphism of A, and suppose that there is an automorphism ξ of Γ as well as a map $\iota : \Gamma \to (A^0)^\times$ satisfying $\iota(\gamma_2\gamma_1) = \rho(\xi(\gamma_2))(\iota(\gamma_2)) \cdot \iota(\gamma_1)$, such that

$$\rho(\xi(\gamma))(U(a)) = \iota(\gamma) U(\rho(\gamma)(a)) \iota(\gamma)^{-1} \tag{4.4}$$

holds. Informally, this means that U commutes with the group actions in a twisted sense given by ξ, and up to inner automorphism which are conjugations with the ι's. Then one can define an induced automorphism of $A \rtimes \Gamma$ by

$$(U \rtimes \xi)(a \otimes \gamma) = U(a)\iota(\gamma)^{-1} \otimes \xi(\gamma). \tag{4.5}$$

Of course, ι is not unique: for instance, passing from $\iota(\gamma)$ to $\iota(\gamma)\chi(\gamma)$ for some $\chi \in \Gamma^*$ amounts to composing $U \rtimes \xi$ with $\rho_{\Gamma^*}(\chi)$.

In a slightly different direction, note that $A \rtimes \Gamma$ is linear over the semisimple ring $\mathbb{C}\Gamma$. For simplicity, assume that Γ is abelian, so that $\mathbb{C}\Gamma \cong R_{|\Gamma|}$ has a basis of orthogonal idempotents, which are the normalized characters $u_\chi = |G|^{-1}\chi$. This means that we can consider $A \rtimes \Gamma$ as a category with objects χ and morphism spaces

$$Hom(\chi_1, \chi_2) = u_{\chi_2}(A \rtimes \Gamma)u_{\chi_1} = A^{\chi_1/\chi_2} \otimes \mathbb{C}u_{\chi_1}.$$

where χ_2/χ_1 is the (pointwise) division in Γ^*, and A^χ is the subspace on which G acts as on the representation corresponding to χ. This suggests an obvious generalization, which is to consider an action of an abelian group Γ on a linear graded category C; here and in related discussions below, the standing assumption is that the action on the set of objects is trivial. Then, the semidirect product category $C \rtimes \Gamma$ has objects $(X, \chi) \in Ob\, C \times \Gamma^*$, and morphisms $Hom((X_1, \chi_1), (X_2, \chi_2)) = Hom(X_1, X_2)^{\chi_1/\chi_2}$. By construction, if C has only finitely many objects and A it its total morphism algebra, then $A \rtimes \Gamma$ is the total morphism algebra of $C \rtimes \Gamma$.

(4b) A Γ-action on an A_∞-algebra $\mathcal{A}$ is a linear action on A, such that all the $\mu^d_{\mathcal{A}}$ are equivariant. The semidirect product A_∞-algebra $\mathcal{A} \rtimes \Gamma$ is then defined as $\mathcal{A} \otimes \mathbb{C}\Gamma$ with the compositions

$$\begin{aligned} &\mu^d_{\mathcal{A} \rtimes \Gamma}(a_d \otimes \gamma_d, \dots, a_1 \otimes \gamma_1) = \\ &\qquad = \mu^d_{\mathcal{A}}(a_d, \rho(\gamma_d)(a_{d-1}), \rho(\gamma_d\gamma_{d_1})(a_{d-2}), \dots) \otimes \gamma_d \cdots \gamma_1. \end{aligned}$$

This generalizes to A_∞-categories with Γ-actions; if there are only finitely many objects, these are the same as R_m-linear Γ-actions on R_m-linear A_∞-algebras.

Most of the discussion above applies to the A_∞-case as well, the only problematic parts being those involving invertible elements and the associated inner automorphisms, since that notion does not carry over in a simple way. In the R_m-linear situation there is at least a rudimentary substitute. Namely, for $r = (r_1, \dots, r_m) \in I = (\mathbb{C}^*)^m$ and $a \in u_j A u_k$, define $r \cdot a = r_j a$ and $a \cdot r = a r_k$. In a slight abuse of terminology, we call $a \mapsto r \cdot a \cdot r^{-1}$ an *elementary inner automorphism*. One can use these instead of actual xax^{-1}'s in formulae like (4.1), (4.4), and this gives a valid A_∞-version of the relevant observations for algebras, even though it is so reduced in scope as to be basically trivial.

Given an A_∞-category $\mathcal{C}$ with a Γ-action, one can define the A_∞-category $Tw_\Gamma \mathcal{C}$ of equivariant twisted complexes. This also carries a Γ-action, and if we forget that action, it becomes equivalent to a full A_∞-subcategory of the ordinary $Tw\,\mathcal{C}$. The construction requires the following changes in the two basic steps: (1) the equivariant version $\Sigma_\Gamma \mathcal{C}$ of the additive completion has objects which are formal direct sums

$$\bigoplus_{f \in F} (X_f \otimes V_f)[\sigma_f] \tag{4.6}$$

where V_f is some finite-dimensional representation of Γ. The morphism spaces are correspondingly

$$hom_{\Sigma_\Gamma \mathcal{C}}\Big(\bigoplus_f (X_f \otimes V_f)[\sigma_f], \bigoplus_g (Y_g \otimes W_g)[\tau_g]\Big) = \\ = \bigoplus_{f,g} hom_{\mathcal{C}}(X_f, Y_g)[\tau_g - \sigma_f] \otimes Hom_{\mathbb{C}}(V_f, W_g)$$

and they carry the obvious tensor product Γ-actions. (2) In the definition of an equivariant twisted complex, the differential δ has to belong to the Γ-invariant part of hom^1. The "equivariant derived category" $D^b_\Gamma(\mathcal{C}) = H^0(Tw_\Gamma\,\mathcal{C})$ carries, in addition to the usual shift functor [1], automorphisms $-\otimes\chi$ for any character $\chi \in \Gamma^*$, which change all V_f in (4.6) by tensoring with the corresponding one-dimensional representation.

Remark 4.1. $D^b_\Gamma(\mathcal{C})$ *is* not *triangulated, since mapping cones exist only for the Γ-invariant morphisms. Fortunately, the evaluation maps ev, $ev^\vee$ used in the definition of the twist and untwist operations are invariant, so $T_X(Y)$, $T^\vee_Y(X)$ are well-defined in $D^b_\Gamma(\mathcal{C})$ (at least assuming that the necessary finite-dimensionality conditions hold). In particular, the theory of exceptional collections and mutation works well in the equivariant context.*

(4c) Because we are working over a characteristic zero field, taking Γ-invariants is an exact functor. This means that the classification theory from the previous section extends in a straightforward way to the equivariant situation: one simply replaces all "obstruction groups" by their Γ-fixed parts. More concretely, suppose that we have a finite Γ acting on a graded algebra A. Then $HH^*(A,A)^\Gamma$

is the cohomology of the subcomplex of $CC^*(A,A)^\Gamma$ of equivariant Hochschild cochains. There is a canonical map $J^\Gamma : HH^*(A,A)^\Gamma \to HH^*(A \rtimes \Gamma, A \rtimes \Gamma)$, induced by the cochain homomorphism

$$\begin{aligned} &(J^\Gamma\tau)(a_s \otimes \gamma_s, \dots, a_1 \otimes \gamma_1) = \\ &\qquad = \tau(a_s, \rho(\gamma_s)(a_{s-1}), \rho(\gamma_s\gamma_{s-1})(a_{s-2}), \dots) \otimes \gamma_s \cdots \gamma_1. \end{aligned} \tag{4.7}$$

This is compatible with the Lie bracket on HH^{*+1}, and has an obvious deformation theoretic meaning: Γ-equivariant A_∞-deformations of A induce deformations of the semidirect product. In fact, (4.7) is always injective. One can see this in several different ways. For a classical homological algebra argument, note that if P is an A-bimodule carrying a Γ-action which is compatible with that on A, then $P \rtimes \Gamma$ is an $A \rtimes \Gamma$-bimodule, which for any $A \rtimes \Gamma$-bimodule M satisfies

$$Hom_{A\rtimes\Gamma - A\rtimes\Gamma}(P \rtimes \Gamma, M) \cong Hom_{A-A}(P,M)^\Gamma;$$

here the Γ-action on the right hand side is obtained from that on P and the diagonal action on M, which sends x to $(1 \otimes \gamma)x(1 \otimes \gamma^{-1})$. In particular, if P is projective then so is $P \rtimes \Gamma$. Applying this observation to a projective A-bimodule resolution $P^* \to A$ which carries a compatible Γ-action, one obtains

$$\begin{aligned} HH^*(A \rtimes \Gamma, A \rtimes \Gamma) &= Ext^*_{A\rtimes\Gamma - A\rtimes\Gamma}(A \rtimes \Gamma, A \rtimes \Gamma) \\ &= Hom^*_{A\rtimes\Gamma - A\rtimes\Gamma}(P^* \rtimes \Gamma, A \rtimes \Gamma) \\ &= Hom^*_{A-A}(P^*, A \rtimes \Gamma)^\Gamma. \end{aligned}$$

Under the diagonal action, $A \rtimes \Gamma$ splits into summands corresponding to conjugacy classes $K \subset \Gamma$, and this induces a splitting of $HH^*(A \rtimes \Gamma, A \rtimes \Gamma)$. In particular, the trivial class contributes $Hom^*_{A-A}(P^*, A)^\Gamma = HH^*(A,A)^\Gamma$, which is precisely the image of (4.7). More generally, the summand corresponding to K is the direct sum of the graph bimodules $Graph(\rho(\gamma))$ for $\gamma \in K$, so that

$$HH^*(A \rtimes \Gamma, A \rtimes \Gamma) = \bigoplus_K Ext^*_{A-A}\Big(A, \bigoplus_{\gamma \in K} Graph(\rho(\gamma))\Big)^\Gamma. \tag{4.8}$$

Altenatively, one can approach the injectivity of J^Γ directly in terms of the Hochschild complex. This involves looking at $A \rtimes \Gamma$ as an algebra linear over the semisimple ring $\mathbb{C}\Gamma$. In the abelian case, this can be put more forcefully by saying that we consider the Γ^*-action on $HH^*(A \rtimes \Gamma, A \rtimes \Gamma)$ induced from that on $A \rtimes \Gamma$ itself. The decomposition into different types of Γ^*-modules is precisely (4.8), with the invariant part being the image of J^Γ. This ultimately finds a natural explanation in terms of the duality theory for group actions, which says in particular that $(A \rtimes \Gamma) \rtimes \Gamma^*$ is Morita equivalent to A [8].

The HKR theorem describes the Hochschild cohomology of polynomial algebras, and more general smooth commutative algebras. There is a generalization to graded algebras [42, Proposition 5.4.6], including exterior algebras as a special case, for which the statement is as follows. Take $A = \Lambda V$ for some finite-dimensional complex vector space V, with the usual grading. Let $SV^\vee$ be the symmetric algebra of the dual space. Then there is an explicit isomorphism

$$\phi_{HKR} : S^s V^\vee \otimes \Lambda^{s+t} V \longrightarrow HH^{s+t}(A,A)^t. \tag{4.9}$$

For instance, the cocycle $[\tau] = \phi_{HKR}(p \otimes 1)$ corresponding to a homogeneous polynomial p of degree s is (in terms of a basis v_j and dual basis $v_j^\vee$)

$$\tau(a_s, \dots, a_1) = \frac{1}{s!} \sum_{j_s,\dots,j_1} \frac{\partial p}{\partial v_{j_1} \dots \partial v_{j_s}} (v_{j_s}^\vee \lrcorner\, a_s) \wedge \dots \wedge (v_{j_1}^\vee \lrcorner\, a_1);$$

and conversely, if $\tau \in CC^0(A,A)^{-s}$ is any cocycle, we recover a homogeneous polynomial $p \otimes 1 = \phi_{HKR}^{-1}([\tau])$ by setting

$$p(v) = \tau(v, \dots, v). \tag{4.10}$$

ϕ_{HKR} is compatible with the actions of $GL(V)$ on both sides (where the action on $V^\vee$ is contragredient to that on V). Moreover, it carries the canonical bracket on $HH^{*+1}(A,A)$ to the Schouten bracket on $SV^\vee \otimes \Lambda V$, which in particular satisfies

$$[v_{i_1}^\vee \dots v_{i_r}^\vee \otimes 1, v_{i_{r+1}}^\vee \dots v_{i_s}^\vee \otimes a] = \sum_{k \leq r} v_{i_1}^\vee \dots \widehat{v_{i_k}^\vee} \dots v_{i_s} \otimes v_{i_k}^\vee \lrcorner\, a \tag{4.11}$$

for $a \in \Lambda V$. The standard proof of the HKR formula is via the Koszul bimodule resolution $P^* \to A$, which in our case consists of $P^s = A \otimes S^s V \otimes A[-s]$ with the differential

$$\begin{aligned} \partial(1 \otimes v_{i_1} \dots v_{i_s} \otimes 1) = \sum_k v_{i_k} \otimes v_{i_1} \dots \widehat{v_{i_k}} \dots v_{i_s} \otimes 1 + \\ + (-1)^s 1 \otimes v_{i_1} \dots \widehat{v_{i_k}} \dots v_{i_s} \otimes v_{i_k}. \end{aligned}$$

Because A is supercommutative, the induced differential on $Hom(P^*, A)$ vanishes, which shows that an abstract isomorphism (4.9) exists; of course, the explicit formula and other properties have to be verified separately, by exhibiting a map between the Koszul and bar resolutions.

We now look at (4.8) in the special case where A is an exterior algebra, and ρ the action of a finite abelian group $\Gamma \subset GL(V)$. Taking the Koszul resolution as P^*, one sees that the γ-summand of $HH^*(A \rtimes \Gamma, A \rtimes \Gamma)$ can be obtained by computing the cohomology of the complex $Hom_{\mathbb{C}}(SV, \Lambda V) \cong SV^\vee \otimes \Lambda V$ with the differential

$$(\delta_\gamma \eta)(v_{i_1} \dots v_{i_s}) = (-1)^s \sum_k \eta(v_{i_1} \dots \widehat{v_{i_k}} \dots v_{i_s}) \wedge (\gamma(v_{i_k}) - v_{i_k}),$$

and then restricting to the Γ-invariant part of that. To carry out the computation, we decompose V into eigenspaces of γ, and consider the corresponding tensor product decomposition of $SV^\vee \otimes \Lambda V$. On the fixed part V^γ, the differential vanishes as before; the remaining components can be identified with classical Koszul complexes, which have one-dimensional cohomology. After chasing the bidegrees and Γ-action through the argument, this amounts to a proof of the following:

PROPOSITION 4.2. *The Hochschild cohomology of $A = \Lambda V \rtimes \Gamma$, for Γ a finite abelian subgroup of $GL(V)$, is*

$$\begin{aligned} HH^{s+t}(A,A)^t &\cong \\ &\cong \bigoplus_{\gamma\in\Gamma} \Big(S^s(V^\gamma)^\vee \otimes \Lambda^{s+t-codim(V^\gamma)}(V^\gamma) \otimes \Lambda^{codim(V^\gamma)}(V/V^\gamma)\Big)^\Gamma. \quad \square \end{aligned}$$

5. COHERENT SHEAVES

To apply the theory from Section 3 to the derived category of coherent sheaves, one needs to introduce an underlying differential graded category. Concretely, this means choosing a class of cochain complexes which compute *Ext* groups, with cochain maps realizing the Yoneda product. Out of the many possibilities offered by the literature, we take the lowest-tech path via Čech cohomology, which is easy to work with but slightly more difficult to relate to the usual derived category (a few readers may prefer to skip that argument, and to take the Čech version as definition of the derived category; in which case some of the properties discussed later on have to be taken for granted).

(5a) Let Y be a noetherian scheme over an algebraically closed field ($\mathbb{C}$ or $\Lambda_{\mathbb{Q}}$, in the cases of interest here). Fix a finite affine open cover $\mathfrak{U}$. We introduce a differential graded category $\check{\mathcal{S}}(Y)$ whose objects are locally free coherent sheaves (algebraic vector bundles), and where the morphism spaces are Čech cochain complexes with values in $\underline{Hom}$ sheaves,

$$hom_{\check{\mathcal{S}}(Y)}(E,F) = \check{C}^*(\mathfrak{U}, \underline{Hom}(E,F)).$$

Composition is the shuffle product together with $\underline{Hom}(F,G) \otimes_{\mathcal{O}_Y} \underline{Hom}(E,F) \rightarrow \underline{Hom}(E,G)$. One can think of any dg category as an A_∞-category with vanishing compositions of order ≥ 3, and therefore define a triangulated category $D^b\check{\mathcal{S}}(Y)$ in the way explained in Section 2, as the cohomological category of twisted complexes (we emphasize that there is no "inverting of quasi-isomorphisms" involved in this process). By refining covers and using standard results about Čech cohomology, one sees that $\check{\mathcal{S}}(Y)$ is independent of $\mathfrak{U}$ up to quasi-isomorphism, hence that $D^b\check{\mathcal{S}}(Y)$ is well-defined up to equivalence of triangulated categories.

LEMMA 5.1. *$D^b\check{\mathcal{S}}(Y)$ is equivalent to the full triangulated subcategory of the usual bounded derived category of coherent sheaves, $D^bCoh(Y)$, which is generated by locally free sheaves. If Y is a regular projective variety, this is the whole of $D^bCoh(Y)$.*

PROOF. The usual approach to dg structures on derived categories of coherent sheaves is through injective resolutions, see for instance [62, Section 2]. In our current language, this can be formulated as follows. For each locally free coherent sheaf E, fix a resolution I_E^* by quasi-coherent injective sheaves. Consider a differential graded category $\mathcal{I}(Y)$ with the same objects as $\check{\mathcal{S}}(Y)$, but where $hom_{\mathcal{I}(Y)}(E,F) = hom_{\mathcal{O}_Y}(I_E^*, I_F^*)$ is the space of sheaf homomorphisms $I_E^* \to I_F^*$ of all degrees, with the obvious differential. If we then define $D^b\mathcal{I}(Y) = H^0(Tw\,\mathcal{I}(Y))$ through twisted complexes, it is equivalent to the subcategory of $D^bCoh(Y)$ appearing in the Lemma. To see that, one associates to each object of $Tw\,\mathcal{I}(Y)$ its total complex; this defines an exact, fully faithful functor from $D^b\mathcal{I}(Y)$ to the chain homotopy category $\mathcal{K}^+(Y)$ of bounded below complexes of injective quasi-coherent sheaves. The image of this functor is the triangulated subcategory generated by resolutions of locally free sheaves, and by standard properties of derived categories, this is equivalent to the triangulated subcategory of $D^bCoh(Y)$ generated by such sheaves. If Y is smooth and projective, all coherent sheaves have finite locally free resolutions, so we get the whole of $D^bCoh(Y)$.

To mediate between Čech and injective resolutions, we use a triangular matrix construction. Given two locally free sheaves E and F, one can consider the Čech complex $\check{C}^*(\underline{Hom}(E, I_F^*))$ with coefficients in the complex of sheaves $\underline{Hom}(E, I_F^*) \cong E^\vee \otimes_{\mathcal{O}_Y} I_F^*$. There are canonical cochain maps

$$hom_{\mathcal{O}_Y}(I_E^*, I_F^*) \longrightarrow \check{C}^*(\underline{Hom}(E, I_F^*)) \longleftarrow \check{C}^*(\underline{Hom}(E,F)). \tag{5.1}$$

The first one is defined by composing a sheaf homomorphism $I_E^* \to I_F^*$ with $E \to I_E^0$, and then restricting the associated section of $\underline{Hom}(E, I_F^*)$ to open subsets in $\mathfrak{U}$. The second one is just the map induced from $F \to I_F^0$. Both maps are quasi-isomorphisms: for the first map, note that the Čech complex computes the hypercohomology of $E^\vee \otimes I_F^*$, which on the other hand that is the ordinary cohomology of $hom_{\mathcal{O}_Y}(E^*, I_F^*)$ because each $E^\vee \otimes I_F^k$ is injective. For the second map one uses a standard spectral sequence comparison argument, together with the fact that the cohomology of $hom_{\mathcal{O}_U}(E|U, I_F^*|U)$ for each open subset U is $Hom_{\mathcal{O}_U}(E|U, F|U)$. Along the same lines, there are canonical homomorphisms

$$\begin{aligned} &hom_{\mathcal{O}_Y}(I_F^*, I_G^*) \otimes \check{C}^*(\underline{Hom}(E, I_F^*)) \longrightarrow \check{C}^*(\underline{Hom}(E, I_G^*)),\\ &\check{C}^*(\underline{Hom}(F, I_G^*)) \otimes \check{C}^*(\underline{Hom}(E,F)) \longrightarrow \check{C}^*(\underline{Hom}(E, I_G^*)). \end{aligned} \tag{5.2}$$

We introduce another dg category $\mathcal{T}$, still keeping the same objects as before, and where the *hom*s are the direct sums of three components

$$hom_{\mathcal{T}}(E,F) = \begin{pmatrix} hom_{\mathcal{O}_Y}(I_E^*, I_F^*) & \check{C}^*(\underline{Hom}(E, I_F^*))[1] \\ 0 & \check{C}^*(\underline{Hom}(E,F)). \end{pmatrix}$$

The differential consists of the given differentials on each summand together with the chain homomorphisms (5.1). Similarly, composition in $\mathcal{T}$ combines the usual compositions of morphisms in $\mathcal{I}(Y)$ and $\check{\mathcal{S}}(Y)$ with the maps (5.2). By definition, this comes with dg functors $\mathcal{T} \to \mathcal{I}(Y)$, $\mathcal{T} \to \check{\mathcal{S}}(Y)$, both of which are quasi-isomorphisms, and therefore one has induced exact equivalences $D^b\check{\mathcal{S}}(Y) \cong D^b\mathcal{T} \cong D^b\mathcal{I}(Y)$. □

Remark 5.2. *At least if one supposes that the ground field has characteristic zero, the entire argument carries over to the derived category of equivariant coherent sheaves with respect to the action of a finite group* Γ *on* Y. *First, by taking intersections over* Γ*-orbits one can find an invariant affine open cover. One then introduces the dg category* $\check{S}_\Gamma(Y)$ *whose objects are locally free sheaves with* Γ*-actions, and whose morphism spaces are the* Γ*-invariant summands of the Čech complexes. In the regular projective case,*

$$D^b\check{\mathcal{S}}_\Gamma(Y) \cong D^b Coh_\Gamma(Y),$$

where $Coh_\Gamma(Y)$ *is the abelian category of* Γ*-equivariant coherent sheaves. We will now make the connection between this observation and the material from Section 4. Suppose for simplicity that* Γ *is abelian. If* E *is a coherent sheaf with a* Γ*-action, then so is* $E \otimes W$ *for any finite-dimensional* Γ*-module* W. *Suppose for simplicity that* Γ *is abelian, and take equivariant locally free sheaves* $E_1, \dots, E_m$. *Then the full dg subcategory* $\mathcal{C} \subset \check{\mathcal{S}}(Y)$ *with objects* E_k *carries a* Γ*-action, and the semidirect product* $\mathcal{C} \rtimes \Gamma$ *can be identified with the full dg subcategory of* $\check{\mathcal{S}}_\Gamma(Y)$ *whose objects are* $E_k \otimes W_\chi$ *for all* k *and all one-dimensional representations* W_χ.

(5b) Let Y_q be a noetherian formal scheme, flat over $\Lambda_\mathbb{N}$. By definition [28, ch. II §9] there is a finite cover $\mathfrak{U}$ by open subsets, on each of which the scheme is "given by" (actually, the formal spectrum of) a commutative algebra $A_{q,U}$ over $\Lambda_\mathbb{N}$ which is complete and torsion-free as a $\Lambda_\mathbb{N}$-module. We consider locally free coherent sheaves over Y_q, which are $\mathcal{O}_{Y_q}$-module sheaves given over each U by sections of a finitely generated free $A_{q,i}$-module. By the same procedure as before, one can make these into objects of a dg category $\check{\mathcal{S}}(Y_q)$ whose morphism spaces are again complete torsion-free $\Lambda_\mathbb{N}$-modules.

Geometrically, Y_q is an infinitesimal one-parameter family of schemes. One can specialize the parameter q to two different "values". Setting $q = 0$ yields the *special fibre* $Y = Y_q \times_{\Lambda_\mathbb{N}} \mathbb{C}$, and on the other hand, inverting q and introducing formal roots of it gives the *general fibre* $Y_q^* = Y_q \times_{\Lambda_\mathbb{N}} \Lambda_\mathbb{Q}$. Both of them are noetherian schemes over algebraically closed fields. There are parallel constructions in the world of our differential graded categories, namely full and faithful dg functors

$$\check{\mathcal{S}}(Y_q) \otimes_{\Lambda_\mathbb{N}} \mathbb{C} \longrightarrow \check{\mathcal{S}}(Y_0), \qquad \check{\mathcal{S}}(Y_q) \otimes_{\Lambda_\mathbb{N}} \Lambda_\mathbb{Q} \longrightarrow \check{\mathcal{S}}(Y_q^*). \tag{5.3}$$

To see what this means concretely, pick a finite collection of locally free sheaves $E_{0,1}, \dots, E_{0,m}$ on Y_0, and suppose that each $E_{0,k}$ admits an extension $E_{q,k}$ to a locally free sheaf on Y_q, whose restriction to Y_q^* we denote by $E_{q,k}^*$. The fact that the first map in (5.3) is full and faithful on the cochain level means that the full dg subcategory of $\check{\mathcal{S}}(Y_q)$ with objects $E_{q,k}$ is a one-parameter deformation of the full dg subcategory of $\check{\mathcal{S}}(Y_0)$ with objects $E_{0,k}$. Moreover, if we take the first of these subcategories and tensor with $\Lambda_\mathbb{Q}$, we obtain the full dg subcategory of $\check{\mathcal{S}}(Y_q^*)$ with objects $E_{q,k}^*$. In this way, deformations of varieties and their effects on coherent sheaves have been brought into the fold of the abstract theory from Section 3. This is still quite crude, for instance it excludes totally those sheaves on Y_0 that might not admit extensions to Y_q. One can give a much more natural and satisfactory treatment using obstructed dg or A_∞ structures, see for instance [16], but for our application the rudimentary version is sufficient.

(5c) We end our discussion by listing two properties of $D^bCoh(Y)$ for a regular projective variety over a field, which are well-known to specialists (I am indebted to Drinfeld for suggesting simplified proofs and giving references). The first result is classical, and explains why passage to D^π is unnecessary.

Lemma 5.3. *$D^bCoh(Y)$ is idempotent closed.*

One way to prove this is to consider the chain homotopy category $\mathcal{K}^+(Y)$, which is idempotent closed by an infinite direct sum trick [45, Proposition 1.6.8] or [41, Lemma 2.4.8.1]. Having done that, one recognizes that the direct summand of any object of $D^b(Y) \subset \mathcal{K}^+(Y)$ again lies in $D^b(Y)$, because its cohomology sheaves are coherent and almost all zero [62, Proposition 2.4]. Alternatively, one can use the stronger result from [9] which says that the bounded derived category is *saturated*.

Lemma 5.4. *If $\iota : Y \hookrightarrow \mathbb{P}^N$ is a projective embedding, and $F_1, \dots, F_m$ are split-generators for $D^bCoh(\mathbb{P}^N)$, their derived restrictions $E_k = L\iota^*(F_k)$ are split-generators for $D^bCoh(Y)$.*

This is due to Kontsevich, and the proof follows the same pattern as Lemma 2.7. By assumption, on $\mathbb{P}^N$ one can express $\mathcal{O}(m)$ for any m as direct summand of a complex built from the F_k. Hence, on Y one can express $\mathcal{O}(m)$ in the same way using E_k. It therefore suffices to show that all the $\mathcal{O}(m)$ together are split-generators for $D^bCoh(Y)$. Any locally free coherent sheaf E on Y has a finite left resolution of the form

$$0 \to E' \to \mathcal{O}(m_l)^{\oplus r_l} \to \cdots \to \mathcal{O}(m_1)^{\oplus r_1} \to E \to 0$$

for some E', and one can make l arbitrarily large. For $l > dim\, Y$ we have $Ext^l(E, E') = H^l(E^\vee \otimes E') = 0$, hence the resulting exact triangle

$$\begin{tikzcd}
\left\{\mathcal{O}(m_l)^{\oplus r_l} \to \cdots \to \mathcal{O}(m_1)^{\oplus r_1}\right\} \arrow[r] & E \arrow[d] \\
& E'[l] \arrow[ul, bend left, "{[1]}"]
\end{tikzcd}$$

splits in such a way that $E \oplus E'[l-1]$ is isomorphic to the top left complex. Once one has split-generated all locally free sheaves from the $\mathcal{O}(m)$, the rest follows as in Lemma 5.1.

6. Symplectic terminology

The purpose of this section is to assemble the necessary nuts and bolts from elementary symplectic geometry. The origin of the main notions, to the author's best knowledge, is as follows. Rational Lagrangian submanifolds have a long history in quantization (going back to Einstein); the first place where they occur in connection with Floer homology is probably Fukaya's paper [15]. Lagrangian submanifolds with zero Maslov class also originated in (Maslov's) quantization; the essentially equivalent notion of grading was introduced by Kontsevich [36]. The symplectic viewpoint on Lefschetz pencils and higher-dimensional linear systems emerged gradually from the work of Arnol'd [1], Donaldson [12, 11], Gompf [23], and Auroux [3, 4]. In particular, matching cycles were invented by Donaldson (unpublished).

(6a) By a *projective Kähler manifold* we mean a compact complex manifold X carrying a unitary holomorphic line bundle o_X, which is positive in the sense that the curvature of the corresponding connection A_X defines a Kähler form

$$\omega_X = \frac{i}{2\pi} F_{A_X}.$$

From a symplectic viewpoint, this integrality condition on the symplectic class leads one to make corresponding restrictions on the other objects which live on X. A Lagrangian submanifold $L \subset X$ is called *rational* if the monodromy of the flat connection $A_X|L$ consists of roots of unity. This means that one can choose a covariantly constant multisection λ_L of $o_X|L$ which is of unit length everywhere (as a multisection, this is the multivalued d_L-th root of a section $\lambda_L^{d_L}$ of $o_X^{\otimes d_L}|L$, for some $d_L \geq 1$). Rationality is invariant under exact Lagrangian isotopy. More precisely, take such an isotopy L_t, and embed it into a Hamiltonian isotopy ϕ^t, with time-dependent Hamiltonian H and vector field Y. We obtain covariantly constant multisections λ_{L_t} by starting with λ_{L_0} and solving the ODE

$$\nabla_{Y_t} \lambda_{L_t} = -2\pi i (H_t|L_t)\, \lambda_{L_t}. \tag{6.1}$$

Rationality takes its name from the effect on the periods of action functionals. Take two rational Lagrangian submanifolds L_0, L_1. For each of them, choose a covariantly constant unit length d_{L_k}-fold multisection λ_{L_k}. The *mod* $\mathbb{Q}$ *action* $\bar{A}(x)$ of a point $x \in L_0 \cap L_1$ is defined by

$$e^{2\pi i \bar{A}(x)} = \lambda_{L_1}(x)/\lambda_{L_0}(x). \tag{6.2}$$

It is unique up to adding a number in $l.c.m.(d_{L_0}, d_{L_1})^{-1}\mathbb{Z} \subset \mathbb{Q}$, whence the name. Now suppose that we have two intersection points x_0, x_1 and a connecting disc between them: this is a smooth map $u : \mathbb{R}\times[0;1] \to X$ with $u(\mathbb{R}\times\{k\}) \subset L_k$, such that $u(s, \cdot)$ converges to the constant paths at x_0, x_1 as $s \to -\infty, +\infty$. By definition of curvature,

$$\int u^*\omega_X \in \bar{A}(x_0) - \bar{A}(x_1) + l.c.m.(d_{L_0}, d_{L_1})^{-1}\mathbb{Z}. \tag{6.3}$$

Similarly, a symplectic automorphism ϕ of X is rational if the monodromy of the induced flat connection on $Hom(o_X, \phi^* o_X)$ consists of roots of unity. This is invariant under Hamiltonian isotopy. We write $Aut(X)$ for the group of such automorphisms, which acts in the obvious way on the set of rational Lagrangian submanifolds.

(6b) Equally important for us will be the relative situation, where in addition to the previous data we have a holomorphic section $\sigma_{X,\infty}$ of o_X such that $X_\infty = \sigma_{X,\infty}^{-1}(0)$ is a divisor with normal crossings, along which $\sigma_{X,\infty}$ vanishes with multiplicity one, and such that each irreducible component $C \subset X_\infty$ is smooth (no self-intersections). We call $M = X \setminus X_\infty$ an *affine Kähler manifold.* One can use $\sigma_{X,\infty}/||\sigma_{X,\infty}||$ to trivialize $o_X|M$, and by writing the connection with respect to this trivialization as $A_X|M = d - 2\pi i\theta_M$, one gets a one-form θ_M which satifies $d\theta_M = \omega_M = \omega_X|M$, so that M is an exact symplectic manifold in the usual sense of the word.

A compact Lagrangian submanifold $L \subset M$ is called *exact* if $\theta_M|L = dK_L$ for some function K_L. This condition is invariant under exact Lagrangian isotopy. Note that if L is exact,

$$\lambda_L = \exp(2\pi i K_L)\frac{\sigma_{X,\infty}}{||\sigma_{X,\infty}||} \tag{6.4}$$

is a covariantly constant section of $o_X|L$, so that must be the trivial flat line bundle. Conversely, suppose that L is some Lagrangian submanifold of M, such that $o_X|L$ is trivial and admits a covariantly constant section which can be written in the form (6.4) for some K_L. Then $dK_L = \theta_M|L$, hence L is exact. We have proved:

LEMMA 6.1. *L is exact iff $o_X|L$ is trivial, and its nonzero covariantly constant sections lie in the same homotopy class of nowhere zero sections as $\sigma_{X,\infty}|L$.* □

Let L_0, L_1 be exact Lagrangian submanifolds of M, and choose functions K_{L_0}, K_{L_1}. One can then define the action of an intersection point $x \in L_0 \cap L_1$ to be

$$A(x) = K_{L_1}(x) - K_{L_0}(x) \in \mathbb{R}. \tag{6.5}$$

For any two such points x_0, x_1 and any connecting disc u in X, the intersection number $u \cdot X_\infty$ is well-defined, and

$$\int u^*\omega_X = A(x_0) - A(x_1) + (u \cdot X_\infty). \tag{6.6}$$

In particular, the action functional is indeed exact if we allow only connecting discs which lie in M.

Let ϕ be a symplectic automorphism of M, which extends smoothly to X in such a way that X_∞ remains pointwise fixed (we denote the extension equally by ϕ). We say that ϕ exact if $\phi^*\theta_M - \theta_M = dK_\phi$ for some function K_ϕ on M. This condition is invariant under Hamiltonian isotopies keeping X_∞ fixed. Exact symplectic automorphisms form a group $Aut(M)$, which acts naturally on the set of exact Lagrangian submanifolds. The analogue of Lemma 6.1 is

Lemma 6.2. *ϕ is exact iff $Hom(o_X, \phi^* o_X)$ is the trivial flat bundle, and every covariantly constant nonzero section has the property that its restriction to M lies in the same homotopy class of nowhere zero sections as $\phi^*\sigma_{X,\infty}/\sigma_{X,\infty}$.* □

Remark 6.3. *The conditions for exactness which appear in the Lemma are automatically satisfied if $H_1(X) = 0$. This is obvious for the first one, and for the second one argues as follows: take the meridian γ in M around a component of X_∞, and bound it by a small disc going through X_∞ once. The homotopy class of covariantly constant sections of $Hom(o_X, \phi^* o_X)|\gamma$ is characterized by the fact that they extend to nowhere zero sections over the disc. On the other hand, $\sigma_{X,\infty}\gamma$ extends to a section with precisely one zero, and so does $\phi^*\sigma_{X,\infty}|\gamma$. Hence, $(\phi^*\sigma_{X,\infty}/\sigma_{X,\infty})|\gamma$ lies in the same homotopy class of nowhere zero sections as the covariantly constant ones.*

The relation between the affine and projective notions is clear: Lemma 6.1 shows that an exact $L \subset M$ is a rational Lagrangian submanifold of X, and Lemma 6.2 does the same for automorphisms. Given a pair of exact Lagrangian submanifolds L_0, L_1, the multivalued action (6.2) is a reduction of the single-valued one (6.5), and (6.6) is a refinement of (6.3).

(6c) Let M be an affine Kähler manifold, coming from X and $\sigma_{X,\infty}$ as always. Suppose that we have a second holomorphic section $\sigma_{X,0}$ of o_X, linearly independent of $\sigma_{X,\infty}$. Let $\{X_z\}_{z\in\mathbb{CP}^1}$ be the pencil of hypersurfaces generated by these two sections. This means that we set

$$\sigma_{X,z} = \sigma_{X,0} - z\sigma_{X,\infty} \quad \text{and} \quad X_z = \sigma_{X,z}^{-1}(0)$$

for $z \in \mathbb{C}$. Assume that $X_0 = \sigma_{X,0}^{-1}(0)$ is smooth in a neighbourhood of the base locus $X_{0,\infty} = X_0 \cap X_\infty$, and that it intersects each stratum of X_∞ transversally; then the same will hold for each X_z, $z \in \mathbb{C}$. Every nonsingular X_z, $z \in \mathbb{C}$, is a projective Kähler manifold with ample line bundle $o_{X_z} = o_X|X_z$, and it carries a preferred holomorphic section $\sigma_{X_z} = \sigma_{X,\infty}|X_z$ whose zero set is $X_{0,\infty}$. By assumption, this is a divisor with normal crossings, so that $M_z = X_z \setminus X_{0,\infty}$ is an affine Kähler manifold. One can also see the M_z as the regular fibres of the holomorphic function

$$\pi_M = \sigma_{X,0}/\sigma_{X,\infty} : M \longrightarrow \mathbb{C}.$$

We denote by $Critv(\pi_M)$ its critical value set, which is finite by Bertini. Given $z, z' \in \mathbb{C} \setminus Critv(\pi_M)$, we denote by $Iso(M_z, M_{z'})$ the space of symplectic isomorphisms $\phi : M_z \rightarrow M_{z'}$ which extend to $X_z \rightarrow X_{z'}$ in such a way that $X_0 \cap X_\infty$ remains pointwise fixed, and such that $\phi^*\theta_{M_{z'}} - \theta_{M_z}$ is exact. These spaces form a topological groupoid under composition, and for $z = z'$ they reduce to the groups $Aut(M_z)$ defined before. Take a smooth path $c : [0; l] \rightarrow \mathbb{C} \setminus Critv(\pi_M)$. The $c'(t)$ can be lifted in a unique way to vector fields $Z_t \in C^\infty(TM|M_{c(t)})$ which are horizontal (orthogonal to $ker(D\pi_M)$ with respect to the symplectic form). By integrating these one gets an exact *symplectic parallel transport* map

$$h_c \in Iso(M_{c(0)}, M_{c(l)}).$$

To show that h_c is well-defined, one embeds M into the graph $\Gamma = \{(z, x) \in \mathbb{C} \times X_z \ : \ x \in X_z\}$. The projection $\pi_\Gamma : \Gamma \rightarrow \mathbb{C}$ is a properification of π_M. It is a standard fact that parallel transport with respect to the product symplectic form $\omega_\Gamma = (\omega_X + \omega_{\mathbb{C}})|\Gamma$ gives a symplectic isomorphism $X_{c(0)} \rightarrow X_{c(l)}$. This extends the original h_c, because adding the pullback of $\omega_{\mathbb{C}}$ to ω_M does not change the vector fields Z_t. Moreover, at a point $(z, x) \in \mathbb{C} \times X_{0,\infty}$ we have

$$T\Gamma_{(z,x)} = \mathbb{C} \times \{\xi \in TX_x \ : \ D\sigma_0(x)\xi - z\, D\sigma_\infty(x)\xi = 0\},$$

and since this splitting is ω_Γ-orthogonal, the parallel transport vector field will lie in $\mathbb{C} \times \{0\}$. This implies that the extension of h_c to $X_{c(0)}$ leaves $X_{0,\infty}$ pointwise fixed. Finally, the exactness of $h_c^*\theta_{M_{c(l)}} - \theta_{M_{c(0)}}$ follows from the Cartan formula $(L_{Z_t}\theta_M) \,|\, M_{c(t)} = d(i_{Z_t}\theta_M) \,|\, M_{c(t)} + i_{Z_t}\omega_M \,|\, M_{c(t)}$, where the last term vanishes because of the horizontality of Z_t.

(6d) We say that $\{X_z\}$ is an *quasi-Lefschetz pencil* if π_M is nondegenerate in the sense of Picard-Lefschetz theory, meaning that it has only nondegenerate critical points, no two of which lie in the same fibre. This is the same as saying that the set of critical points $Crit(\pi_M)$ is regular ($D\pi_M$ is transverse to the zero-section) and $\pi_M : Crit(\pi_M) \rightarrow Critv(\pi_M)$ is bijective. Lefschetz pencils are the special case where X_∞ is smooth.

An *embedded vanishing path* for a quasi-Lefschetz pencil is an embedded path $c : [0; 1] \rightarrow \mathbb{C}$ with $c^{-1}(Critv(\pi_M)) = \{1\}$. By considering the limiting behaviour of parallel transport along $c|[0; t]$ as $t \rightarrow 1$, one defines an embedded Lagrangian disc $\Delta_c \subset M$, the *Lefschetz thimble*, whose boundary is a Lagrangian sphere $V_c \subset M_{c(0)}$, the associated *vanishing cycle*. The Picard-Lefschetz formula [60, Proposition 1.15] says that the monodromy around a loop γ which doubles around c in positive sense is the Dehn twist associated to the vanishing cycle, up to isotopy inside the group of exact symplectic automorphisms:

$$h_\gamma \simeq \tau_{V_c} \in Aut(M_{c(0)}).$$

To be precise, V_c carries a small additional piece of structure, called a framing in [58, 60], and the definition of Dehn twist uses that too. However, framings contain no information if the vanishing cycles are of dimension ≤ 3, so omitting them will not matter for our applications.

Now choose a base point $z_* \in \mathbb{C} \setminus Critv(\pi_M)$ and base path $c_* : [0;\infty) \to \mathbb{C} \setminus Critv(\pi_M)$, meaning an embedded path with $c_*(0) = z_*$ which goes off to infinity, in the sense that there is a $\zeta \in \mathbb{C}^*$ such that $c_*(t) = \zeta t$ for all $t \gg 0$. By a *distinguished basis* of embedded vanishing paths, we mean an ordered collection of such paths $(c_1, \dots, c_r)$, $r = |Critv(\pi_M)|$, starting at z_* and with the following properties: the tangent directions $\mathbb{R}^{\geq 0} c_k'(0)$ are pairwise distinct and clockwise ordered, and different c_k do not intersect except at z_*. Moreover, $\mathbb{R}^{\geq 0} c_*'(0)$ should lie strictly between $\mathbb{R}^{\geq 0} c_r'(0)$ and $\mathbb{R}^{\geq 0} c_1'(0)$, and no c_k may intersect c_* except at z_*. The composition of the corresponding loops γ_k is freely homotopic in $\mathbb{C} \setminus Critv(\pi_M)$ to a circle $\gamma_\infty(t) = R\zeta e^{it}$ of some large radius $R \gg 0$, and hence the product of all the associated Dehn twists is the monodromy around that circle. More precisely, taking into account the change of base point, one has

$$\tau_{V_{c_1}} \dots \tau_{V_{c_r}} \simeq (h_{c_*|[0;R]})^{-1} h_{\gamma_\infty} h_{c_*|[0;R]} \in Aut(M_{z_*}). \tag{6.7}$$

This becomes more familiar for actual Lefschetz pencils, where one can extend the graph to $\mathbb{C}\mathrm{P}^1$,

$$\bar{\Gamma} = \{(z,x) \in \mathbb{C}\mathrm{P}^1 \times X \ : \ x \in X_z\} \xrightarrow{\pi_{\bar{\Gamma}}} \mathbb{C}\mathrm{P}^1, \tag{6.8}$$

so that ∞ is a regular value of $\pi_{\bar{\Gamma}}$. Proceeding as before, one can define symplectic parallel transport for paths in $\mathbb{C}\mathrm{P}^1 \setminus Critv(\pi_M)$. In particular, since γ_∞ can be contracted by passing over ∞, its monodromy is isotopic to the identity, so that (6.7) simplifies to the following relation between Dehn twists in the symplectic mapping class group $\pi_0(Aut(M_{z_*}))$:

$$\tau_{V_{c_1}} \circ \dots \circ \tau_{V_{c_r}} \simeq id. \tag{6.9}$$

Remark 6.4. *It is no problem (except terminologically) to allow several non-degenerate critical points to lie in the same fibre. One should then choose a vanishing path for every critical point. Two paths c_k, c_l going to the same critical value should either coincide or else satisfy $c_k(t) \neq c_l(t)$ for $t \in (0;1)$, and $\mathbb{R}^{\geq 0} c_k'(0) \neq \mathbb{R}^{\geq 0} c_l'(0)$. In the first case, the associated vanishing cycles in M_{z_*} will be disjoint.*

(6e) Let $\{X_z\}$ be a quasi-Lefschetz pencil. Consider a smooth embedded path $d : [-1;1] \to \mathbb{C}$ such that $d^{-1}(Critv(\pi_M)) = \{-1;1\}$. Split it into a pair of vanishing paths $c_-(t) = d(-t)$, $c_+(t) = d(t)$, $t \in [0;1]$, and consider the associated vanishing cycles

$$V_{c_-}, V_{c_+} \subset M_{d(0)}.$$

We say that d is a *matching path* if V_{c_-}, V_{c_+} are isotopic as exact Lagrangian spheres. Actually, there is an additional condition concerning the framings, which we do not want to spell out here, since it is vacuous in the dimensions that we will be interested in. If d is a matching path, one can glue together the Lefschetz thimbles Δ_{c_-}, Δ_{c_+} to obtain a Lagrangian sphere in the total space, the *matching cycle* $\Sigma_d \subset M$. In the naive situation where the two vanishing cycles coincide, this would simply be the set-theoretic union of the Lefschetz thimbles. In general, the construction involves a choice of isotopy between the cycles, and Σ_d might theoretically depend on that choice. Note however that this can never happen if the fibres are Riemann surfaces with boundary, since there, the space of exact Lagrangian submanifolds (circles) having a fixed isotopy class is contractible.

(6f) Let $\sigma_{X,0}, \sigma_{X,\infty}$ be two sections of o_X which generate a quasi-Lefschetz pencil $\{X_z\}$. Suppose that we have yet another section $\sigma'_{X,0}$, linearly independent from the previous ones. We require that its zero set should be smooth in a neighbourhood of $X_0 \cap X_\infty$, and should intersect each stratum of that base locus transversally. Consider the associated map

$$M \xrightarrow{b_M = \left(\frac{\sigma_{X,0}}{\sigma_{X,\infty}}, \frac{\sigma'_{X,0}}{\sigma_{X,\infty}}\right)} \mathbb{C}^2, \tag{6.10}$$

its critical point set $Crit(b_M)$ and critical value set $C = Critv(b_M)$. Concerning this, we make the following additional assumptions:

- $Crit(b_M)$ is regular, and $b_M : Crit(b_M) \rightarrow C$ is an embedding away from finitely many points.

Regularity means that $Db_M \in \Gamma(Hom(TM, \mathbb{C}^2))$ is never zero, and is transverse to the subset of rank one linear maps. The second statement implies that for generic $z \in \mathbb{C} \setminus Critv(\pi_M)$, the curve $Crit(b_M)$ is transverse to $b_M^{-1}(\{z\} \times \mathbb{C})$, and the projection $b_M | Crit(b_M) \times b_M^{-1}(\{z\} \times \mathbb{C}) \rightarrow C \times (\{z\} \times \mathbb{C})$ is a bijection. In this case, the sections $(\sigma'_{X,0}|X_z, \sigma_{X,\infty}|X_z)$ generate a quasi-Lefschetz pencil on X_z, and the critical value set of the associated holomorphic function $q_{M_z} : M_z \rightarrow \mathbb{C}$ is precisely

$$Critv(q_{M_z}) = C \cap (\{z\} \times \mathbb{C}) \subset \mathbb{C}. \tag{6.11}$$

There is a finite subset of $\mathbb{C} \setminus Critv(\pi_M)$ where this fails, and we denote it by $Fakev(\pi_M)$. These "fake critical values" will play a role similar to the real ones, see (6.12) below.

- Let x be a critical point of π_M, $\pi_M(x) = z$. Then the restriction of $D^2\pi_M$ to the complex codimension one subspace $ker(D_x b_M)$ is nondegenerate. Moreover, all critical points of $q_{M_z}|(M_z \setminus \{x\})$ are nondegenerate, and their q_{M_z}-values are pairwise distinct and different from $q_{M_z}(x)$.

As a consequence, $b_M : Crit(b_M) \rightarrow C$ is an embedding on $Crit(b_M) \cap M_z$, and moreover the projection $C \subset \mathbb{C}^2 \rightarrow \mathbb{C}$ to the first variable has an ordinary (double) branch point at $b_M(x)$, and is a local isomorphism at all other points of $C \cap (\{z\} \times \mathbb{C})$.

If the two conditions stated above are satisfied, we say that $\sigma'_{X,0}$ is a *generic auxiliary section* of o_X. Practically, the situation is that we have a quasi-Lefschetz pencil on X, almost every regular fibre X_z of which again admits a quasi-Lefschetz pencil. Moreover, we have a good understanding of how these pencils degenerate as X_z becomes singular. We will now explain the consequences of this for the vanishing cycles of π_M. For that, suppose that our base point $z_* \notin Fakev(\pi_M)$. Alongside the ordinary monodromy of the pencil,

$$\pi_1(\mathbb{C} \setminus Critv(\pi_M), z_*) \longrightarrow \pi_0(Aut(M_{z_*})),$$

there is a "relative" or "braid" monodromy homomorphism

$$\pi_1(\mathbb{C} \setminus (Critv(\pi_M) \cup Fakev(\pi_M)), z_*) \longrightarrow \pi_0(\mathit{Diff}^c(\mathbb{C}, (C \cap \{z_*\} \times \mathbb{C}))), \tag{6.12}$$

which describes how the finite set (6.11) moves in $\mathbb{C}$ as z changes. Take an embedded vanishing path for π_M, $c : [0;1] \to \mathbb{C}$, with the additional property that $c([0;1]) \cap Fakev(\pi_M) = \emptyset$. Let γ be a loop doubling around c in positive sense. As we know, the monodromy around γ is the Dehn twist along V_c. The corresponding statement for (6.12) is that there is an embedded path $d : [-1;1] \to \mathbb{C}$, with $d^{-1}(Critv(q_{M_{z_*}})) = \{\pm 1\}$, such that the image of γ under the relative monodromy is the half-twist along d. The relation between the two is as follows, see [61] for the proof:

Proposition 6.5. *d is a matching path for $q_{M_{z_*}}$, and the vanishing cycle $V_c \subset M_{z_*}$ is, up to Lagrangian isotopy, a matching cycle Σ_d for that path.* $\square$

(6g) We temporarily digress from Kähler to general symplectic geometry. Let (M^{2n}, ω_M) be any connected symplectic manifold with $c_1(M) = 0$. Take an almost complex structure J_M which tames ω_M, and a complex volume form (a nowhere vanishing $(n,0)$-form with respect to J_M) η_M. Let $\mathfrak{L}_M \to M$ be the bundle of (unoriented) Lagrangian Grassmannians. The phase function associated to η_M,

$$\alpha_M : \mathfrak{L}_M \longrightarrow S^1,$$

is defined as follows. Given $x \in M$ and $\Lambda \in \mathfrak{L}_{M,x}$, take any (not necessarily orthonormal) basis $e_1, \dots, e_n$ of Λ, and set

$$\alpha_M(\Lambda) = \frac{\eta_M(e_1 \wedge \dots \wedge e_n)^2}{|\eta_M(e_1 \wedge \dots \wedge e_n)|^2}. \tag{6.13}$$

We distinguish the special case where J_M is ω_M-compatible by calling α_M a *classical phase function*. These are the most commonly used ones, and they are much better behaved. General phase functions will be needed later for technical reasons.

Given α_M, one can associate to a Lagrangian submanifold $L \subset M$ or a symplectic automorphism $\phi : M \to M$ a phase function

$$\begin{cases} \alpha_L : L \longrightarrow S^1, \quad \alpha_L(x) = \alpha_M(TL_x), \\ \alpha_\phi : \mathfrak{L}_M \longrightarrow S^1, \quad \alpha_\phi(\Lambda) = \alpha_M(D\phi(\Lambda))/\alpha_M(\Lambda). \end{cases}$$

A grading of L or ϕ is a real-valued lift $\tilde{\alpha}_L$ resp. $\tilde{\alpha}_\phi$, where the convention is that $\mathbb{R}$ covers S^1 by $a \mapsto \exp(2\pi i a)$. Pairs $\tilde{L} = (L, \tilde{\alpha}_L)$, $\tilde{\phi} = (\phi, \tilde{\alpha}_\phi)$ are called graded Lagrangian submanifolds resp. graded symplectic automorphisms. The latter form a group, with a natural action on the set of graded Lagrangian submanifolds. The composition law and action are defined by

$$\begin{cases} \tilde{\alpha}_{\psi\circ\phi} = \tilde{\alpha}_\psi \circ D\phi + \tilde{\alpha}_\phi, \\ \tilde{\alpha}_{\phi(L)} = \tilde{\alpha}_L \circ \phi^{-1} + \tilde{\alpha}_\phi \circ D\phi^{-1}. \end{cases} \tag{6.14}$$

In particular, for $k \in \mathbb{Z}$ we have the k-fold shift $[k] = (\phi = \mathrm{id}_M, \tilde{\alpha}_\phi = -k)$, which reduces the grading of each Lagrangian submanifold by k. All the "graded" notions actually depend only on the homotopy class of η_M^2 as a trivialization of K_M^2, or equivalently on the cohomology class $\mu_M = [\alpha_M] \in H^1(\mathfrak{L}_M)$, called a global Maslov class in [57]. Namely, suppose that we have two cohomologous phase functions, which can be written as $\alpha'_M = \alpha_M e^{2\pi i\chi}$ for some $\chi : \mathfrak{L}_M \to \mathbb{R}$. Gradings with respect to the two are related in an obvious way,

$$\begin{cases} \tilde{\alpha}'_L(x) = \tilde{\alpha}_L(x) + \chi(TL_x), \\ \tilde{\alpha}'_\phi(\Lambda) = \tilde{\alpha}_\phi(\Lambda) + \chi(D\phi(\Lambda)) - \chi(\Lambda). \end{cases} \tag{6.15}$$

Note that χ is unique only up to an integer constant. This is irrelevant for $\tilde{\alpha}'_\phi$; it affects $\tilde{\alpha}'_L$ but not in a really important way, since changing χ shifts the grading of all Lagrangians simultaneously.

REMARK 6.6. *There is a canonical square root* $\alpha_M^{1/2} : \mathfrak{L}^{\mathrm{or}} \to \mathbb{R}$ *of* α_M *on the bundle* $\mathfrak{L}^{\mathrm{or}}$ *of oriented Lagrangian Grassmannians, defined by removing the squares from* (6.13) *and using positively oriented bases* $e_1, \dots, e_n$. *Any graded Lagrangian submanifold carries a preferred orientation, characterized by*

$$\exp(\pi i \tilde{\alpha}_L(x)) = \alpha_M^{1/2}(TL_x).$$

(6h) There is an obvious notion of graded symplectic fibre bundle, which is a locally trivial symplectic fibration $p : E \to B$ with an almost complex structure J_{E_b} and complex volume form η_{E_b} on each fibre E_b. The monodromy maps of such a fibration are naturally graded: given a loop $\gamma : [0;l] \to B$ and the corresponding parallel transport maps $\phi_t = h_{\gamma|[0;t]} : E_{\gamma(0)} \to E_{\gamma(t)}$, one takes the unique smooth family of functions $\tilde{a}_t$ on the Lagrangian Grassmannian bundle of $E_{\gamma(0)}$ with $\tilde{a}_0 = 0$ and

$$\exp(2\pi i \tilde{a}_t) = (\alpha_{E_{\gamma(t)}} \circ D\phi_t)/\alpha_{E_{\gamma(0)}}.$$

The graded symplectic monodromy is $\tilde{h}_\gamma = (\phi_l, \tilde{\alpha}_{\phi_l} = \tilde{a}_l)$. An example which is relevant for our purpose is where we have a trivial fibre bundle $E = \mathbb{C}^* \times M$ over $B = \mathbb{C}^*$ with a varying complex volume form $\eta_{E_z} = z^\mu \eta_M$. The "invisible singularity" of this family at $z = 0$ leads to a monodromy which, for a circle γ turning around the origin in clockwise direction, is the shift

$$\tilde{h}_\gamma = [2\mu]. \tag{6.16}$$

(6i) Gradings make it possible to assign absolute Maslov indices to Lagrangian intersection points. This depends on the following notion from symplectic linear algebra: a smooth path of Lagrangian subspaces Λ_t $(0 \leq t \leq 1)$ in a fixed symplectic vector space is *crossingless* if it satisfies $\Lambda_0 \cap \Lambda_t = \Lambda_0 \cap \Lambda_1$ for all $t > 0$, and the crossing form [51] on $\Lambda_0/\Lambda_0 \cap \Lambda_1$ associated to $(d\Lambda_t/dt)_{t=0}$ is negative definite. Nonintersecting paths with given endpoints exist and are unique up to homotopy, see again [51].

LEMMA 6.7. *Suppose that α_M is classical. Let Λ_t be a crossingless path in TM_x for some x. Choose $\tilde{a} : [0;1] \to \mathbb{R}$ such that $\exp(2\pi i \tilde{a}_t) = \alpha_M(\Lambda_t)$. Then*

$$\tilde{a}_1 - \tilde{a}_0 \in (-n;0].$$

PROOF. We can identify $\Phi : TM_x \to \mathbb{C}^n$ in such a way that the complex structure and symplectic form become standard, $\Phi(\Lambda_0) = \mathbb{R}^n$ and $\Phi(\Lambda_1) = e^{i\pi c_1}\mathbb{R} \times \cdots \times e^{i\pi c_n}\mathbb{R}$ with $c_k \in (-1;0]$. For the obvious crossingless path $\Lambda_t = e^{i\pi t c_1}\mathbb{R} \times \cdots \times e^{i\pi t c_n}\mathbb{R}$ we have $\tilde{a}_1 - \tilde{a}_0 = c_1 + \cdots + c_n \in (-n;0]$. Because of homotopy uniqueness, the same holds for any other crossingless path. □

Take two graded Lagrangian submanifolds $\tilde{L}_0, \tilde{L}_1$, and a point $x \in L_0 \cap L_1$. Choose a crossingless path from $\Lambda_0 = TL_0$ to $\Lambda_1 = TL_1$, and take $\tilde{a}_t \in \mathbb{R}$ such that $\exp(2\pi i \tilde{a}_t) = \alpha_M(\Lambda_t)$. The absolute Maslov index is defined as

$$I(x) = (\tilde{\alpha}_{L_1}(x) - \tilde{a}_1) - (\tilde{\alpha}_{L_0}(x) - \tilde{a}_0) \in \mathbb{Z}. \tag{6.17}$$

Lemma 6.7 implies that this is approximately the difference of the phases:

LEMMA 6.8. *If α_M is classical, $I(x) - \tilde{\alpha}_{L_1}(x) + \tilde{\alpha}_{L_0}(x) \in [0;n)$.* □

This was pointed out to me by Joyce, but it is known to many other people in mathematics and physics, see for instance the proof of [65, Theorem 4.3]. In fact, ideas of this kind can be traced back at least to [56]. Another easy consequence of Lemma 6.7, this time for symplectic automorphisms, will be useful later:

LEMMA 6.9. *Suppose that α_M is classical. Let $\tilde{\phi} = (\phi, \tilde{\alpha}_\phi)$ a graded symplectic automorphism of M. For any two Lagrangian subspaces $\Lambda_0, \Lambda_1 \in \mathfrak{L}_{M,x}$ at the same point,*

$$|\tilde{\alpha}_\phi(\Lambda_1) - \tilde{\alpha}_\phi(\Lambda_0)| < n.$$

PROOF. Choose a crossingless path Λ_t. Lemma 6.7 shows that as we go along this path, $\alpha_M(\Lambda_t)$ changes by an angle in $(-2\pi n;0]$, and so does $\alpha_M(D\phi(\Lambda_t))$. The angular change in $\alpha_M(D\phi(\Lambda_t))/\alpha_M(\Lambda_t)$ is the difference between the two, hence lies in $(-2\pi n; 2\pi n)$. □

(6j) We now return to the case of an affine Kähler manifold $M = X \setminus X_\infty$. We say that M is *affine Calabi-Yau* if it comes with a preferred isomorphism of holomorphic line bundles,

$$\beta_X : \mathcal{K}_X \xrightarrow{\cong} o_X^{\otimes -m_X} \tag{6.18}$$

for some $m_X \in \mathbb{Z}$ (called the monotonicity index). On M we have a preferred trivialization of o_X given by σ_∞, hence through β_X a holomorphic volume form η_M. In the special case where $m_X = 0$, X itself is Calabi-Yau, and η_M extends to a holomorphic volume form η_X on it. We denote by $\widetilde{Aut}(M)$ the group of exact symplectic automorphisms of M equipped with a grading. Similarly, if $m_X = 0$, we write $\widetilde{Aut}(X)$ for the graded rational automorphisms of X.

Remark 6.10. *In the $m_X = 0$ case, there is a canonical embedding*

$$\widetilde{Aut}(M) \longrightarrow \widetilde{Aut}(X).$$

We already know that every $\phi \in Aut(M)$ extends to a rational symplectic automorphism of X. There are no obstructions to extending gradings over subsets of codimension ≥ 2, so the grading of ϕ extends uniquely from M to X.

If $\{X_z \subset X\}$ is a quasi-Lefschetz pencil, we have for $z \in \mathbb{C} \setminus Critv(\pi_M)$ a canonical isomorphism $\beta_{X_z} : \mathcal{K}_{X_z} \cong (\mathcal{K}_X \otimes o_X)|X_z \cong o_X^{\otimes -m_X+1}$, so the M_z are again affine Calabi-Yaus, with monotonicity index $m_{X_z} = m_X - 1$. More explicitly, the complex volume form is obtained as a quotient η_M/dz, which means that it is the unique solution of

$$dz \wedge \eta_{M_z} = \eta_M. \tag{6.19}$$

The first consequence of this is that the monodromy maps along loops γ in $\mathbb{C} \setminus Critv(\pi_M)$ have canonical gradings, $\tilde{h}_\gamma \in \widetilde{Aut}(M_{\gamma(0)})$. If c is a vanishing path, the associated vanishing cycle V_c admits a grading; this is trivial if its dimension is ≥ 2, and otherwise follows from the fact that it is bounded by a Lagrangian disc in M. Hence, the associated Dehn twist τ_{V_c} then has a canonical grading $\tilde{\tau}_{V_c}$, which is zero outside a neighbourhood of V_c itself. Taking the loop γ which doubles c, we have the following graded version of the Picard-Lefschetz theorem:

$$\tilde{h}_\gamma \simeq \tilde{\tau}_{V_c} \in \widetilde{Aut}(M_{c(0)}).$$

The analogue of (6.9) is

Lemma 6.11. *Suppose that $\{X_z\}$ is a Lefschetz pencil, and $c_1, \dots, c_m$ a distinguished basis of vanishing cycles. Then*

$$\tilde{\tau}_{V_{c_1}} \circ \cdots \circ \tilde{\tau}_{V_{c_r}} \simeq [4 - 2m_X] \in \widetilde{Aut}(M_{z_*}).$$

Proof. What we need to show is that the graded monodromy around a large circle γ_∞ is $[4 - 2m_X]$. For simplicity, change variables to $\zeta = z^{-1}$, so that γ_∞ becomes a small circle going clockwise around $\zeta = 0$. Recall that η_{M_z} is obtained by dividing η_M, which has a zero or pole of order $-m_X$ along X_∞, by $dz = d\zeta/\zeta^2$, which has order -2. Hence, after introducing the compactified graph $\bar{\Gamma}$ as in (6.8), one has a locally trivial symplectic fibration near $\zeta = 0$, with a family of complex volume forms on the fibres that grows like ζ^{2-m_X}. Now we are in the same situation as in (6.16). □

7. Monodromy and negativity

Continuing the discussion above, suppose that $m_X = 1$, so that the fibres X_z are Calabi-Yau manifolds. In the Lefschetz pencil case, Lemma 6.11 tells us that the "large complex structure limit monodromy" is a downward shift. We will need a generalization of this statement to the case when X_∞ has normal crossings, where the geometry of the monodromy map becomes more complicated. Intuition is provided by two aspects of mirror symmetry: ideas related to the SYZ conjecture suggest that large complex structure limit monodromy maps should be fibrewise translations in a Lagrangian torus fibration; and in the context of homological mirror symmetry we have a formula for the mirror automorphism of the derived category of sheaves which goes back to Kontsevich, see [2]. Because of technical difficulties, our result will finally be limited to $K3$ surfaces; but large parts of the argument are more general, and it seems likely that with some more effort, the dimensional restriction could be lifted altogether.

(7a) We start with some generalities. Let M^{2n} be a compact connected symplectic manifold (possibly with boundary), with a tame almost complex structure J_M, complex volume form η_M, and associated phase function α_M.

Definition 7.1. *A graded symplectic automorphism $\tilde{\phi}$ of M is* negative *if there is a $d_0 > 0$ such that*

$$\tilde{\alpha}_{\phi^{d_0}}(\Lambda) < 0 \quad \textit{for all } \Lambda \in \mathfrak{L}_M.$$

Here, the grading of ϕ^{d_0} is defined by the composition rule (6.14). The same rule shows that if negativity holds, the grading of the iterates goes to $-\infty$. Namely, the grading of ϕ^{d_0} is $\leq -\epsilon$ for some $\epsilon > 0$, and the grading of ϕ is $\leq C$ for some $C > 0$. By writing $d = kd_0 + l$ with $0 \leq l \leq d_0 - 1$ one sees that

$$\tilde{\alpha}_{\phi^d}(\Lambda) \leq Cl - k\epsilon < Cd_0 - \epsilon[d/d_0], \tag{7.1}$$

as claimed. A closely related property is that the negativity of a given graded symplectic automorphism depends only on the global Maslov class μ_M, and not on the choice of α_M within that class. For this take two cohomologous phase functions $\alpha'_M = e^{2\pi i\chi}\alpha_M$. Since χ is bounded, it follows from (6.15) that the gradings of ϕ^d with respect to α_M, α'_M differ by a uniformly bounded amount, which in view of (7.1) gives the desired property.

Lemma 7.2. *Let $\tilde{\phi}$ be a negative graded symplectic automorphism, and $\tilde{L}_0, \tilde{L}_1$ two graded Lagrangian submanifolds. For any $k \in \mathbb{Z}$ one can find a $d > 0$ and a small exact perturbation L_1' of L_1 such that for each $x \in L_0 \cap \phi^d(L_1')$, the absolute Maslov index is*

$$I(x) < k.$$

Proof. Since negativity is independent of the choice of phase function, we may assume that that function is classical. Suppose that the grading of L_1 is bounded above by C_1, and that of L_0 bounded below by C_0. Choose some d such that $C_1 - C_0 + Cd_0 - \epsilon[d/d_0] \leq k - n$. From (7.1) and (6.14) it follows that

$$\tilde{\alpha}_{\phi^d(L_1)}(x) - \tilde{\alpha}_{L_0}(x) < k - n \tag{7.2}$$

at every intersection point x. This estimate continues to hold if one perturbs L_1 slightly, to make the intersections transverse. Now apply Lemma 6.8. ☐

Negativity is well-behaved with respect to abelian symplectic reduction (this fact is related to the quotient construction of special Lagrangian submanifolds, see for instance [24]). The situation is as follows: $M = M^{2n}$ carries a free Hamiltonian T^k-action, with moment map $\mu : M \to \mathbb{R}^k$. We have an almost complex structure J_M which is ω_M-compatible and T^k-invariant, and a T^k-invariant complex volume form η_M, with associated (classical and T^k-invariant) phase function α_M. Denote by $K_1, \dots, K_k$ the Killing fields of the action. Let ϕ be a symplectic automorphism which preserves μ, hence is T^k-equivariant; and $\tilde{\alpha}_\phi$ a grading of it (necessarily T^k-invariant). For every moment value r we have the symplectic quotient $M^{\mathrm{red},r} = \mu^{-1}(r)/T^k$. This carries the reduced symplectic form $\omega_{M^{\mathrm{red},r}}$ and a compatible almost complex structure $J_{M^{\mathrm{red},r}}$, the latter of which is obtained by taking the complexified tangent spaces along the orbits, and identifying their orthogonal complement with the tangent spaces of $M^{\mathrm{red},r}$. Moreover, ϕ induces symplectic automorphisms $\phi^{\mathrm{red},r}$. In our situation, we also have a reduced complex volume form $\eta_{M^{\mathrm{red},r}}$, defined by restricting $i_{K_1} \dots i_{K_r} \eta_M$ to the same orthogonal complement as before. Projection identifies Lagrangian subspaces $\Lambda \subset TM$ containing the tangent space to the orbit with Lagrangian subspaces $\Lambda^{\mathrm{red}} \subset TM^{\mathrm{red},r}$. The phase functions defined by η_M, $\eta_{M^{\mathrm{red},r}}$ are related by $\alpha_{M^{\mathrm{red},r}}(\Lambda^{\mathrm{red}}) = \alpha_M(\Lambda)$. In particular, we can define a grading of the reduced symplectic automorphisms by

$$\tilde{\alpha}_{\phi^{\mathrm{red},r}}(\Lambda^{\mathrm{red}}) = \tilde{\alpha}_\phi(\Lambda). \tag{7.3}$$

Lemma 7.3. *Suppose that the $\tilde{\phi}^{\mathrm{red},r}$ are all negative. Then $\tilde{\phi}$ is negative.*

Proof. Choose some r. By assumption there is a d_0 such that the grading of $(\tilde{\phi}^{\mathrm{red},r})^{d_0}$ is negative, and then the same thing holds for neighbouring r. By a covering argument, one sees that there is a d_0 such that the gradings of the d_0-th iterates of all reduced maps are negative. Applying (7.1) shows that for some large d, the grading of each $(\tilde{\phi}^{\mathrm{red},r})^d$ will be $< -n$ everywhere. By (7.3), this means that at each point of M there are some Lagrangian tangent subspaces whose $\tilde{\phi}^d$-grading is $< -n$. From this and Lemma 6.9 one gets the desired conclusion. □

(7b) We will now look at the local models for the "large complex structure limit" monodromy, first under the rather unrealistic assumption that both the symplectic and complex structures are standard, which allows one to integrate the monodromy vector field explicitly.

Assumptions 7.4. *Fix* $n \geq 2$ *and* $2 \leq k \leq n$. *We take* $Y = \mathbb{C}^n$ *with the standard complex structure and symplectic form* ω_Y, *and the map*

$$p : \mathbb{C}^n \longrightarrow \mathbb{C}, \quad y \longmapsto y_1 y_2 \dots y_k.$$

Y should carry the complex volume form (away from the zero fibre)

$$\frac{dy_1}{y_1} \wedge \cdots \wedge \frac{dy_k}{y_k} \wedge dy_{k+1} \wedge \cdots \wedge dy_n.$$

The monodromy vector field Z is the unique horizontal lift of the clockwise rotational field $-i\zeta\partial_\zeta$. Explicitly,

$$Z = -i\frac{\left(\frac{y_1}{|y_1|^2}, \dots, \frac{y_k}{|y_k|^2}, 0, \dots, 0\right)}{\frac{1}{|y_1|^2} + \cdots + \frac{1}{|y_k|^2}},$$

In particular, since each $|y_j|^2$ is invariant, the flow ψ_t of Z is well-defined for all times, in spite of the noncompactness of the fibres. As in our discussion of Lefschetz pencils, we define complex volume forms η_{Y_z} on the fibres by (formally) dividing η_Y by some complex one-form on the base, which in this case will be $d\zeta/\zeta^2$. As a consequence, if at some point $y \in Y_\zeta = p^{-1}(\zeta)$ we have a Lagrangian subspace $\Lambda \subset TY_y$ with $Dp(\Lambda) = a\mathbb{R}$, $a \in S^1$, then the phase of the vertical part $\Lambda^v = \Lambda \cap \ker(Dp)$ inside Y_ζ is related to the phase of Λ by

$$\alpha_{Y_\zeta}(\Lambda^v) = \frac{\zeta^4}{a^2|\zeta|^4}\alpha_Y(\Lambda). \tag{7.4}$$

For each point y with $p(y) = \zeta \neq 0$, consider the Lagrangian tangent subspace $\Lambda_y = \mathbb{R}iy_1 \oplus \cdots \oplus \mathbb{R}iy_k \oplus \mathbb{R}^{n-k} \subset TY_y$. This satisfies $\alpha_Y(\Lambda_y) = (-1)^k$ and $Dp(\Lambda_y) = i\zeta\mathbb{R}$, hence (7.4) says that

$$\alpha_{Y_z}(\Lambda_y^v) = \frac{\zeta^2}{|\zeta|^2}(-1)^{k-1}. \tag{7.5}$$

ψ_t takes the Λ_y to each other, and rotates the base coordinate ζ, hence it also maps the Λ_y^v to each other. Let $h_\zeta = \psi_{2\pi}|Y_z : Y_z \to Y_z$ be the monodromy on the fibre over ζ. By definition and (7.5), the canonical grading of this map satisfies

$$\begin{aligned}\tilde{\alpha}_{h_\zeta}(\Lambda^v_y) &= \frac{1}{2\pi}\int_0^{2\pi} \frac{d}{dt}\arg(\alpha_{Y_z}(D\psi_t(\Lambda^v_y)))\,dt \\ &= \frac{1}{2\pi}\int_0^{2\pi} \frac{d}{dt}\arg((-1)^{k-1}e^{-2it})\,dt = -2.\end{aligned} \tag{7.6}$$

If one passes to a sufficiently large iterate such as h_ζ^n, the grading on Λ^v_y is $-2n$, hence by Lemma 6.9 the grading of any Λ is $< -n-1$. In a slight abuse of terminology (since Y_ζ is not compact), we conclude that the monodromy is a negative graded symplectic automorphism, in a way which is uniform for all ζ.

(7c) A different local model appears at the singular points of the base locus of our pencil. In Ruan's work on the SYZ conjecture, see in particular [54], these are responsible for the singularities of the Lagrangian torus fibration. For closely related reasons, the negativity of the monodromy breaks down at such points.

Assumption 7.5. *Fix $n \geq 3$ and $3 \leq k \leq n$. We again take $Y = \mathbb{C}^n$ with the standard complex and symplectic forms, but now with the rational map*

$$p : Y \dashrightarrow \mathbb{C}, \qquad p(y) = \frac{y_2 \cdots y_k}{y_1},$$

in the presence of the complex volume form (away from the closure of the zero fibre)

$$\eta_Y = dy_1 \wedge \frac{dy_2}{y_2} \wedge \cdots \wedge \frac{dy_k}{y_k} \wedge dy_{k+1} \wedge \cdots \wedge dy_n.$$

Denote by Y_ζ the closure of $p^{-1}(y)$. For $\zeta \neq 0$, each such fibre can be identified with $\mathbb{C}^{n-1}$ by projecting to $(y_2, \dots, y_n)$. We will use this identification freely, hoping that this does not cause too much confusion. For instance, the symplectic form on $Y_\zeta \cong \mathbb{C}^{n-1}$ is given by

$$\omega_{Y_\zeta} = \frac{i}{2}\partial\bar{\partial}\Big(\frac{|y_2 \dots y_k|^2}{|\zeta|^2} + |y_2|^2 + \cdots + |y_n|^2\Big) \tag{7.7}$$

As in our previous model, we write Z for the clockwise monodromy vector field (defined by passing to the graph of p). In view of our identification, this becomes a ζ-dependent vector field on $\mathbb{C}^{n-1}$, namely:

$$Z = -i\frac{\Big(\frac{y_2}{|y_2|^2}, \dots, \frac{y_k}{|y_k|^2}, 0, \dots, 0\Big)}{\frac{1}{|y_1|^2} + \frac{1}{|y_2|^2} + \cdots + \frac{1}{|y_k|^2}}$$

where $y_1 = y_2 \dots y_k/\zeta$. Again, this preserves $|y_2|^2, \dots, |y_n|^2$, so that the flow ψ_t is well-defined for all times.

Taking η_Y on the total space and $d\zeta/\zeta^2$ on the base, one finds that the induced volume form η_{Y_ζ} is the standard form on $\mathbb{C}^{n-1}$. At each point $y \in Y_\zeta$ with

$y_2 \dots y_k \neq 0$, consider the Lagrangian subspace $\Lambda^v_y = \mathbb{R}iy_2 \oplus \cdots \oplus \mathbb{R}iy_k \oplus \mathbb{R}^{n-k}$. These subspaces are preserved by the flow ψ_t, and a computation similar to (7.6) shows that the canonical grading of the monodromy map $h_\zeta = \psi_{2\pi}|Y_\zeta$ satisfies

$$\tilde{\alpha}_{h_\zeta}(\Lambda^v_y) = -2\frac{1}{1+\frac{1}{\frac{|y_1|^2}{|y_2|^2}+\cdots+\frac{|y_1|^2}{|y_k|^2}}}.$$

This is obviously negative, but of course we also need to consider the grading at the points where Λ^v_y is not defined. Suppose for simplicity that the sizes of the first $k-1$ coordinates of $y \in Y_\zeta$ are ordered, $|y_2| \leq |y_3| \leq \dots |y_{k-1}|$. Then

$$\tilde{\alpha}_{h_\zeta}(\Lambda_y) \leq -2\frac{1}{1+\frac{|\zeta|^2}{|y_3|^{2(k-1)}}},$$

which means that the grading is bounded above by something which depends on the size of the second smallest coordinate. As a consequence, if one has a sequence

$$y^{(m)} \to y^{(\infty)}, \quad y_2^{(m)} \dots y_k^{(m)} \neq 0 \text{ for all } m,$$

and such that only one of the $(y_2^{(\infty)}, \dots, y_k^{(\infty)})$ vanishes, the $\tilde{\alpha}_{h_\zeta}(\Lambda^v_{y^{(m)}})$ are bounded above by a negative constant, hence cannot go to zero. Roughly speaking, this means that h_ζ is negative away from the subset where at least two of the $y_2, \dots, y_k$ vanish. In the special case $k = n = 3$, this subset is a point $0 \in Y_\zeta$; one can then apply (6.14) and Lemma 6.9 much as in our previous local model, to get the following statement:

Lemma 7.6. *Suppose that $k = n = 3$. Let $W \subset Y_\zeta$ be some open ball around the origin. Then $h_\zeta(W) = W$, and there is a $d > 0$ such that the grading of $h^d_\zeta|(Y_\zeta \setminus W)$ is negative.* ☐

Remark 7.7. *As mentioned before, ideas associated to the SYZ conjecture suggest that the monodromy should be a fibrewise translation in a SLAG fibration. From this point of view, the negativity of the monodromy is based on the idea that the phases of the tangent spaces to the fibration should be shifted by -2. In the situation of Assumptions 7.4, this is indeed the case: $L_r \cong T^{k-1} \times \mathbb{R}^{n-k} \subset Y_\zeta$, given by*

$$L_r = \{|y_1|^2 = r_1, \dots, |y_k|^2 = r_k, \mathrm{im}(y_{k+1}) = r_{k+1}, \dots, \mathrm{im}(y_n) = r_n\}$$

with $r_1 \dots r_k = |\zeta|^2$, are special Lagrangian with respect to η_{Y_ζ}. The monodromy translates each of them, and the shift is computed in (7.6). *For Assumptions 7.5, there is a standard SLAG fibration with singularities on $Y_\zeta \cong \mathbb{C}^{n-1}$ (with respect to* (7.7) *and the standard complex volume form), see e.g.* [24]*:*

$$L_r = \{|y_3|^2 - |y_2|^2 = r_1, \dots |y_k|^2 - |y_2|^2 = r_{k-2}, \\ \mathrm{im}(y_2 \dots y_k) = r_{k-1}, \mathrm{im}(y_{k+1}) = r_k, \dots, \mathrm{im}(y_n) = r_{n-1}\}. \tag{7.8}$$

However, the monodromy preserves this only asymptotically far away from the singular locus, and hence the grading is only asymptotically equal to -2. *To see more precisely what happens, we apply a symplectic reduction procedure, which means mapping* $Y_\zeta \rightarrow \mathbb{C}$ *by sending* y *to* $w = y_2 y_3 \dots y_k$. *The fibres of the SLAG fibration lie over horizontal lines in the base* $\mathbb{C}$. *The clockwise monodromy also fibres over a diffeomorphism of* $\mathbb{C}$, *which rotates each point* w *by some* $|w|$*-dependent angle. For* $|w| \gg 0$, *this is almost a full clockwise rotation, hence it approximately preserves the horizontal lines, but near* $w = 0$ *the angle of rotation undergoes a full change* 0 *to* -2π.

On the other hand, as pointed out to me by Mark Gross, in the $K3$ *case one can construct a symplectic automorphism which preserves all the fibres of a SLAG fibration. This is obtained by hyperkähler rotation from a complex automorphism, and it is presumably isotopic to the monodromy map, even though that has not strictly speaking been proved. So, it may well be that the failure of our local models to preserve the fibrations* (7.8), *and the corresponding breakdown in negativity, could be repaired by an isotopy which is more global in nature.*

(7d) We now relax our previous assumptions on the complex and symplectic structures, for the first local model. We choose to work in Darboux coordinates, and with a complex structure and holomorphic function which are not standard, but which still preserve a T^k symmetry.

ASSUMPTIONS 7.8. *Fix* $n \geq 2$ *and* $2 \leq k \leq n$.

- $Y \subset \mathbb{C}^n$ *is an open ball around the origin, carrying the standard symplectic form* ω_Y *and the standard diagonal* T^k*-action*
$$\rho_s(y) = (e^{is_1} y_1, \dots, e^{is_k} y_k, y_{k+1}, \dots, y_n),$$
with moment map $\mu : Y \rightarrow (\mathbb{R}^{\geq 0})^k$. *For any regular moment value* r, *the quotient* $Y^{\mathrm{red},r}$ *can be identified with an open subset of* $\mathbb{C}^{n-r}$, *with the standard symplectic form* $\omega_{Y^{\mathrm{red},r}}$. *We denote points in the reduced spaces by* $y^{\mathrm{red}} = (y_{k+1}, \dots, y_n)$.
- J_Y *is a complex structure which is tamed by* ω_Y. *At the point* $y = 0$ *(but not necessarily elsewhere), it is* ω_Y*-compatible and* T^k*-invariant.*
- $p : Y \rightarrow \mathbb{C}$ *is a* J_Y*-holomorphic function with the following properties:*
  (i) $p(\rho_s(y)) = e^{i(s_1 + \cdots + s_k)} p(y)$.
  (ii) $\partial_{y_1} \dots \partial_{y_k} p$ *is nonzero at the point* $y = 0$.
- η_Y *is a* J_Y*-complex volume form on* $Y \setminus p^{-1}(0)$, *such that* $p(y)\eta_Y$ *extends to a smooth form on* Y, *which is nonzero at* $y = 0$. *Let* α_Y *be the corresponding phase function.*

Lemma 7.9. *One can write*

$$p(y) = 2^{-k/2} y_1 y_2 \dots y_k \, q(\tfrac{1}{2}|y_1|^2, \dots, \tfrac{1}{2}|y_k|^2, y_{k+1}, \dots, y_n)$$

for some smooth function q *satisfying* $q(0) \neq 0$.

Proof. Consider the Taylor expansion of p around $y = 0$. From property (i) of p one sees that each monomial which occurs with nonzero coefficient in that expansion contains one of the factors y_l for each $1 \leq l \leq k$. As a consequence, $p(y)/(y_1 \dots y_k)$ extends smoothly over $y = 0$. The same argument can be applied to points where only certain of the coordinates $y_1 \dots y_k$ vanish. Again applying (i), one sees that $p(y)/(y_1 \dots y_k)$ is T^k-invariant, which means that it can be written in the form stated above. Finally, $q(0) \neq 0$ is property (ii). □

Lemma 7.10. *After making* Y *smaller if necessary, one has*

$$p(y) = 0 \iff y_l = 0 \text{ for some } l = 1, \dots, k;$$
$$Dp(y) = 0 \iff y_l = 0 \text{ for at least two } l = 1, \dots, k.$$

Proof. Suppose that $q(y) \neq 0$ on the whole of Y. Then the first statement is obvious, and so is the $\Leftarrow$ implication in the second one. Conversely, suppose that y is a critical point. Property (i) implies that $p(y) = 0$, so that $y_l = 0$ for some $l = 1, \dots, k$. From $0 = Dp_y = y_1 \dots \hat{y}_l \dots y_k \, q(y) \, dy_l$ one sees that another of the $y_1, \dots, y_k$ has to vanish too. □

We will assume from now on that the conclusions of Lemma 7.10 hold. Consider the function $H(y) = -\frac{1}{2}|p(y)|^2$, its Hamiltonian vector field X and flow ϕ_t. Since H is T^k-invariant, the flow (where defined) is equivariant and preserves the level sets of the moment map. For every regular moment value r, we can consider the induced function

$$H^{\mathrm{red},r}(y^{\mathrm{red}}) = -\tfrac{1}{2} r_1 \dots r_k \, |q(r_1, \dots, r_k, y_{k+1}, \dots, y_n)|^2, \tag{7.9}$$

on $Y^{\mathrm{red},r}$, its vector field $X^{\mathrm{red},r}$ and flow $\phi_t^{\mathrm{red},r}$. If the moment value is small, so are (7.9) and all its y^{red}-derivatives, which means that the reduced flow moves very slowly. We need a quantitative version of this:

Lemma 7.11. *Given* $\epsilon_1, \epsilon_2 > 0$, *there are* $\delta_1, \delta_2, \delta_3 > 0$ *such that for all*

$$\begin{cases} r \in (\mathbb{R}^{>0})^k & \text{with } ||r|| < \delta_1, \\ y^{\mathrm{red}} \in Y^{\mathrm{red},r} & \text{with } ||y^{\mathrm{red}}|| < \delta_2, \\ t \in \mathbb{R} & \text{with } |t| < \delta_3 r_1^{-1} \dots r_k^{-1}, \end{cases}$$

the following holds: $\phi_t^{\mathrm{red},r}$ *is well-defined near* y^{red}, *and satisfies*

$$||\phi_t^{\mathrm{red},r}(y^{\mathrm{red}})|| < \epsilon_1, \tag{7.10}$$

$$||(D\phi_t^{\mathrm{red},r})_{y^{\mathrm{red}}} - \mathrm{Id}|| < \epsilon_2. \tag{7.11}$$

PROOF. By Lemma 7.9 there are $\delta_1 > 0$, $\delta_2 \in (0; \min(\epsilon_1/2, \epsilon_2/2))$ and a $C > 0$, such that

$$||X^{\mathrm{red},r}_{y^{\mathrm{red}}}||,\ ||DX^{\mathrm{red},r}_{y^{\mathrm{red}}}|| \leq Cr_1 \dots r_k$$

for each $||r|| < \delta_1$, $||y^{\mathrm{red}}|| < 2\delta_2$ (we use the Hilbert-Schmidt norm for matrices $DX^{\mathrm{red},r}$, normalized to $||\mathrm{Id}|| = 1$). We claim that one can take $\delta_3 = \delta_2/C$. The first estimate (7.10) is clear: if one starts at a point y^{red} of norm less than δ_2, and moves with speed at most $Cr_1 \dots r_k$ for a time less than $\delta_2 C^{-1} r_1^{-1} \dots r_k^{-1}$, the endpoint will be of norm less than $2\delta_2 < \epsilon_1$. Next, use the linearized flow to get an exponential growth bound for the time interval t relevant to us: $||D\phi^{\mathrm{red},r}_t|| \leq \exp(Cr_1 \dots r_k t) \leq \exp(\delta_2)$. One may assume that $\exp(\delta_2) \leq 2$. By plugging this back into the linearized equation, one gets the linear estimate $||D\phi^{\mathrm{red},r}_t - \mathrm{Id}|| \leq 2Cr_1 \dots r_k t \leq 2\delta_2 < \epsilon_2$, from which (7.11) follows. □

We now need to deal with the fact that J_Y and η_Y are only approximately T^k-invariant. Let J'_Y be the constant complex structure on Y which agrees with J_Y at $y = 0$. This is ω_Y-compatible and T^k-invariant, so

$$J'_Y = \begin{pmatrix} i \cdot 1_k & 0 \\ 0 & J'_{Y^{\mathrm{red}}} \end{pmatrix}.$$

$J'_{Y^{\mathrm{red}}}$ is the induced complex structure on $Y^{\mathrm{red},r}$ for all r. Similarly, we consider the J'_Y-complex volume form η'_Y which is obtained by taking the value of $p(y)\eta_Y$ at the point $y = 0$, and dividing that by $y_1 \dots y_k$. This can be written as

$$\eta'_Y = i^k \frac{dy_1}{y_1} \wedge \dots \wedge \frac{dy_k}{y_k} \wedge \eta'_{Y^{\mathrm{red}}},$$

hence is T^k-invariant. $\eta'_{Y^{\mathrm{red}}}$, which is the induced complex volume form on the reduced spaces, is again constant and r-independent. Let α'_Y be the phase function associated to (J'_Y, η'_Y) (unlike α_Y, this is a classical phase function), and $\alpha'_{Y^{\mathrm{red},r}}$ the induced phase functions on the quotients.

LEMMA 7.12. *Given $\epsilon_3 > 0$, there is $\delta_4 > 0$ such that for all $||y|| < \delta_4$ with $p(y) \neq 0$, and all $\Lambda \in \mathfrak{L}_{Y,y}$,*

$$\Big|\frac{1}{2\pi} \arg(\alpha'_Y(\Lambda)/\alpha_Y(\Lambda))\Big| < \epsilon_3$$

(by this we mean that there is a branch of the argument with that property).

PROOF. The quotient $\alpha'_Y(\Lambda)/\alpha_Y(\Lambda)$ does not change if we multiply both η_Y, η'_Y with $y_1 \dots y_k$. But by construction,

$$y_1 \dots y_k \eta_Y = 2^{-k/2} q(\tfrac{1}{2}|y_1|^2, \dots, \tfrac{1}{2}|y_k|^2, y_{k+1}, \dots, y_n)^{-1} \sigma_y,$$
$$y_1 \dots y_k \eta'_Y = 2^{-k/2} q(0)^{-1} \sigma_0$$

where σ is some smooth n-form with $\sigma_0 \neq 0$, so the associated phases also become close as $y \to 0$. □

ϕ_t has a canonical grading $\tilde{\alpha}_{\phi_t}$ with respect to α_Y, obtained by starting with the trivial grading for ϕ_0 and extending continuously. Denote by $\tilde{\alpha}'_{\phi_t}$ the grading obtained in the same way, but with respect to α'_Y. There are corresponding gradings $\tilde{\alpha}'_{\phi_t^{\mathrm{red},r}}$ on the reduced spaces, with respect to $\alpha'_{Y^{\mathrm{red},r}}$.

Lemma 7.13. *Given $\epsilon_4 > 0$, one can find $\delta_5, \delta_6, \delta_7 > 0$ such that for all*

$$\begin{cases} r \in (\mathbb{R}^{>0})^k & \textit{with } ||r|| < \delta_5, \\ y^{\mathrm{red}} \in Y^{\mathrm{red},r} & \textit{with } ||y^{\mathrm{red}}|| < \delta_6, \\ \Lambda^{\mathrm{red}} \in \mathfrak{L}_{Y^{\mathrm{red},r},y^{\mathrm{red}}} & \textit{and} \\ t \in \mathbb{R} & \textit{with } |t| < \delta_7 r_1^{-1} \dots r_k^{-1}, \end{cases}$$

$\phi_t^{\mathrm{red},r}$ is well-defined near y^{red}, and its grading is $|\tilde{\alpha}'_{\phi_t^{\mathrm{red},r}}(\Lambda^{\mathrm{red}})| < \epsilon_4$.

Proof. This is an easy consequence of Lemma 7.11: one can achieve that the derivative $(D\phi_t^{\mathrm{red},r})_{y^{\mathrm{red}}}$ remains very close to the identity in the whole range of t's that we are considering. Since the volume form $\eta_{Y^{\mathrm{red},r}}$ is constant and independent of r, it follows that the grading remains small. □

Lemma 7.14. *Given $\epsilon_5 > 0$, one can find $\delta_8, \delta_9, \delta_{10} > 0$ such that for all*

$$\begin{cases} y \in Y & \textit{with } 0 < |p(y)| < \delta_8 \textit{ and } ||y|| < \delta_9, \\ \Lambda \in \mathfrak{L}_{Y,y}, & \textit{and} \\ t \in \mathbb{R} & \textit{with } |t| < \delta_{10}|p(y)|^{-2}, \end{cases}$$

ϕ_t is well-defined near y, and its grading satisfies $|\tilde{\alpha}_{\phi_t}(\Lambda)| < n + \epsilon_5$.

Proof. Apply Lemma 7.11 to the corresponding point y^{red} in the quotient. In particular, one can choose $\delta_8, \delta_9, \delta_{10}$ in such a way that the reduced flow exists near y^{red} for $|t| < \delta_{10}|p(y)|^{-2}$. Since the fibres of the quotient map are compact (tori), it follows that ϕ_t is well-defined near y in the same range of t. We now apply the same basic reasoning as in Lemma 7.3. After possibly making δ_8, δ_9 smaller, Lemma 7.13 ensures that there is a $\Lambda \in \mathfrak{L}_{Y,y}$ such that $|\tilde{\alpha}'_{\phi^t}(\Lambda)| < \epsilon_5/2$. By Lemma 6.9, the grading is $< n + \epsilon_5/2$ for all other Lagrangian subspaces at the same point. Finally, one uses Lemma 7.12 to pass from the grading associated to η'_Y to that for η_Y, again making the δs smaller so that the phase difference becomes $< \epsilon_5/2$. □

The next step is to relate ϕ_t to the monodromy. As usual, we take the vector field $-i\zeta\partial_\zeta$ on $\mathbb{C}^*$, and lift it in the unique way to a horizontal vector field Z on $Y \setminus p^{-1}(0)$. This is well-defined because ω_Y tames J_Y, hence the regular fibres Y_ζ, $\zeta \neq 0$, are symplectic submanifolds. We claim that

$$Z = fX \tag{7.12}$$

for some positive function f on $Y \setminus p^{-1}(0)$. First, because H is constant on each fibre, the associated vector field X is horizontal. Secondly, because $\omega_Y(X, Z) = 0$ and the space of horizontal vectors TY^h is two-dimensional, it is true that $X = f^{-1}Z$ for some f. A little more thought shows that

$$f = \frac{\omega_Y | TY^h}{p^*\omega_{\mathbb{C}} | TY^h} > 0, \tag{7.13}$$

where TY^h is the ω_Y-horizontal subspace (which is two-dimensional, so any two two-forms on it differ by a scalar). Let (ψ_s) be the flow of Z, so that $h_\zeta = \psi_{2\pi}|Y_\zeta$ is the clockwise monodromy (of course, like ϕ_s this flow is only partially defined). As a consequence of (7.12), the unparametrized flow lines of ϕ_t and ψ_t agree, so

$$\psi_t(y) = \phi_{g_t(y)}(y), \quad \text{where} \quad g_t(y) = \int_0^t f(\psi_\tau(y))d\tau. \tag{7.14}$$

Lemma 7.15. *For every $d > 0$, $\epsilon_6 > 0$ there are $\delta_{11}, \delta_{12} > 0$ such that the following holds. For all $0 < |\zeta| < \delta_{11}$ and all $y \in Y_\zeta$ with $||y|| < \delta_{12}$, the d-fold monodromy h_ζ^d is well-defined, and (7.14) holds for $t = 2\pi d$ with*

$$g_{2\pi d}(y) \leq \frac{\epsilon_6}{|\zeta|^2}.$$

Proof. Using property (i) of p and its J_Y-holomorphicity, one obtains that $Dp_y(J_Y(-iy_1, 0, \dots, 0)) = p(y)$ and hence $dH_y(J_Y(-iy_1, 0, \dots, 0)) = |p(y)|^2 = |\zeta|^2$. It follows that for small y, $||X_y|| \geq c_1|\zeta|^2/||y||$ for some $c_1 > 0$. Hence there is a $C_2 > 0$ such that

$$C_2^{-1}|\zeta|^4/||y||^2 \leq \omega_Y(X, J_Y X) = f^{-2}\omega_Y(Z, J_Y Z).$$

On the other hand, $\omega_Y(Z, J_Y Z) = |\zeta|^2 f$ by definition of Z and (7.13), so finally,

$$f(y) \leq C_2 \frac{||y||^2}{|\zeta|^2}.$$

Integrating gives a similar inequality $g_{2\pi d}(y) \leq C_3||y||^2/|\zeta|^2$. Clearly, by restricting to smaller y one can make the right hand side less than $\epsilon_6/|\zeta|^2$ for any desired ϵ_6 (this, by the way, is the crucial estimate in our whole computation). □

We use J_Y-complex volume forms η_{Y_ζ} on the fibres obtained in the usual way from η_Y, so that (7.4) holds. Take $d > 0$, $y \in Y_\zeta$, and a Lagrangian subspace $\Lambda \subset TY_y$ with $Dp(\Lambda) = i\zeta\mathbb{R}$. This is necessarily of the form $\Lambda = \Lambda^v \oplus \mathbb{R}Z_y$ for some Lagrangian subspace $\Lambda^v \subset T_y(Y_\zeta)$. By (7.12) and (7.14) one has

$$D(\psi_t)_y = D(\phi_{g_t})_y + X_{\psi_t(y)} \otimes dg_t = D(\phi_{g_t})_y + Z_{\psi_t(y)} \otimes \frac{dg_t}{f(\psi_t(y))} \tag{7.15}$$

Since Λ contains Z_y, and ψ_t is the clockwise parallel transport flow, $D(\psi_t)_y(\Lambda)$ contains $Z_{\psi_t(y)}$. From this and (7.15) it follows that

$$D\psi_t(\Lambda) = D(\phi_{g_t})(\Lambda), \qquad D(\psi_t|Y_\zeta)(\Lambda^v) = D(\phi_{g_t})(\Lambda) \cap \ker(Dp),$$

and hence by (7.4) that $\alpha_{\psi_t|Y_\zeta}(\Lambda^v) = e^{-4\pi i t}\alpha_{\phi_{g_t}}(\Lambda)$. By considering the behaviour of this over the time $t \in [0; 2\pi d]$ one finds that the canonical grading of the monodromy satisfies

$$\tilde{\alpha}_{h^d_\zeta}(\Lambda^v) = \tilde{\alpha}_{\phi_{g_{2\pi d}(y)}}(\Lambda) - 2d. \tag{7.16}$$

Lemma 7.16. *For every $d > 0$ and $\epsilon_7 > 0$ there are $\delta_{13}, \delta_{14} > 0$ such that the following holds. For every $y \in Y_\zeta$ with $0 < |\zeta| < \delta_{13}$ and $||y|| < \delta_{14}$, and every Lagrangian subspace $\Lambda^v \subset T(Y_\zeta)_y$, we have that h^d_ζ is well-defined near y, and its canonical grading satisfies*

$$\tilde{\alpha}_{h^d_\zeta}(\Lambda^v) < n - 2d + \epsilon_7.$$

Proof. Lemma 7.14 says that one can choose δ_{13}, δ_{14} so small that there is a $\delta_{15} > 0$ with

$$|\tilde{\alpha}_{\phi_t}(\Lambda)| < n + \epsilon_7 \tag{7.17}$$

for all $|t| < \delta_{15}/|p(y)|^2$. On the other hand, after possibly making δ_{13}, δ_{14} smaller, one knows from Lemma 7.15 that $|g_{2\pi d}| < \delta_{15}/|p(y)|^2$, so (7.17) applies to the values of t that are relevant for our monodromy map. (7.16) completes the proof. □

This shows that the basic feature of our first local model, the uniform negativity of the monodromy, survives even if one generalizes from the idealized situation which we considered first (Assumptions 7.4) to a more realistic one (Assumptions 7.8). In principle, one should carry out a parallel discussion for the second local model. This would be rather more delicate, since the grading becomes zero along a submanifold; an arbitrary small perturbation can potentially make it positive, thereby spoiling the entire argument. Fortunately, in the $K3$ case the second model occurs only at isolated points, and one can achieve that the local picture near those points is standard (satisfies Assumptions 7.5), so that the previous discussion suffices after all.

(7e) We will now explain how to patch together the local models to understand of the "large complex structure limit" monodromy. This is somewhat similar to the strategy used by Ruan to construct (singular) Lagrangian torus fibrations [54]. We consider a Fano threefold X with $o_X = \mathcal{K}_X^{-1}$, and a pair of sections $\sigma_{X,0}, \sigma_{X,\infty}$ which generate a quasi-Lefschetz pencil of hypersurfaces $\{X_z\}$. From $\sigma_{X,\infty}$ we get a rational complex volume form η_X with a pole along X_∞. There are canonical induced complex volume forms η_{X_z} on the fibres X_z (if that is smooth and $z \neq \infty$), which are therefore Calabi-Yau surfaces. In fact, they must be $K3$s, since $H^1(X) = 0$ for any Fano variety, hence $H^1(X_z) = 0$ by the Lefschetz hyperplane theorem.

Lemma 7.17. *One can find a Kähler form ω'_X in the same cohomology class as ω_X, with the following properties. (i) Near every point where three components of X_∞ meet, $(X, \omega'_X, 1/\pi)$ is modelled on (Y, ω_Y, p) from Assumptions 7.4, with $n = k = 3$. (ii) Near every point where X_0 meets two components of X_∞, $(X, \omega'_X, 1/\pi)$ is modelled on (Y, ω_Y, p) from Assumptions 7.5, with $n = k = 3$.*

This is easy because the set of such points (which we will call "lowest stratum points" in future) is finite. Near each of them, there are local holomorphic coordinates in which $(X, 1/\pi)$ becomes (Y, p). One can then locally modify the Kähler form to make it standard, see e.g. [54, Lemma 7.3] or [60, Lemma 1.7].

Lemma 7.18. *One can find a Kähler form ω''_X in the same cohomology class as ω_X, which agrees with ω'_X in a neighbourhood of the lowest stratum points, and such that in addition, any two components of X_∞ meet orthogonally.* □

Ruan proves a much more general result in [54, Theorem 7.1]. The main idea is to apply the techniques from the previous Lemma to each fibre of the normal bundle ν to the intersection of two components of X_∞.

Before proceeding, we need some more notation. Denote by C the irreducible components of X_∞, and by $L_C = \mathcal{O}(-C)$ the corresponding line bundles. Let γ_C be their canonical sections, with a zero along C. Take two components $C \neq C'$, and choose a small neighbourhood $U_{C,C'}$ of $C \cap C'$. On this consider the vector bundle

$$N_{C,C'} = L_C \oplus L_C^{-1},$$

We want to equip L_C and hence $N_{C,C'}$ with hermitian metrics and the corresponding connections, with the following prescribed behaviour near the lowest stratum pointsin $U_{C,C'}$:

- If x is a point where $C \cap C'$ meets another component C'', we choose local coordinates for X as given by Lemma 7.17, so that Assumptions 7.4 hold, and a local trivialization of L_C such that $\gamma_C(y) = y_1$. Take the trivial metric on L_C with respect to this trivialization.

- If x is a point where $C \cap C'$ meets X_0, we choose local coordinates for X as given by Lemma 7.17, so that Assumptions 7.5 hold, and a local trivialization of L_C such that $\gamma_C(y) = y_2$. Again, we take the trivial metric.

Take the Kähler metric associated to ω''_X, and use its exponential map to define a retraction $r_{C,C'} : U_{C,C'} \to C \cap C'$ of an open neighbourhood onto $C \cap C'$.

A corresponding use of parallel transport on $N_{C,C'}$, with respect to our chosen connection, yields an isomorphism $R_{C,C'} : N_{C,C'}|U_{C,C'} \to r^*_{C,C'}(N_{C,C'}|C \cap C')$. Taking the two objects together, we have a map

$$\nu_{C,C'} : U_{C,C'} \setminus X_0 \xrightarrow{\xi_{C,C'}=(\gamma_C,\pi_M^{-1}\gamma_C^{-1})} N_{C,C'}|U_{C,C'} \xrightarrow{R_{C,C'}} N_{C,C'}|C \cap C'.$$

Lemma 7.19. *After making $U_{C,C'}$ smaller if necessary, it carries a unique T^2-action $\sigma_{C,C'}$ which, under $\nu_{C,C'}$, corresponds to the obvious fibrewise T^2-action on $N_{C,C'}|C \cap C'$. In particular,*

$$\pi_M(\sigma_{C,C',s}(x)) = e^{-i(s_1+s_2)}\pi_M(x). \tag{7.18}$$

Moreover, at each point of $C \cap C'$, the action preserves ω''_X and the complex structure; the same holds in a neighbourhood of the lowest stratum points.

Proof. It is sufficient to verify the existence and uniqueness statement locally. First, take a point $x \in C \cap C'$ which is not a lowest stratum point. γ_C^{-1} has a simple pole along C, and π_M^{-1} has a simple zero along $C \cup C'$, hence $\pi_M^{-1}\gamma_C^{-1}$ has a simple zero along C'. It follows that $\xi_{C,C'}$ is transverse to the zero-section at x, so $\nu_{C,C'}$ is a local diffeomorphism; we simply pull back the T^2-action by it. By definition of the connection on $N_{C,C'}$, there is a commutative diagram

$$\begin{tikzcd}
U_{C,C'} \setminus X_0 \arrow[r, "\xi_{C,C'}"] \arrow[dr, "\pi_M^{-1}"'] & N_{C,C'}|U_{C,C'} \arrow[r, "R_{C,C'}"] \arrow[d] & N_{C,C'}|C \cap C' \arrow[dl] \\
& \mathbb{C} &
\end{tikzcd}$$

where the unlabeled arrows are the canonical pairings $L_C \oplus L_C^{-1} \to \mathbb{C}$. This shows that our pullback T^2-action satisfies (7.18). At the point x itself,

$$D\nu_{C,C'} : TX_x \longrightarrow T(C \cap C')_x \oplus N_x \tag{7.19}$$

is given by orthogonal projection to $C \cap C'$ with respect to the $o_{X,2}$-metric, together with $D\xi_x : TX_x \to N_x$. This is $\mathbb{C}$-linear, which implies that $\sigma_{C,C',x}$ preserves the complex structure. As for the symplectic structure, observe that by Lemma 7.18 TX_x splits into three complex one-dimensional pieces, namely $T(C \cap C')_x$ and the normal directions to that inside C, C'. On the right hand side of (7.19), this corresponds to the splitting of N_x into line bundles. The symplectic structures on the each summand do not correspond, but they differ only by a scalar, which is enough to conclude that $\sigma_{C,C'}$ is symplectic at x.

It remains to deal with the situation near the lowest stratum points. For this, we choose local coordinates and the trivialization of $N_{C,C'}$ as arranged before. Near a point of $C \cap C' \cap C''$, one then finds that $\nu_{C,C'}(y) = (y_3, y_1, y_2y_3)$, where the first component is the coordinate on $C \cap C'$. Hence, our $\sigma_{C,C'}$ near that point is the standard T^2-action in the variables y_1, y_2. Similarly, near a point of

$C \cap C' \cap X_0$, $\nu_{C,C'}(y) = (y_1, y_2, y_3/y_1)$, so that one ends up with the T^2-action in the variables y_2, y_3. These obviously have the desired properties. □

We now pick an ordering of every pair $\{C, C'\}$ of components, and will use only the T^2-actions $\sigma_{C,C'}$ in that order.

Lemma 7.20. *There is a symplectic form ω_X''' which tames the complex structure, such that for each intersection $C \cap C'$ of two components of X_∞, there is a neighbourhood $U_{C,C'}$ such that the local T^2-action $\rho_{C,C'}$ leaves ω_X''' invariant. Moreover, ω_X''' lies in the same cohomology class as ω_X'', and agrees with it at each point $x \in C \cap C'$, as well as in a neighbourhood of the lowest stratum points.*

Proof. Let $\bar{\omega}_{C,C'} \in \Omega^2(U_{C,C'})$ be the result of averaging ω_X'' with respect to $\rho_{C,C'}$. This is a closed two-form; it agrees with ω_X'' along $CcapC'$, and also in a neighbourhood of the lowest stratum points. Using the Poincaré lemma one can write $\bar{\omega}_{C,C'} - \omega_X'' = d\beta_{C,C'}$, where $\beta_{C,C'}$ is a one-form that vanishes to second order along $C \cap C'$, and which is again zero in a neighbourhood of the lowest stratum point. Take a compactly supported cutoff function $\psi_{C,C'}$ on $U_{C,C'}$, which is equal to one near $C \cap C'$. One sees easily that as one rescales the normal directions to make the support of $\psi_{C,C'}$ smaller, $d(\psi_{C,C'}\beta_{C,C'})$ becomes arbitrarily small in C^0-sense. It follows that for sufficiently small support, $\omega_X'' + d(\psi_{C,C'}\beta_{C,C'})$ is a symplectic form which tames the complex structure. This agrees with ω_X'' near the lowest stratum points, and for that reason, one can carry out the construction simultaneously for all intersections $C \cap C'$. □

Lemma 7.21. *Let $x \in C \cap C'$ be a point where two components of X_∞ meet, which is not a lowest stratum point. Then there are local $\rho_{C,C'}$-equivariant Darboux coordinates near x, in which X with the map $1/\pi$, the symplectic structure ω_X''', the given complex structure J_X, and the meromorphic complex volume form η_X, is isomorphic to $(Y, p, \omega_Y, J_Y, \eta_Y)$ as in Assumptions 7.8.* □

This is straightforward checking, which we leave to the reader. The outcome of the whole construction is this:

Proposition 7.22. *Let $\gamma_\infty : [0; 2\pi] \to \mathbb{C}$ be a circle of large radius $R \gg 0$. Then the graded monodromy*

$$\tilde{h}_{\gamma_\infty} \in \widetilde{Aut}(M_R)$$

is isotopic (within that group) to a graded symplectic automorphism $\tilde{\phi}$ with the following property. There is a finite set $\Sigma \subset X_z \cap X_\infty$ which admits arbitrarily small open neighbourhoods $W \subset X_z$, such that $\phi(W) = W$ and $\tilde{\phi}|(X_z \setminus W)$.

Proof. The main point is to replace the given Kähler form ω_X by ω_X''', and we need to convince ourselves that this is permitted. For each regular X_z there is a symplectic isomorphism

$$\phi_z : (X_z, \omega_{X_z}) \longrightarrow (X_z, \omega_X'''|X_z)$$

which carries $X_{0,\infty}$ to itself. This follows from Moser's Lemma together with the isotopy theorem for symplectic surfaces with orthogonal normal crossings, see [54, Theorem 6.5] for a more general statement valid in all dimensions. Next, an inspection of the construction of monodromy maps for π_M shows that they can be defined using any symplectic form which tames the complex structure, and in particular ω_X'''. The resulting diagram of symplectic maps

$$\begin{tikzcd}
(X_R, \omega_{X_R}) \arrow[r, "\phi_R"] \arrow[d, "h_{\gamma_\infty}"] & (X_R, \omega_X'''|X_R) \arrow[d, "h'''_{\gamma_\infty}"] \\
(X_R, \omega_{X_R}) \arrow[r, "\phi_R"] & (X_R, \omega_X'''|X_R)
\end{tikzcd}$$

commutes up to symplectic isotopy rel $X_{R,\infty}$. Since X_R is a $K3$ surface, Remark 6.3 shows that h_{γ_∞} is isotopic to $\phi_R^{-1} \circ h'''_{\gamma_\infty} \circ \phi_R$ within $Aut(M_R)$. It is no problem to take the gradings into consideration. Hence, if h'''_{γ_∞} has the desired negativity property for large R, so will h_{γ_∞}.

Let Σ be the set of those points where two components of X_∞ intersect each other and X_0. This lies in the base locus of our pencil, hence in every fibre X_z. Around each point $x \in \Sigma$, choose a neighbourhood $U_x \subset X$ which is an open ball around the origin in local coordinates as in Assumptions 7.5, and set $U = \bigcup_x U_x$. We know from our discussion of the local model that U is invariant under parallel transport along large circles. The claim is that for sufficiently large R, the grading of

$$\tilde{\psi}_R = (\tilde{h}'''_{\gamma_\infty})^2$$

(with respect to ω_X''', the given complex structure and complex volume form η_{X_R}) is negative on $X_R \setminus U$. This is proved by contradiction: suppose that we have a sequence $R_k \to \infty$, and points $x_k \in X_{R_k} \setminus U$ such that the grading of $\tilde{\psi}_R$ is not negative on some Lagrangian subspace in the tangent space at x_k. Consider the limit point of a subsequence, $x_\infty \in X_\infty \setminus U$. If x_∞ is a smooth point of X_∞, the grading of $\tilde{\psi}_R$ at x_{z_k} converges to -2 for the same reason as in the case of Lefschetz pencils, see Lemma 6.11. If x_∞ is a point where several components of X_∞ meet, Lemma 7.16 says that the grading of $\tilde{\psi}_R$ at x_{z_k} will be bounded above by $-1/2$ for large k (when applying this local result, the reader should keep in mind the change of coordinates on the base, $\zeta = 1/z$ and $dz = d\zeta/\zeta^2$).

Fix some R such that the claim made above applies. Take a neighbourhood $W \subset U \cap X_R$ of Σ. After making W smaller, we may again assume that in the local coordinates of Assumption 7.5, this is the intersection of the fibre $Y_{1/R}$ with an open ball around the origin. Lemma 7.6 tells us that $h'''_{\gamma_\infty}(W) = W$, and that h'''_{γ_∞} is negative on $U \setminus W$. $\square$

8. FUKAYA CATEGORIES

There are Fukaya categories for affine and projective Calabi-Yaus, as well as a relative version which interpolates between the two. All of them are c-unital A_∞-categories, but with different coefficient rings:

	notation for category	notation for objects	coefficient ring
affine	$\mathcal{F}(M)$	$L^\flat = (L, \tilde{\alpha}_L, \$_L)$	$\mathbb{C}$
relative	$\mathcal{F}(X, X_\infty)$	$L^\natural = (L, \tilde{\alpha}_L, \$_L, J_L)$	$\Lambda_\mathbb{N}$
projective	$\mathcal{F}(X)$	$L^\sharp = (L, \tilde{\alpha}_L, \$_L, \lambda_L, J_L)$	$\Lambda_\mathbb{Q}$

While our definition of affine Fukaya categories is fairly general, in the projective and relative setups only Calabi-Yau surfaces ($K3$ and tori) will be considered. This restriction allows us to avoid using virtual fundamental chains, and it also simplifies the relation between the different categories: up to quasi-equivalence, $\mathcal{F}(X, X_\infty)$ reduces to $\mathcal{F}(M)$ if the deformation parameter q is set to zero, while tensoring with $\Lambda_\mathbb{Q}$ gives a full subcategory of $\mathcal{F}(X)$ (for the definition of $\mathcal{F}(X)$ in higher dimensions, see [18]; and for its presumed relation with $\mathcal{F}(M)$, [59]). We emphasize that all properties of Fukaya categories which we use are quite simple ones, and largely independent of the finer details of the definition.

(8a) The following is a summary of [61, Chapter 2], with minor changes in notation and degree of generality. Let $M = X \setminus X_\infty$ be an affine Calabi-Yau. Objects of $\mathcal{F}(M)$ are *exact Lagrangian branes*

$$L^\flat = (L, \tilde{\alpha}_L, \$_L), \tag{8.1}$$

where $L \subset M$ is an oriented exact Lagrangian submanifold, $\tilde{\alpha}_L$ a grading which is compatible with the orientation (see Remark 6.6), and $\$_L$ a *Spin* structure. To define the morphism spaces, choose for each pair of objects a *Floer datum*

$$(H, J) = (H_{L_0^\flat, L_1^\flat}, J_{L_0^\flat, L_1^\flat}).$$

This consists of a function $H \in C^\infty_c([0;1] \times M, \mathbb{R})$, as well as a family $J = J_t$ of ω_X-compatible almost complex structures parametrized by $t \in [0;1]$, which for all t agree with the given complex structure in a neighbourhood of X_∞. The condition on H is that the associated Hamiltonian isotopy (ϕ^t_H) should satisfy $\phi^1_H(L_0) \pitchfork L_1$. We then define $\mathfrak{C}(L_0^\flat, L_1^\flat)$ to be the set of trajectories $x : [0;1] \rightarrow M$, $x(t) = \phi^t_H(x(0))$, with boundary conditions $x(k) \in L_k$. This is obviously bijective to $\phi^1_H(L_0) \cap L_1$, hence finite. To each such x one can associate a Maslov index $I(x)$ and orientation set $or(x)$. For $I(x)$, one observes that the grading of L_0 induces one of $\phi^1_H(L_0)$, and then uses the index of an intersection point as defined in (6.17). $or(x)$ is a set with two elements, the two possible "coherent orientations" of x; we refer to [61, 18] for its definition, which uses the *Spin* structures on L_0, L_1. The Floer cochain space is the graded

$\mathbb{C}$-vector space

$$CF^j_M(L^\flat_0, L^\flat_1) = \bigoplus_{\substack{x \in \mathfrak{C}(L^\flat_0, L^\flat_1) \\ I(x)=j}} \mathbb{C}_x, \tag{8.2}$$

where $\mathbb{C}_x$ is defined as the quotient of the two-dimensional space $\mathbb{C}[or(x)]$ by the relation that the two elements of $or(x)$ must sum to zero (it is of course isomorphic to $\mathbb{C}$, but not canonically). In the Fukaya category, CF^*_M is the space of *hom*s from $L^\flat_0$ to $L^\flat_1$.

Given $x_0, x_1 \in \mathfrak{C}(L^\flat_0, L^\flat_1)$, we now consider the moduli spaces $\mathfrak{M}^1_M(x_0, x_1)$ of solutions $u : \mathbb{R} \times [0;1] \to M$ to Floer's "gradient flow" equation with limits x_0, x_1, divided by the $\mathbb{R}$-action by translation. As part of the definition of Floer datum, we require that these spaces should all be regular. Assume from now on that $I(x_1) = I(x_0) - 1$. By an index formula $\mathfrak{M}^1_M(x_0, x_1)$ is zero-dimensional, and a compactness theorem says that it is a finite set. A few words about this: since the action functional is exact (6.6) and there is no bubbling, we only have to worry about sequences of solutions which become increasingly close to X_∞. To see that this does not happen, one uses a maximum principle argument based on the function $-\log ||\sigma_{X,\infty}||^2$, which goes to $+\infty$ at X_∞. It is plurisubharmonic since

$$\partial\bar{\partial} \log ||\sigma_{X,\infty}||^2 = -F_{\nabla_X} = 2\pi i \omega_M. \tag{8.3}$$

Each $u \in \mathfrak{M}^1_M(x_0, x_1)$ comes with a preferred identification $or(x_1) \cong or(x_0)$, which in turn defines a map $\mathbb{C}_{x_1} \to \mathbb{C}_{x_0}$. The sum of these over all u is the Floer differential $\mu^1_M : CF^j_M(L^\flat_0, L^\flat_1) \to CF^{j+1}_M(L^\flat_0, L^\flat_1)$, which is the first composition map in the Fukaya category. Its cohomology is the Floer cohomology $HF^*_M(L^\flat_0, L^\flat_1)$, which is therefore the space of morphisms in the cohomological category $H\mathcal{F}(M)$.

To define the rest of the A_∞-structure, one proceeds as follows. Take some $d \geq 2$. A $(d+1)$-*pointed disc with Lagrangian labels*

$$(S, \zeta_0, \dots, \zeta_d, L^\flat_0, \dots, L^\flat_d) \tag{8.4}$$

is a Riemann surface S isomorphic to the closed disc with $d+1$ boundary points removed. Moreover, these points at infinity should have a preferred numbering $\zeta_0, \dots, \zeta_d$, compatible with their natural cyclic order. Denote the connected components of ∂S by $I_0, \dots, I_d$, again in cyclic order and starting with the component I_0 which lies between ζ_0 and ζ_1. Finally, we want to have exact Lagrangian branes $L^\flat_0, \dots, L^\flat_d$ attached to the boundary components. A set of *strip-like ends* for S consists of proper holomorphic embeddings

$$\begin{aligned} \epsilon_{S,0} : \mathbb{R}^{\leq 0} \times [0;1] &\longrightarrow S, & \lim_{s \to -\infty} \epsilon_{S,0}(s, \cdot) &= \zeta_0, \\ \epsilon_{S,1}, \dots, \epsilon_{S,d} : \mathbb{R}^{\geq 0} \times [0;1] &\longrightarrow S, & \lim_{s \to +\infty} \epsilon_{S,k}(s, \cdot) &= \zeta_k \end{aligned}$$

taking $\mathbb{R}^{\leq 0} \times \{0;1\}$, $\mathbb{R}^{\geq 0} \times \{0;1\}$ to ∂S and which have disjoint images. A *perturbation datum* on S is a pair (K_S, J_S) consisting of a family $\{J_{S,z}\}_{z \in S}$ of compatible almost complex structures on X, which agree with the given complex structure in a neighbourhood of X_∞, and a one-form $K_S \in \Omega^1(S, C_c^\infty(M,\mathbb{R}))$ with values in smooth functions, subject to the condition that $K_S(\xi)|L_k = 0$ for $\xi \in T(I_k) \subset T(\partial S)$. The behaviour of (K_S, J_S) over the strip-like ends is fixed by the previously chosen Floer data:

$$J_{S,\epsilon_{S,k}(s,t)} = \begin{cases} J_{L_0^\flat,L_d^\flat,t} & k=0,\\ J_{L_{k-1}^\flat,L_k^\flat,t} & k>0 \end{cases} \quad \text{and} \quad \epsilon_{S,k}^* K_S = \begin{cases} H_{L_0^\flat,L_d^\flat,t}\,dt & k=0,\\ H_{L_{k-1}^\flat,L_k^\flat,t}\,dt & k>0. \end{cases}$$

Suppose that we have made a global choice of strip-like ends and perturbation data. This means that for every $(d+1)$-pointed disc with Lagrangian labels we have chosen a set of strip-like ends (K_S, J_S) (the dependence on the other data in (8.4) is suppressed from the notation for the sake of brevity). This needs to satisfy two additional kinds of conditions: first of all, it needs to vary smoothly if we change the complex structure on S in a smooth way, which means that it is defined on the universal family of such discs. Secondly, it needs to be well-behaved with respect to the compactification of that family, which includes degenerations to discs with nodes. This in fact establishes recursive relations between the choices of perturbation data for different d, see [61] for details.

Supposing that all these conditions have been met, take $x_0 \in \mathfrak{C}(L_0^\flat, L_d^\flat)$ as well as $x_k \in \mathfrak{C}(L_{k-1}^\flat, L_k^\flat)$ for $k = 1, \dots, d$. Define $\mathfrak{M}_M^d(x_0, \dots, x_d)$ to be the space of equivalence classes of pairs consisting of a $(d+1)$-marked disc with labels $L_0^\flat, \dots, L_d^\flat$, and a solution of the following generalization of Floer's equation for a map $u : S \to M$:

$$\left\{ \begin{array}{l} u(I_k) \subset L_k, \\ (du(z) - Y_{S,z,u(z)}) + J_{S,z,u(z)} \circ (du - Y_{S,z,u(z)}) \circ i = 0, \\ \lim_{s \to \pm\infty} u(\epsilon_{S,k}(s, \cdot)) = x_k. \end{array} \right. \tag{8.5}$$

Here $Y_S \in \Omega^1(S, C_c^\infty(TM))$ is the Hamiltonian-vector-field valued one-form determined by K_S. The equivalence relation on pairs is isomorphism of the underlying Riemann surfaces, compatible with all the additional structure, and composition of maps u with that isomorphism. The resulting quotient space is (locally) the zero-set of a Fredholm section of a suitably defined Banach vector bundle. Generically, this will be transverse, and then $\mathfrak{M}_M^d(x_0, \dots, x_d)$ is smooth of dimension $I(x_0) - I(x_1) - \cdots - I(x_d) + d - 2$. Moreover, the zero-dimensional spaces are finite. It is maybe worth while stating the basic energy equality which underlies the compactness argument. Choose primitives K_{L_k} for $\theta|L_k$. Define the action of the limits x_k by a generalization of (6.5),

$$A(x_k) = -\int x_k^*\theta + \\ + \begin{cases} \int H_{L_0^\flat,L_d^\flat}(x_0(t))\,dt + K_{L_d}(x_0(1)) - K_{L_0}(x_0(0)) & k=0,\\ \int H_{L_{k-1}^\flat,L_k^\flat}(x_k(t))\,dt + K_{L_k}(x_k(1)) - K_{L_{k-1}}(x_k(0)) & k=1,\dots,d. \end{cases}$$

Then for $u \in \mathfrak{M}^d_M(x_0, \dots, x_d)$,

$$\begin{aligned} E(u) &\overset{\text{def}}{=} \tfrac{1}{2}\int_S |du - Y|^2 \\ &= A(x_0) - \sum_{k=1}^{d} A(x_d) - \int_S u^*(-dK_S + \tfrac{1}{2}\{K_S, K_S\}). \end{aligned} \tag{8.6}$$

The integral on the right hand side can be bounded by a constant which is independent of u, because the curvature term $-dK_S + \frac{1}{2}\{K_S, K_S\}$ vanishes over the strip-like ends of S. Of course (8.3) also plays a role in the proof of compactness, as was the case for the Floer differential. To continue with the main story: any element of a zero-dimensional space $\mathfrak{M}^d_M(x_0, \dots, x_d)$ defines a canonical bijection

$$or(x_0) \cong or(x_1) \times_{\mathbb{Z}/2} \cdots \times_{\mathbb{Z}/2} or(x_d). \tag{8.7}$$

The order d composition map μ^d_M in $\mathcal{F}(M)$ is the sum of the corresponding maps $\mathbb{C}_{x_d} \otimes \dots \otimes \mathbb{C}_{x_1} \to \mathbb{C}_{x_0}$ (the order is reversed according to our notational conventions for A_∞-categories). There are additional signs which involve the Maslov indices $I(x_k)$, but since they depend on the conventions about determinant lines used in setting up the orientation spaces, we prefer not to spell them out here. $\mathcal{F}(M)$ is independent of all the choices (Floer data, strip-like ends, perturbation data) made in its construction up to quasi-equivalence, and indeed quasi-isomorphism acting trivially on the objects.

Remark 8.1. *Suppose that no irreducible component of X_∞ is rational. Then one can use a wider class of almost complex structures, namely all those ω_X-compatible ones such that each irreducible component $C \subset X_\infty$ is an almost complex submanifold. Instead of using* (8.3) *to prevent solutions of Floer's equation and its generalization from escaping into X_∞, one now argues as follows: if the solution stays inside M, its intersection number with C is zero. The limit of a sequence of solutions has the same property, and nonnegativity of intersection multiplicities shows that the limit is disjoint from X_∞ (more precisely, one thinks of a solution of* (8.5) *as a map $u : S \to S \times X$ which is pseudo-holomorphic for a suitable almost complex structure, and considers its intersection with $S \times X_\infty$). Similarly, in the inhomogeneous terms one can use Hamiltonian functions on X whose values and first derivatives vanish on X_∞.*

(8b) Let M be an affine Calabi-Yau, equipped with a homomorphism

$$\rho : \pi_1(M) \longrightarrow \Gamma$$

into some finitely generated abelian group Γ. Consider the associated abelian covering $M_\Gamma \to M$. Denote by $\mathcal{F}(M)_\rho$ the full subcategory of $\mathcal{F}(M)$ consisting of those branes $L^\flat$ such that the composition $\pi_1(L) \to \pi_1(M) \to \Gamma$ is trivial. For each of these choose a lift $L_\Gamma \subset M_\Gamma$. Once one has done that, the *hom*-spaces split as

$$CF^*_M(L_0^\flat, L_1^\flat) = \bigoplus_{\gamma \in \Gamma} CF^*_M(L_0^\flat, L_1^\flat)_\gamma,$$

where the γ-summand consists of those $x \in \mathfrak{C}(L_0^\flat, L_1^\flat)$ whose unique lift $x_\Gamma : [0;1] \rightarrow M_\Gamma$ with $x_\Gamma(0) \in L_{0,\Gamma}$ satisfies $\gamma^{-1}(x_\Gamma(1)) \in L_{1,\Gamma}$. Because each pseudo-holomorphic polygon can be lifted to M_Γ, these splittings are compatible with the compositions, in the sense that

$$CF^*_M(L_{d-1}^\flat, L_d^\flat)_{\gamma_d} \otimes \dots \otimes CF^*_M(L_0^\flat, L_1^\flat)_{\gamma_1} \xrightarrow{\mu^d_M} CF^*_M(L_0^\flat, L_d^\flat)_{\gamma_1 \dots \gamma_d}.$$

This constitutes a Γ-*coaction* on our A_∞-category. For general nonsense reasons, it defines an *action* of the character group $\Gamma^* = Hom(\Gamma, \mathbb{C}^*)$: elements $\chi \in \Gamma^*$ act on each $CF^*_M(L_0^\flat, L_1^\flat)_\gamma$ by scalar multiplication with $\chi(\gamma) \in \mathbb{C}^*$.

What we actually need is a parallel but slightly different geometric setup. Let $M = X \setminus X_\infty$ be an affine Calabi-Yau, and Γ a finite abelian group acting on X by Kähler isometries. This should come with a lift of the action to o_X, which preserves A_X and is compatible with the induced action on $\mathcal{K}_X$ via the isomorphism β_X (6.18). Finally, we require that the action preserves $\sigma_{X,\infty}$ up to multiplication with $|m_X|$-th roots of unity (for $m_X = 0$ these may be arbitrary constants in S^1), and that it is free on M. As a consequence of these assumptions, the one-form θ_M and the holomorphic volume form η_M are Γ-invariant. Let $\bar{M} = M/\Gamma$ be the quotient, with its inherited symplectic form $\omega_{\bar{M}}$, one-form $\theta_{\bar{M}}$ and holomorphic volume form $\eta_{\bar{M}}$. Note that $-\log ||\sigma_{X,\infty}||^2$ also descends to a proper subharmonic function on $\bar{M}$. One can therefore define the Fukaya category $\mathcal{F}(\bar{M})$ in the same way as for M itself, using almost complex structures which agree with the given complex structure outside a compact subset (this is in spite of the fact that the obvious compactification $\bar{X} = X/\Gamma$ may be singular). There is a tautological homomorphism $\rho : \pi_1(\bar{M}) \rightarrow \Gamma$, whose associated covering is $\bar{M}_\Gamma = M$. The observation made above applies to this case as well, so one has a Γ^*-action on $\mathcal{F}(\bar{M})_\rho$. Take the semidirect product A_∞-category, as defined in Section 4b.

LEMMA 8.2. *$\mathcal{F}(\bar{M})_\rho \rtimes \Gamma^*$ is quasi-isomorphic to the full subcategory of $\mathcal{F}(M)$ consisting of those branes in M which satisfy $\gamma(L) \cap L = \emptyset$ for all nontrivial $\gamma \in \Gamma$.* □

The proof is straightforward, and we leave it to the reader. Note that the Γ^*-action is not quite unique, since it depends on choosing lifts for the branes; but different choices lead to group actions related by (elementary) inner automorphisms as in (4.1), which explains why the semidirect product is the same for all choices.

(8c) Let X be a projective Calabi-Yau surface, and $L \subset X$ a Lagrangian submanifold which admits a grading. An ω_X-compatible almost complex structure

J is called *regular with respect to* L if there are no non-constant J-holomorphic spheres, and no non-constant J-holomorphic discs with boundary on L.

LEMMA 8.3. *Almost complex structures which are regular with respect to a given* L *are dense within the set of all* ω_X*-compatible almost complex structures.*

PROOF. Since $c_1(X) = 0$, the virtual dimension of the space of unparametrized J-holomorphic spheres at a point v is $4 + 2\langle c_1(X), [v]\rangle - 6 = -2$. Standard transversality results tell us that for generic J, there are no J-holomorphic spheres that are somewhere injective; and since any non-constant sphere is a multiple cover of a somewhere injective one, there are no non-constant spheres at all. The argument for discs goes along the same lines: because of the grading, the Maslov number $\mu(w)$ of any disc in (X, L) vanishes, and the virtual dimension is

$$2 + \mu(w) - 3 = -1. \tag{8.8}$$

Again, this shows that for generic J, there are no somewhere injective discs. The decomposition theorem of Kwon-Oh [38] and Lazzarini [39] says that if there is a non-constant J-holomorphic disc, there is also a somewhere injective one, which implies the desired result. □

Objects of $\mathcal{F}(X)$ are *rational Lagrangian branes*

$$L^\sharp = (L, \tilde{\alpha}_L, \$_L, \lambda_L, J_L) \tag{8.9}$$

where L is an oriented Lagrangian submanifold of X with a grading $\tilde{\alpha}_L$ and *Spin* structure $\$_L$; λ_L is a covariantly constant multisection of the circle bundle of $o_X|L$ of some degree $d_L \geq 1$; and J_L is an ω_X-compatible almost complex structure which is regular with respect to L. For any pair of objects, we take a Floer datum

$$(H, J) = (H_{L_0^\sharp, L_1^\sharp}, J_{L_0^\sharp, L_1^\sharp})$$

as before, except that the conditions on the behaviour of H and J near X_∞ are omitted, and the following new requirements imposed instead: J_t must be equal to J_{L_0}, J_{L_1} for $t = 0, 1$; and there should be no non-constant J_t-holomorphic spheres for any t (this is still true generically, because the virtual dimension of the parametrized moduli space is -1). Define the set $\mathfrak{C}(L_0^\sharp, L_1^\sharp)$ as before; and for any $x : [0; 1] \to X$ in that set, write $\bar{A}(x) = \bar{A}(x(1))$ for the mod $\mathbb{Q}$ action (6.2) of $x(1)$ as an intersection point of $\phi_H^1(L_0) \cap L_1$ (where we have used (6.1) to get a multisection of $o_X|\phi_H^1(L_0)$ from λ_{L_0}). To spell this out, we have

$$\exp(2\pi i \bar{A}(x)) = \frac{\lambda_{L_1}(x(1))}{\kappa(\lambda_{L_0}(x(0)))} \exp\Big(2\pi i \int_0^1 H_t(x(t))\, dt\Big),$$

where κ is parallel transport on o_X along x. For any number $r \in \mathbb{R}/\mathbb{Q}$, we define a one-dimensional $\Lambda_{\mathbb{Q}}$-vector space $q^r\Lambda_{\mathbb{Q}}$ whose elements are formal series $f(q) = \sum_m a_m q^m$, with m running over all real numbers in $r + \frac{1}{d}\mathbb{Z}$ for some natural number d which depends on f, and $a_m \in \mathbb{C}$ vanishing for sufficiently negative m. Write $\Lambda_{\mathbb{Q},x}$ for the quotient of $q^{\bar{A}(x)}\Lambda_{\mathbb{Q}}[or(x)]$ by the relation that the two elements of $or(x)$ sum to zero. Then the Floer cochain space is

$$CF^j_X(L_0^\sharp, L_1^\sharp) = \bigoplus_{\substack{x \in \mathfrak{C}(L_0^\sharp, L_1^\sharp) \\ I(x)=j}} \Lambda_{\mathbb{Q},x}.$$

Denote by $\mathfrak{M}^1_X(x_0,x_1)$ the space of solutions of Floer's equation in X with limits x_0, x_1 (this means we drop the previous assumption that the solutions stay in M). As before, we require that this is regular, and in particular zero-dimensional in the case $I(x_1) = I(x_0) - 1$. Take a solution u in such a zero-dimensional moduli space, and let $E(u) = \int |\partial u/\partial s|^2 \, ds\, dt > 0$ be its energy. A version of (6.3) shows that $E(u) \in \bar{A}(x_0) - \bar{A}(x_1) + l.c.m.(d_{L_0}, d_{L_1})^{-!}\mathbb{Z}$, so that we get a well-defined $\Lambda_{\mathbb{Q}}$-module map

$$\pm q^{E(u)} : \Lambda_{\mathbb{Q},x_1} \longrightarrow \Lambda_{\mathbb{Q},x_0}.$$

The sign of this is determined by the induced bijection of orientation sets, as before, and summing over all u yields the Floer differential μ^1_X in $\mathcal{F}(X)$. The sum is no longer finite, but it gives a well-defined $\Lambda_{\mathbb{Q}}$-module map by Gromov compactness; the basic point being that there is no bubbling off of holomorphic discs or spheres, by assumption on the Floer datum. Denote the resulting Floer cohomology by $HF^*_X(L_0^\sharp, L_1^\sharp)$.

The definition of the higher order compositions is largely similar. One may use only perturbation data (K_S, J_S) where $J_S = J_{L_k}$ over the boundary component $I_k \subset \partial S$. It is no longer possible to avoid $J_{S,z}$-holomorphic spheres for some values of $z \in int(S)$, but the argument from [30] shows that for a generic choice of J_S, these spheres will never occur as bubbles in the Gromov compactification of the spaces $\mathfrak{M}^d_X(x_0, \dots, x_d)$ as long as these are of dimension ≤ 1, which is all that we will ever use. Equation (8.6) still holds if we replace all the actions by their mod $\mathbb{Q}$ counterparts. We define the contribution of a point (S,u) in a zero-dimensional moduli space $\mathfrak{M}^d_X(x_0,\dots,x_d)$ to μ^d_X to be

$$\pm q^{E(u) + \int_S u^*(-dK_S + \frac{1}{2}\{K_S,K_S\})} : \Lambda_{\mathbb{Q},x_d} \otimes \cdots \otimes \Lambda_{\mathbb{Q},x_1} \longrightarrow \Lambda_{\mathbb{Q},x_0}, \tag{8.10}$$

with the sign again depending on (8.7). There is a uniform lower bound for the $\int_S$ term. This, together with Gromov compactness, ensures that the sum is a well-defined $\Lambda_{\mathbb{Q}}$-multilinear map.

Remark 8.4. *The choice of λ_L serves only to fix the action: the Floer cohomology groups for different choices are isomorphic in an obvious way, by multiplication with q^m for some $m \in \mathbb{Q}$. In particular, changing λ_L does not affect the quasi-isomorphism class of the resulting object in $\mathcal{F}(X)$.*

(8d) Take X as before, but now equipped with a section $\sigma_{X,\infty}$ of o_X whose zero set X_∞ is a reduced divisor with normal crossings, so that $M = X \setminus X_\infty$ is an affine Calabi-Yau surface. Following Remark 8.1, we also assume that X_∞ has no rational components. The objects of the relative Fukaya category $\mathcal{F}(X, X_\infty)$ are a mixture of the two previously introduced classes, (8.1) and (8.9). One considers *relative Lagrangian branes*

$$L^\natural = (L, \tilde{\alpha}_L, \$_L, J_L) \tag{8.11}$$

where $(L, \tilde{\alpha}_L, \$_L)$ is an exact Lagrangian brane in M; and J_L is an ω_X-compatible almost complex structure on X such that each component of X_∞ is an almost complex submanifold, and which is regular with respect L. The restriction on $J_L|X_\infty$ does not interfere with the regularity theory for somewhere injective pseudo-holomorphic spheres $v : S^2 \to X$, since $v^{-1}(X_\infty)$ is necessarily a finite set, while the set of somewhere injective points is open; and the same applies to pseudo-holomorphic discs. For any two objects, choose a Floer datum

$$(H, J) = (H_{L_0^\natural, L_1^\natural}, J_{L_0^\natural, L_1^\natural})$$

such that $H \in C^\infty([0;1] \times X, \mathbb{R})$ vanishes along $[0;1] \times X_\infty$, along with its first derivative; and $J_t|X_\infty$ makes the components of X_∞ almost complex. In addition, we assume that $J_k = J_{L_k}$ for $k = 0, 1$, and that there are no non-constant J_t-holomorphic spheres for any t.

We have the usual set $\mathfrak{C}(L_0^\natural, L_1^\natural)$ of H-trajectories joining L_0 to L_1, and for x in that set we define $\Lambda_{\mathbb{N},x}$ in the same way as $\mathbb{C}_x$, just using $\Lambda_\mathbb{N}$ instead of $\mathbb{C}$ as the ground ring. The Floer cochain space in $\mathcal{F}(X, X_\infty)$ is

$$CF^j_{X,X_\infty}(L_0^\natural, L_1^\natural) = \bigoplus_{\substack{x \in \mathfrak{C}(L_0^\natural, L_1^\natural) \\ I(x) = j}} \Lambda_{\mathbb{N},x}.$$

For the Floer differential, each $u \in \mathfrak{M}^1_X(x_0, x_1)$ contributes with $\pm q^{u \cdot X_\infty} : \Lambda_{\mathbb{N},x_1} \to \Lambda_{\mathbb{N},x_0}$. This makes sense for a single u since the number $u \cdot X_\infty$ is nonnegative; and Gromov compactness, together with a version of (6.6), says that the possibly infinite sum over all u gives rise to a well-defined map μ^1_{X,X_∞}. Denote the resulting Floer cohomology by $HF^*_{X,X_\infty}(L_0^\natural, L_1^\natural)$. The definition of the higher order compositions works in the same way, and we will therefore omit the details.

Remark 8.5. *Let X, X' be two Calabi-Yau surfaces, with sections $\sigma_{X,\infty}$, $\sigma_{X',\infty}$ of their ample line bundles o_X, $o_{X'}$ which define reduced divisors with normal crossings, having no rational components. Suppose that there is a symplectic isomorphism*

$$\phi : (X, X_\infty) \longrightarrow (X', X'_\infty)$$

with the following additional properties: $Hom(o_X, \phi^* o_{X'})$ is the trivial flat line bundle, and over $M = X \setminus X_\infty$, its covariantly constant sections lie in the same homotopy class of nowhere zero sections as $\phi^*\sigma_{X',\infty}/\sigma_{X,\infty}$. Moreover, under the isomorphism (unique up to homotopy) $\mathcal{K}_X \cong \phi^*\mathcal{K}_{X'}$ induced by deforming the almost complex structure on X to the pullback of the one from X', the nowhere zero sections η_X and $\phi^*\eta_{X'}$ are homotopic. Then, by suitably choosing *the Floer and perturbation data, one can define an induced quasi-isomorphism of A_∞-categories over $\Lambda_{\mathbb{N}}$,*

$$\phi_* : \mathcal{F}(X, X_\infty) \to \mathcal{F}(X', X'_\infty).$$

In the case where $H_1(X) = 0$, all the additional conditions imposed on ϕ are automatically satisfied, for the same reasons as in Remark 6.3.

(8e) Finally, we need to establish the relation between the different kinds of Fukaya categories. Let X be a Calabi-Yau surface with a suitable section $\sigma_{X,\infty}$, so that $\mathcal{F}(M)$, $\mathcal{F}(X)$ and $\mathcal{F}(X, X_\infty)$ are all defined.

Proposition 8.6. *Consider $\mathbb{C}$ as a $\Lambda_{\mathbb{N}}$-module in the obvious way, with q acting trivially. Then the $\mathbb{C}$-linear A_∞-category obtained from $\mathcal{F}(X, X_\infty)$ by reduction of constants, $\mathcal{F}(X, X_\infty) \otimes_{\Lambda_{\mathbb{N}}} \mathbb{C}$, is quasi-equivalent to $\mathcal{F}(M)$.*

Proposition 8.7. *Consider $\Lambda_{\mathbb{Q}} \supset \Lambda_{\mathbb{N}}$ as a $\Lambda_{\mathbb{N}}$-module in the obvious way. Then the $\Lambda_{\mathbb{Q}}$-linear category obtained from $\mathcal{F}(X, X_\infty)$ by extension of constants, $\mathcal{F}(X, X_\infty) \otimes_{\Lambda_{\mathbb{N}}} \Lambda_{\mathbb{Q}}$, is quasi-isomorphic to a full A_∞-subcategory of $\mathcal{F}(X)$. This subcategory consists of all those $L^\sharp$ such that $L \cap X_\infty = \emptyset$, $o_X|L$ is the trivial flat bundle, and its covariantly constant sections lie in the same homotopy class as $\sigma_{X,\infty}|L$.*

Both statements are essentially obvious. For Proposition 8.6, one notices that setting $q = 0$ means that we count only those solutions of Floer's equation or its generalization which have zero intersection number with, hence are disjoint from, X_∞. Then, the only difference between $\mathcal{F}(M)$ and $\mathcal{F}(X, X_\infty) \otimes_{\Lambda_{\mathbb{N}}} \mathbb{C}$ is that the latter category has more objects, consisting of the same exact Lagrangian brane with different choices of J_L. However, all these objects are isomorphic, whence the quasi-equivalence. For Proposition 8.7 one has to choose a function K_L, $dK_L = \theta_M|L$, for each exact Lagrangian brane in M. The associated object in $\mathcal{F}(X)$ then carries the multisection λ_L from (6.4), and one identifies

$$CF^*_{(X,X_\infty)}(L_0^\natural, L_1^\natural) \otimes_{\Lambda_{\mathbb{N}}} \Lambda_{\mathbb{Q}} \xrightarrow{\cong} CF^*_X(L_0^\sharp, L_1^\sharp) \tag{8.12}$$

by using the map $q^{A(x)} : \Lambda_{\mathbb{N},x} \otimes_{\Lambda_{\mathbb{N}}} \Lambda_{\mathbb{Q}} \to \Lambda_{\mathbb{Q},x}$ on each one-dimensional subspace. The compositions in the two categories are based on the same moduli spaces, and the fact that the weights used to count solutions correspond under (8.12) is a consequence of (6.6) and a suitable generalization. Clearly, $\mathcal{F}(X, X_\infty) \otimes_{\Lambda_{\mathbb{N}}} \Lambda_{\mathbb{Q}}$ is not all of $\mathcal{F}(X)$, since only the objects which are exact Lagrangian submanifolds in M occur.

(8f) While the basic properties of affine Fukaya categories are familiar and quite simple, see for instance [61], the relative and projective versions reserve some small surprises. These are all covered by the powerful general theory from [18], but we prefer a more elementary approach. The relative situation is the more important one for the arguments later on, and we will consider it primarily.

Fix an affine Calabi-Yau surface $M = X \setminus X_\infty$, without rational components in X_∞. Take an exact Lagrangian brane $(L, \tilde{\alpha}_L, \$_L)$ in M, and a one-parameter family J_t $(0 \leq t \leq 1)$ of compatible almost complex structures, all of which make the components of X_∞ almost complex, and are regular with respect to L. These give rise to a family of objects of $\mathcal{F}(X, X_\infty)$, which we denote by $L_t^\natural$. By suitably adapting the construction of Piunikhin-Salamon-Schwarz [48], one defines a canonical isomorphism

$$H\Psi_{X,X_\infty} : H^*(L;\mathbb{C}) \otimes_{\mathbb{C}} \Lambda_{\mathbb{N}} \xrightarrow{\cong} HF^*_{X,X_\infty}(L_0^\natural, L_1^\natural). \tag{8.13}$$

In the special case where J_t is constant, one can use this to obtain a distinguished element $e = H\Psi_{X,X_\infty}(1) \in HF^0_{X,X_\infty}(L_0^\natural, L_0^\natural)$, which is in fact the cohomological unit for $L_0^\natural$ in $H\mathcal{F}(X, X_\infty)$. The more general isomorphisms (8.13), for different choices of almost complex structures, carry the ordinary cup product to the pair of pants product. This eventually shows that $L_0^\natural$, $L_1^\natural$ are quasi-isomorphic objects. A slight generalization, which in fact is already implicit in our use of Floer data, then proves the following

Lemma 8.8. *Let L_t be a family of Lagrangian submanifolds of M, and $L_t^\natural$ a family of relative brane structures on them, varying smoothly with t. Then all $L_t^\natural$ are quasi-isomorphic objects of $\mathcal{F}(X, X_\infty)$.* ☐

Note that because all the L_t are supposed to have relative brane structures, they are in particular exact in M, so that the isotopy is Hamiltonian.

(8g) We will now summarize the definition of $H\Psi_{X,X_\infty}$. Pick a cohomology class $c \in H^j(L;\mathbb{Z})$, and represent its Poincaré dual by a map $\gamma : C \to L$ from a closed oriented manifold C (recall that L is a surface). Take the Riemann surface $S = \mathbb{H}$, the closed upper half plane, and consider a generalized Floer equation for pairs (u, y) consisting of a map $u : S \to M$ and a point $y \in C$,

$$\begin{cases} u(\partial S) \subset L, \\ u(0) = \gamma(c), \\ (du - Y_{S,u(z),z}) + J_{S,z,u(z)} \circ (du - Y_{S,u(z),z} \circ i) = 0, \\ \lim_{s\to+\infty} u(e^{\pi(s+it)}) = x_0(1-t), \quad x_0 \in \mathfrak{C}(L_0^\natural, L_1^\natural). \end{cases} \tag{8.14}$$

The choice of almost complex structures J_S and inhomogeneous term Y_S has to obey a whole series of conditions, of which we mention only the most relevant one: along the boundary $\partial S = \mathbb{R}$, $J_{S,z} = J_{\psi(z)}$ for some function $\psi : \mathbb{R} \to [0;1]$ with $\psi(s) = 0$ for very negative s; $\psi(s) = 1$ for very positive s; and $\psi'(s) > 0$ for all s such that $\psi(s) \in (0;1)$.

Assume that the moduli spaces $\mathfrak{M}^{\Psi}_{X,X_\infty}(x_0;\gamma)$ of solutions of (8.14) are all regular. If $I(x_0) = j$, the relevant moduli space is zero-dimensional, and every (u, y) in it comes with a canonical choice of point in $or(x_0)$. This will define an element $\pm q^{u\cdot X_\infty} \in \Lambda_{\mathbb{N},x_0}$, and the sum of all these is denoted by

$$\Psi_{X,X_\infty}(\gamma) \in CF^j_{X,X_\infty}(L_0^\natural, L_1^\natural).$$

The proof that this is a Floer cocycle relies on a standard gluing argument: one looks at the (one-dimensional) moduli spaces $\mathfrak{M}^{\Psi}_{X,X_\infty}(x_0;\gamma)$ with $I(x_0) = j+1$, and proves that points in the boundary of their Gromov compactification are pairs consisting of an isolated solution which contributes to $\Psi_{X,X_\infty}(\gamma)$ and a Floer gradient trajectory. The main point is that due to the choice of J_S, there is no bubbling off of pseudo-holomorphic discs. By definition, the Floer cohomology class of $\Psi_{X,X_\infty}(\gamma)$ is $H\Psi_{X,X_\infty}(c \otimes 1)$.

(8h) The assumption made up to now, that all the J_t are regular, is not generic: while a single almost complex structure will typically admit no nonconstant pseudo-holomorphic discs with boundary on L, these can no longer be avoided in one-parameter families, as one can see from (8.8). In other words, the space of all almost complex structures is divided into chambers by codimension one walls consisting of non-regular ones. Because there is an infinite number of possible homotopy classes of discs, there is potentially an infinite number of walls, which might well be everywhere dense. We will look at the simplest instance of the behaviour of Floer cohomology under wall-crossing (a complete treatment is beyond our scope here, since it requires techniques that can deal with the lack of transversality for multiply-covered discs, see [18]).

ASSUMPTIONS 8.9. *Fix an exact Lagrangian brane $(L, \tilde{\alpha}_L, \$_L)$ in M, and a family (J_t) of compatible almost complex structures, each of which makes the components of X_∞ almost complex. Suppose that this family has the following properties:*

- *J_0, J_1 are regular with respect to L;*
- *for all t, there are no J_t-holomorphic spheres $v : S^2 \to X$ such that $v \cdot X_\infty = 1$;*
- *there is a $t_0 \in (0;1)$ such that for all $t \neq t_0$, no J_t-holomorphic discs $w : (D, \partial D) \to (X, L)$ with $w \cdot X_\infty = 1$ exist. For J_{t_0}, there is exactly one such disc, which we denote by w_0. This is parametrized-regular, and $[w_0|\partial D] \in H_1(L)$ is nonzero.*

The terminology in the last sentence may require some explanation. Denote by $D\bar{\partial}_{w,X} : \Gamma(w^*TX, w^*TL) \to \Omega^{0,1}(w^*TX)$ the standard linearized operator associated to a J_t-holomorphic disc $w : (D, \partial D) \to (X, L)$. Here $\Gamma(E, F)$ means sections of the vector bundle $E \to D$ with boundary values in the totally real subbundle $F \subset E|\partial D$. The parametrized version,

$$D\bar{\partial}^{par}_{w,X} : \mathbb{R} \times \Gamma(w^*TX, w^*TL) \longrightarrow \Omega^{0,1}(w^*TX),$$

adds a first component $1 \mapsto \frac{1}{2}(\partial J_t/\partial t) \circ dw \circ i$. If this is onto in suitable Sobolev completions, we say that w is parametrized-regular.

Consider the equation (8.14) as before, choosing ψ in such a way that $\psi(0) = t_0$. For solutions $(u, y) \in \mathfrak{M}^{\Psi}_{X,X_\infty}(x_0; \gamma)$ with $I(x_0) = j$, which occur in zero-dimensional moduli spaces, one can achieve by a suitable generic choice of (K_S, J_S) that $u(z)$ does not meet the boundary of $J_{\psi(z)}$-holomorphic discs in (X, L), for any $z \in \mathbb{R}$. Hence one can define Floer cochains $\Psi_{X,X_\infty}(\gamma)$ as before. Expand these and the Floer differential on $CF^*_{X,X_\infty}(L^\natural_0, L^\natural_1)$ into powers of q,

$$\begin{aligned}\Psi_{X,X_\infty}(\gamma) &= \Psi_0(\gamma) + \Psi_1(\gamma)q + \Psi_2(\gamma)q^2 + \dots,\\ \mu^1_{X,X_\infty} &= \mu_0 + \mu_1 q + \mu_2 q^2 + \dots\end{aligned}$$

Claim 8.10. $\mu_0\Psi_0(\gamma) = 0$*; indeed,* Ψ_0 *is an isomorphism between* $H^*(L; \mathbb{C}) \otimes \Lambda_{\mathbb{N}}$ *and the cohomology of* $CF^*_{X,X_\infty}(L^\natural_0, L^\natural_1)$ *with only* μ_0 *as differential.*

This is simply the statement of the Piunikhin-Salamon-Schwarz isomorphism for L as an exact Lagrangian submanifold in the affine Calabi-Yau M: by Proposition 8.6, $\mu_0 = \mu^1_M$ is the differential in $\mathcal{F}(M)$, and similarly Ψ_0 counts solutions of (8.14) which stay inside M.

Claim 8.11. *Suppose that* $(C, \gamma) = (L, \mathrm{id})$ *is the whole surface. Then*

$$\mu_1\Psi_0(\gamma) + \mu_0\Psi_1(\gamma) = \pm\Psi_0(w_0|\partial D).$$

The $\Psi_0(w_0|\partial D)$ term comes from a new stratum in the compactification of the one-dimensional moduli spaces, which consists of solutions of (8.14) with the disc bubble w_0 attached to it at the point $z = 0$. Under the regularity condition on w_0 that we have imposed, these are smooth boundary points of the compactification.

A straightforward computation starting from the two claims above shows that

$$HF^0_{X,X_\infty}(L^\natural_0, L^\natural_1) = 0.$$

In particular $L^\natural_0$, $L^\natural_1$ are not quasi-isomorphic as objects of $\mathcal{F}(X, X_\infty)$. Moreover, if we cut off the q-series after the first order, then by the same kind of computation as before the natural map

$$H^0(CF^*_{X,X_\infty}(L^\natural_0, L^\natural_1) \otimes_{\Lambda_{\mathbb{N}}} \mathbb{C}[q]/q^2) \otimes_{\mathbb{C}[q]/q^2} \mathbb{C} \longrightarrow H^0(CF^*_M(L^\flat_0, L^\flat_1)) \tag{8.15}$$

vanishes. More precisely, both sides are isomorphic to $\mathbb{C}$, but the left hand side is represented by the cocycle $q\Psi_0([L])$, which maps to zero. As a consequence,

Corollary 8.12. *Suppose that there is some* $(L, \tilde{\alpha}_L, \$_L)$ *such that Assumptions 8.9 hold. Then* $\mathcal{F}(X, X_\infty)$*, seen as a one-parameter deformation of* A_∞*-categories, is nontrivial at first order in* q*.* □

(8i) Our assumption that $[w_0|\partial D] \in H_1(L)$ is nontrivial obviously excludes Lagrangian spheres, and indeed wall-crossing in trivial in that case:

LEMMA 8.13. *Up to quasi-isomorphism, a graded Lagrangian sphere* $\tilde{L}$ *in* M *gives rise to a unique object of* $\mathcal{F}(X, X_\infty)$.

PROOF. Take two objects $L_0^\natural, L_1^\natural$ obtained by various choices of relative brane structures on L (keeping the same grading). Clearly, these two objects are quasi-isomorphic in $\mathcal{F}(X, X_\infty) \otimes_{\Lambda_\mathbb{N}} \mathbb{C} \cong \mathcal{F}(M)$. Because there is no odd-dimensional cohomology, the quasi-isomorphism between them survives to an element of $HF^0_{X,X_\infty}(L_0^\natural, L_1^\natural)$, and an obvious spectral sequence argument shows that this is a quasi-isomorphism in $\mathcal{F}(X, X_\infty)$. □

REMARK 8.14. *Essentially all of our observations concerning relative Fukaya categories carry over to the projective case. First of all, we have an analogue of Lemma 8.8, saying that* the isomorphism class of an object of $\mathcal{F}(X)$ remains constant under smooth deformation. *Secondly,* crossing a wall on which a single lowest area holomorphic disc w_0 sits does change the isomorphism class of an object*, assuming as before that* w_0 *is parameterized-regular and* $[w_0|\partial D]$ *is nontrivial. Finally, for graded Lagrangian spheres in* X *we have the same statement as in Lemma 8.13. The proofs are quite similar, except that the filtration by intersection numbers with* X_∞ *has to be replaced by a filtration by values of the action functional. This follows methods that were pioneered by Oh* [47] *and vastly developed in* [18]. *In our case, the rationality assumption simplifies the technical details somewhat, since it makes the filtration discrete.*

(8j) As shown by Corollary 8.12, one can use wall-crossing phenomena in the relative Fukaya category to prove its nontriviality in the sense of deformation theory; provided of course that one can come up with a Lagrangian submanifold for which wall-crossing actually occurs. To make that easier, we now suppose that the Calabi-Yau surface under discussion is the fibre of a quasi-Lefschetz pencil, so that the desired Lagrangian submanifold can be first constructed in the degenerate fibre at ∞.

For this discussion, we change notation as follows: X is a Fano threefold with $o_X = \mathcal{K}_X^{-1}$, carrying a pair of sections $\sigma_{X,0}, \sigma_{X,\infty}$ which generate a quasi-Lefschetz pencil of Calabi-Yau surfaces $\{X_z\}$. To be able to define the relative Fukaya categories, we assume that $X_{0,\infty}$ has no rational components.

Fix some irreducible component C of X_∞, and denote by C' the union of all the other components. There is an obvious short exact sequence

$$0 \longrightarrow TC \longrightarrow TX|C \xrightarrow{D(\sigma_{X,\infty}/\sigma_{X,0})} \mathcal{O}_C(C \cap X_0 - C \cap C') \longrightarrow 0. \tag{8.16}$$

From the resulting isomorphism $\mathcal{K}_C \cong \mathcal{K}_X|C \otimes \mathcal{K}_X^{-1}(-C')|C \cong \mathcal{O}_C(-C' \cap C)$ one obtains a meromorphic two-form η_C on C with simple poles along $C \cap C'$. Similarly, from $A_X|C$ and $\sigma_{X,0}|C$ one obtains a one-form θ_C on $C \setminus X_0$ with $d\theta_C = \omega_C = \omega_X|C$.

Assumptions 8.15. *Suppose that we have the following data L_0, w_0:*

- *$L_0 \subset C$ is a Lagrangian submanifold, disjoint from $C' \cup X_0$, exact with respect to θ_C, and graded with respect to η_C. This implies that the Maslov number and area of any holomorphic disc $w : (D, \partial D) \to (C, L_0)$ can be computed as intersection numbers:*

$$\begin{aligned} \mu(w) &= 2(w \cdot C'), \\ \textstyle\int w^* \omega_C &= w \cdot X_0. \end{aligned} \tag{8.17}$$

- *$w_0 : (D, \partial D) \to (C, L_0)$ is a specific holomorphic disc with $w_0 \cdot X_0 = w_0 \cdot C' = 1$. The point $w_0(0)$ should lie in $C' \cap X_0$, and $[w_0|\partial D] \in H_1(L_0)$ should be nontrivial. Take the linearized operator of w_0, and restrict its domain to those sections ξ with $\xi(0) = 0$. We ask that this restriction, denoted by*

$$D\bar{\partial}^{res}_{w_0,C} : \Gamma(w_0^*TC, w_0^*TL_0)^{res} \longrightarrow \Omega^{0,1}(w_0^*TC), \tag{8.18}$$

  should have one-dimensional cokernel (in suitable Sobolev spaces).
- *Moreover,*
  - (i) *every holomorphic disc w in (C, L_0) with $w \cdot C' = w \cdot X_0 = 1$ and $w(0) \in C' \cap X_0$ is equal to w_0, up to a possible action of $S^1 \subset Aut(D)$;*
  - (ii) *there are no holomorphic discs w in (C, L_0) with $w \cdot C' = 0$, $w \cdot X_0 = 1$;*
  - (iii) *there are no holomorphic spheres v in C with $\int v^* \omega_C = 1$.*

One can put L_0 into a smooth family $\{L_s\}$ (for $s \in \mathbb{R}$, $|s|$ small) of Lagrangian submanifolds, $L_s \subset X_{1/s}$, such that the union $L = \bigcup_s L_s$ is Lagrangian (but of course not closed) in X. In fact, the condition on L determines the L_s uniquely, since it means that they get mapped into each other by symplectic parallel transport in $\mathbb{R}$-direction.

Lemma 8.16. *For each $s \neq 0$, L_s is an exact Lagrangian submanifold in the affine Kähler surface $M_{1/s}$, and graded with respect to $\eta_{X_{1/s}}$.*

Proof. Supposing that $s \neq 0$ remains sufficiently small, one has $L_s \cap X_0 = \emptyset$, and since $X_{1/s} \cap X_0 = X_{1/s} \cap X_\infty$, also $L_s \cap X_\infty = \emptyset$. Because $o_X|L_\infty$ is trivial and L is Lagrangian, $o_X|L_s$ is trivial for any s. Moreover, its covariantly constant sections lie in the same homotopy class of nowhere zero sections as $\sigma_{X,0}|L_s$, because that is true for $s = 0$. On the other hand, $\sigma_{X,\infty}/\sigma_{X,0}$ is constant on L_s, so the covariantly constant sections are also homotopic to $\sigma_{X,\infty}|L_s$, which is equivalent to exactness. The argument with the grading is essentially the same. □

Exactness of L_0 with respect to θ_C means that the area of holomorphic discs in (C, L_0) is integral. Our disc w_0 has minimal area, since by assumption

$\int w_0^*\omega_C = w_0 \cdot X_0 = 1$. By [38, 39], w_0 has a somewhere injective point. Standard transversality arguments tell us that there is a family $(J_{C,r})$, for $r \in \mathbb{R}$ with $|r|$ small, of compatible almost complex structures on C such that $J_{C,0}$ is the given complex structure, and which makes w_0 parametrized-regular. Since the set of somewhere injective points is open, and w_0 intersects $C' \cup X_0$ only in $w_0(0)$, one can assume that all $J_{C,r}$ are equal to $J_{C,0}$ near $C \cap (C' \cup X_0)$. Having that, one can extend them to a family $(J_{X,r})$ of compatible almost complex structures on X, such that again $J_{X,0}$ is the given complex structure, and with the following two properties for all r: $J_{X,r} = J_{X,0}$ near X_0; and the projection $\pi^{-1} = \sigma_{X,\infty}/\sigma_{X,0} : X \setminus X_0 \to \mathbb{C}$ is $J_{X,r}$-holomorphic.

Consider the moduli space $\mathfrak{M}^{para}$ of pairs (r, w), where $w : (D, \partial D) \to (X, L)$ is a $J_{X,r}$-holomorphic disc such that $w \cdot X_0 = 1$ and $w(0) \in X_0 \cap C'$, divided by the action of S^1 by rotation on D.

LEMMA 8.17. *For every (r, w) in this moduli space, the image of w is contained in some fibre $X_{1/s}$ of the Lefschetz pencil, with $s \in \mathbb{R}$.*

PROOF. Consider the meromorphic map $\pi^{-1} \circ w : D \dashrightarrow \mathbb{C}$. w intersects X_0 only at $w(0)$ and with multiplicity one. At the same point it intersects X_∞, and this implies that $\pi^{-1} \circ w$ is in fact holomorphic. Since its boundary values are real, it must be constant. ☐

LEMMA 8.18. *Let (r_k, w_k) be a sequence of pairs, where $w_k : (D, \partial D) \to (X, L)$ is a J_{X,r_k}-holomorphic disc with $\int w_k^*\omega_X = 1$, whose image is contained in X_{1/s_k} for some $s_k \neq 0$. Suppose that $r_k, s_k \to 0$; then the w_k, after a possible reparametrization by automorphisms of D and passing to a subsequence, converge to w_0.*

PROOF. Gromov compactness tells us that the limit of a subsequence is a stable disc in X_∞, but in our case each non-constant component contributes an integral amount of energy, which means that there can be only one of them. Furthermore, the case where the non-constant component is a sphere is excluded by part (iii) of our assumptions. We conclude that after passing to a subsequence, the convergence is in C^∞-sense towards a limit which is a holomorphic disc $w : (D, \partial D) \to (C, L_0)$.

We know that $w \cdot X_0 = 1$, and (ii) says that w must intersect C'. Every $z \in D$ such that $w(z) \in C'$ must also satisfy $w(z) \in X_0$, otherwise the disc w could not be the smooth limit of discs in the fibres X_{1/s_k}. In fact, a closer look at the local geometric situation shows that $w \cdot C' \leq w \cdot X_0$. Hence $w \cdot X_0 = w \cdot C' = 1$, and after reparametrization we may assume that $w(0) \in X_0 \cap C'$, so an appeal to (i) finishes the argument. ☐

For the next step, we need to compare the linearized operators associated to w_0 as a holomorphic disc in C and X. From (8.16) one gets a diagram with exact columns,

(8.19)
$$\begin{tikzcd}
0 \arrow[d] & 0 \arrow[d] \\
\Gamma(w_0^*TC, w_0^*TL_0) \arrow[r, "D\bar\partial_{w_0,C}"] \arrow[d] & \Omega^{0,1}(w_0^*TC) \arrow[d] \\
\Gamma(w_0^*TX, w_0^*TL) \arrow[r, "D\bar\partial_{w_0,X}"] \arrow[d] & \Omega^{0,1}(w_0^*TX) \arrow[d] \\
\Gamma(w_0^*\mathcal{O}_C(C \cap X_0 - C \cap C'), w_0^*\Lambda) \arrow[r, "\bar\partial"] \arrow[d] & \Omega^{0,1}(w_0^*\mathcal{O}_C(C \cap X_0 - C \cap C')) \arrow[d] \\
0 & 0
\end{tikzcd}$$

where Λ is the canonical real subbundle of $\mathcal{O}_C(C \cap X_0 - C \cap C')|L_0$, whose existence is due to the fact that L_0 is disjoint from $X_0 \cup C'$. As usual, we use suitable Sobolev spaces to make each row elliptic. The bottom row is a $\bar\partial$-operator on a line bundle over a disc: in fact, since w_0 intersects both X_0 and C' only at $z = 0$ at with multiplicity one, this is just the standard $\bar\partial$-operator for functions with real boundary conditions. Hence, its Sobolev completion is onto with one-dimensional kernel, consisting of nowhere zero functions. Without losing exactness of the columns, we may restrict the domains of the top two $D\bar\partial$-operators to sections with $\xi(0) \in T(X_0 \cap C')$ (this tangent space is zero for the first row, and two-dimensional for the second one). Kernels and cokernels of the restrictions will fit into a long exact sequence

$$0 \to \ker D\bar\partial^{res}_{w_0,C} \to \ker D\bar\partial^{res}_{w_0,X} \to \mathbb{R} \to \operatorname{coker} D\bar\partial^{res}_{w_0,C} \to \operatorname{coker} D\bar\partial^{res}_{w_0,X} \to 0.$$

From this and (8.17) one sees that the expected or virtual dimension of the moduli space $\mathfrak{M}^{para}$ at $(0, w_0)$ is

$$\text{v.dim}\, \mathfrak{M}^{para} = \operatorname{ind} D\bar\partial^{res}_{w_0,X} + 1 - 1 = \operatorname{ind} D\bar\partial_{w_0,C} - 3 = 2 + \mu(w_0) - 3 = 1.$$

(in the second expression, the $+1$ comes from the added parameter r, and the -1 from the action of S^1 by rotations). By assumption $\operatorname{coker} D\bar\partial^{res}_{w_0,C} \cong \mathbb{R}$, and by construction of $\{J_{C,r}\}$, this cokernel is spanned by $(\partial J_{C,r}/\partial r)_{r=0} \circ dw_0 \circ i$. Hence, the cokernel of $D\bar\partial^{res}_{w_0,X}$ is spanned by $(\partial J_{X,r}/\partial r)_{r=0} \circ dw_0 \circ i$. This means that the corresponding parametrized linearized operator is onto, so that $(0, w_0)$ is a regular point of $\mathfrak{M}^{para}$.

Lemma 8.19. *There is a family (r_t, w_t) of regular points of $\mathfrak{M}^{para}$, for $t \in \mathbb{R}$ with $|t|$ small, with $r_0 = 0$ and w_0 our given holomorphic disc. This satisfies $w_t(D) \subset X_{1/s_t}$ for some $s_t \in \mathbb{R}$ smoothly depending on t, and with $s_0 = 0$, $(ds_t/dt)_{t=0} \neq 0$.*

Proof. We know that $\mathfrak{M}^{para}$ is locally a smooth one-manifold, so take a family (r_t, w_t) which parametrizes it locally. The fact that $w_t \in X_{1/s_t}$ for some s_t comes from Lemma 8.17, and the smoothness of $t \mapsto s_t$ is obvious. Now suppose that $(ds_t/dt)_{t=0}$ vanishes. Then $(\partial r_t/\partial t, \partial w_t/\partial t)_{t=0}$ lies in the kernel of the parametrized version of $D\bar\partial^{res}_{w_0,C}$. By construction this is onto, and for index reasons the only nontrivial kernel elements are those obtained from infinitesimal rotations of D. This contradicts the fact that we started with a local chart for $\mathfrak{M}^{para}$. □

Lemma 8.20. *For all small $t \neq 0$, w_t is parametrized-regular as a J_{r_t}-holomorphic disc in $(X_{1/s(t)}, L_{s(t)})$.*

The proof is by a comparison argument using a diagram of $\bar\partial$-operators similar to (8.19); we leave it to the reader. Now fix some small $\epsilon > 0$ and $0 \neq t \ll \epsilon$. Then in the family $(J_{X,r}|X_{1/s(t)})$, $r \in [-\epsilon; \epsilon]$, of almost complex structures there is no pseudo-holomorphic disc in $(X_{1/s(t)}, L_{s(t)})$ with unit area other than w_t. In fact, the contrary assumption for very small ϵ and t leads to a contradiction with Lemma 8.18 and the local structure of the moduli space $\mathfrak{M}^{para}$. By Lemma 8.20, this situation persists under a small perturbation of our family of almost complex structures; by such a perturbation, we can make it regular with respect to $L_{s(t)}$ at the endpoints $r = \pm\epsilon$, and eliminate pseudo-holomorphic spheres in $X_{1/s(t)}$ for all values of r. We are now in the situation of Corollary 8.12, which shows that $\mathcal{F}(X_{1/s(t)}, X_{1/s(t),\infty})$ is nontrivial at first order. Using Remark 8.5 and symplectic parallel transport one can extend this to other fibres of our pencil, so the conclusion is:

Corollary 8.21. *Suppose that Assumptions 8.15 hold. Take a smooth fibre X_z of our pencil of Calabi-Yau surfaces. Then its relative Fukaya category $\mathcal{F}(X_z, X_{0,\infty})$, seen as a one-parameter deformation of A_∞-categories, is nontrivial at first order.* □

9. Computations in Fukaya categories

To effect the link between symplectic geometry and homological algebra, we basically rely on two tools. The first is the correspondence between Dehn twists and algebraic twists, which is a reformulation of the long exact sequence in Floer cohomology. The version in the literature [60, 61] is for the exact or affine situation, but as will be outlined below, the proof can be easily adapted to the projective and relative cases. Our second tool is a dimensional induction machine based on the notion of matching cycle, developed in [61]; this is strictly limited to the affine context, which is the reason why affine varieties appear in this paper at all. We will treat dimensional induction as a black box, stating its properties in an axiomatic way only; ultimately it is again an application of the long exact sequence, but the way in which this works is too complicated to be summarized briefly.

(9a) Let M be an affine Calabi-Yau of dimension $n > 1$, and $L \subset M$ a Lagrangian sphere. Equip it with a grading and the unique *Spin* structure, making it into an object $L^\flat$ of $\mathcal{F}(M)$. Let τ_L be the Dehn twist along L, and $\tilde{\tau}_L$ its canonical grading (again, there is the small matter of framing, which we ignore). Consider also the algebraic twist $T_{L^\flat}$ along $L^\flat$, which is a map of objects of $D^b\mathcal{F}(M)$. Let $L_1^\flat$ be any exact Lagrangian brane, with underlying Lagrangian submanifold L_1. Clearly $\tau_L^{-1}(L_1)$ carries an induced brane structure, making it into an object $\tilde{\tau}_L^{-1}(L_1^\flat)$ of the Fukaya category. What [61] says is that

Proposition 9.1. *$T_{L^\flat}(\tilde{\tau}_L^{-1}(L_1^\flat))$ is isomorphic to $L_1^\flat$ in $D^b\mathcal{F}(M)$.*

On the algebraic side, this is an application of Lemma 2.6, where W is some other exact Lagrangian brane $L_0^\flat$; we need to explain how a suitable moduli space setup provides the necessary pair (k,h). This is a variation of the construction in [60], based on pseudo-holomorphic sections of exact symplectic Lefschetz fibrations with Lagrangian boundary conditions. Specifically, taking $S = \mathbb{H}$ the upper half plane, there is such a fibration $E_L \to S$, which has a single critical point somewhere over $int(S)$. The regular fibres of E_L are isomorphic to M; the monodromy around the singularity is of course τ_L; and there is a canonical trivialization away from a compact subset of S. Given any exact Lagrangian brane in M such as $L_1^\flat$, one can define a subbundle $F_{L_1} \subset E_L|\partial S$ which, in the canonical trivialization, satisfies

$$F_{L_1} \cap E_s = \begin{cases} \tau_L^{-1}(L_1) & s \ll 0, \\ L_1 & s \gg 0. \end{cases}$$

By considering an equation similar to (8.5) for sections of E_L with boundary values on F_{L_1}, one then obtains a “relative invariant” in the style of [48] taking values in $HF^0(\tilde{\tau}_L^{-1}(L_1^\flat), L_1^\flat)$. Take k to be the cocycle representing it, which of course depends on the specific choices of almost complex structures etc. used in the definition.

For h one uses a parametrized version of the same construction, applied to a family of Lefschetz fibrations E^t with boundary conditions F^t, $t \in \mathbb{R}$. Figure 1 shows the asymptotic behaviour of this family: for $t \ll 0$, (E^t, F^t) is the fibre connected sum of (E_L, F_{L_1}) and a trivial fibration over the three-punctured disc, with boundary conditions over the different boundary components given by the Lagrangian submanifolds L, $\tau_L^{-1}(L_1)$ and L_1. The invariant associated to the degenerate limit $t = -\infty$ is therefore the composition

$$\begin{aligned} CF^*_M(L^\flat, \tilde{\tau}_L^{-1}(L_1^\flat)) &\longrightarrow CF^*_M(L^\flat, L_1^\flat), \\ a &\longmapsto \mu^2_M(a,k). \end{aligned} \tag{9.1}$$

For $t \gg 0$, (E^t, F^t) again decomposes as a connected sum. Its first component is E_L as before, but now with boundary condition F_L; and the second component is trivial, with boundary conditions given by L, $\tau_L^{-1}(L_1)$ and again L. In principle, the associated invariant is

$$\begin{aligned} CF^*_M(L^\flat, \tilde{\tau}_L^{-1}(L_1^\flat)) \cong CF^*_M(\tilde{\tau}_L(L^\flat), L_1^\flat) &\longrightarrow CF^*_M(L^\flat, L_1^\flat), \\ a &\longmapsto \mu^2_M(l, a), \end{aligned} \tag{9.2}$$

where $l \in CF^0(L^\flat, \tilde{\tau}_L(L^\flat)) \cong CF^0(\tilde{\tau}^{-1}(L^\flat), L^\flat)$ is obtained from (E_L, F_L). However, since $\tilde{\tau}_L(\tilde{L}) = \tilde{L}[1-n]$ with $n > 1$, we have $HF^0(L^\flat, \tilde{\tau}_L(L^\flat)) = HF^{1-n}(L^\flat, L^\flat) \cong H^{1-n}(S^n; \mathbb{C}) = 0$, and by suitably choosing the Floer data, one can achieve that $CF^0(L^\flat, \tilde{\tau}_L(L^\flat))$ vanishes too. Hence l and the expression (9.2) are both zero, and from this one deduces that the parametrized moduli space of sections associated to the deformation from $t = -\infty$ to $t = +\infty$ yields a chain homotopy h between (9.1) and zero, as desired (this vanishing argument, which relies on the grading, is different from the one in [60, Proposition 2.13]; we use it because it extends easily to the projective and relative cases).



Figure 1.

Since all Lagrangian submanifolds involved are exact, there are real-valued action functionals for all the Floer cochain groups involved, so that in the relevant complex (2.5) one can introduce an $\mathbb{R}$-filtration by the values of the action. Its acyclicity is then proved by a spectral sequence argument. Briefly, one replaces the $\mathbb{R}$-filtration by a $\mathbb{Z}$-subfiltration, which divides the action values into intervals of some suitably chosen small size $\epsilon > 0$. After passing to the associated graded space of the filtration, one finds that the Floer differentials μ^1 and the terms $\mu^2(h(a_2), a_1)$, $\mu^3(k, a_2, a_1)$ disappear, while the remaining terms $\mu^2(a_2, a_1)$ and $\mu^2(k, b)$ make the complex exact, hence acyclic. For this to work, some care needs to be observed when choosing the map τ_L and the cochain level representatives for k and h, see [60, Chapter 3] for details.

(9b) We now turn to the consequences of Proposition 9.1, combining arguments from [61] with the notion of negative graded symplectic automorphisms introduced in Section 7.

Lemma 9.2. *Suppose that $m_X = 0$, so M is the affine part of a projective Calabi-Yau X. Let $L_1, \dots, L_m \subset M$ be Lagrangian spheres, and suppose that*

$$\tilde{\tau}_{L_1} \dots \tilde{\tau}_{L_m} \in \widetilde{Aut}(M)$$

is isotopic, within that group, to a graded symplectic automorphism $\tilde{\phi}$ whose extension to X is negative (see Remark 6.10). Then $L_1^\flat, \dots, L_m^\flat$ are split-generators for $D^\pi\mathcal{F}(M)$.

Proof. Let $L^\flat$ be any exact Lagrangian brane. From Lemma 7.2 we know that if d is sufficiently large, a small exact perturbation L' of L will cause the Maslov index of any intersection point $x \in L' \cap \phi^d(L)$ to be strictly negative. By definition of the Floer cochain space (8.2), it follows that

$$Hom_{H^0\mathcal{F}(M)}(L^\flat, \tilde{\phi}^d(L^\flat)) = HF^0_M(L^\flat, \tilde{\phi}^d(L^\flat)) = 0.$$

On the other hand, $\tilde{\phi}^d(L^\flat)$ is isomorphic in $D^b\mathcal{F}(M)$ to $(T_{L_1^\flat} \dots T_{L_m^\flat})^d(L^\flat)$. Lemma 2.7 says that $L^\flat$ lies in the subcategory of $D^\pi\mathcal{F}(M)$ split-generated by the $L_k^\flat$; by construction of that category, this implies that these objects are split-generators. □

Remark 9.3. *In Lemma 9.2, one can replace the negativity of $\tilde{\phi}$ by the following slightly weaker assumption. There is a closed subset $\Sigma \subset X$ which is a finite union of real submanifolds of dimension $< n$, which admits arbitrarily small open neighbourhoods W satisfying $\phi(W) = W$, and such that $\tilde{\phi}$ is negative on $X \setminus W$. The reason is that by a generic perturbation, one can make Λ disjoint from Σ. Then it is disjoint from a sufficiently small W, and the same will hold for all iterates $\phi^d(\Lambda)$, so that the breakdown of negativity on Σ can be disregarded.*

(9c) To adapt Proposition 9.1 to the case of a projective Calabi-Yau surface X, with L a Lagrangian sphere and $L_1^\sharp$ a rational brane, two points have to be taken into account. Firstly, more care needs to be exercised when defining the invariants derived from pseudo-holomorphic sections, to avoid bubbling off of holomorphic discs. This is not really new, since we have dealt with the same problem in the construction of $\mathcal{F}(X)$. Note that up to shifts, L gives rise to a unique object $L^\sharp$ of the Fukaya category, by Lemma 8.13. For $\tilde{\tau}_L^{-1}(L_1^\sharp)$, one needs to choose the regular almost complex structure which is the τ_L-pullback of that associated to the L_1. The second modification concerns the filtration argument. Our ground ring is now $\Lambda_{\mathbb{Q}}$, and each pseudo-holomorphic section contributes $\pm q^e$ as in (8.10). In the $\mathbb{R}$-filtration of the Floer cochain groups, one now uses powers of q as a replacement for values of the action functional. Since only a finite number of rational branes are involved ($L^\sharp$, $L_0^\sharp$, $L_1^\sharp$), one can actually work over a subfield $\Lambda_{(1/d)\mathbb{Z}} \subset \Lambda_{\mathbb{Q}}$ containing only roots of q of some fixed order d. An essentially equivalent and more familiar formulation is to say that all action functionals together have only a discrete set of periods. This allows one to carry over the previous argument, with its use of a $\mathbb{Z}$-subfiltration: it suffices to make sure that $\epsilon \ll 1/d$ (actually, this is the only point in the entire paper that would have to be modified substantially if one wished to include non-rational branes in $\mathcal{F}(X)$). The outcome is

PROPOSITION 9.4. *$T_{L^\flat}(\tilde{\tau}_L^{-1}(L_1^\flat))$ is isomorphic to $L_1^\flat$ in $D^b\mathcal{F}(X)$.* $\square$

There is also a version for the relative Fukaya category, where L is again a Lagrangian sphere in M and $L_1^\natural$ an object of that category. The definition of the relevant version of (k,h) is as in the projective case, except sections u now contribute with integer powers of q according to their intersection with X_∞. As for the acyclicity of (2.5), one filters it by powers of q, and then taking the associated graded space brings one back to the affine situation, where we already know that acyclicity holds.

PROPOSITION 9.5. *$T_{L^\natural}(\tilde{\tau}_L^{-1}(L_1^\natural))$ is isomorphic to $L_1^\natural$ in $D^b\mathcal{F}(X, X_\infty)$.* $\square$

The argument from Lemma 9.2 carries over to the projective and relative cases without any modifications, and so does Remark 9.3. By combining this with Proposition 7.22, one arrives at the following conclusion:

COROLLARY 9.6. *Let X be a projective Kähler threefold which is Fano, with $o_X = \mathcal{K}_X^{-1}$, equipped with sections $\sigma_{X,0}$, $\sigma_{X,\infty}$ which generate an almost Lefschetz pencil of Calabi-Yau surfaces $\{X_z\}$. Assume that $X_0 \cap X_\infty$ has no rational components. Pick some base point $z_* \in \mathbb{C} \setminus Critv(\pi_M)$, base path c_*, and a distinguished basis of vanishing cycles $V_{c_1}, \dots, V_{c_r} \subset M_{z_*}$. Then the V_{c_i}, equipped with some gradings, are split-generators for all versions of the split-closed derived Fukaya category: $D^\pi\mathcal{F}(M_{z_*})$, $D^\pi\mathcal{F}(X_{z_*})$ and $D^\pi\mathcal{F}(X_{z_*}, X_{z_*,\infty})$.* $\square$

(9d) Let $M = X \setminus X_\infty$ be an affine Calabi-Yau of dimension $n > 1$, equipped with a quasi-Lefschetz pencil $\{X_z\}$ and the associated holomorphic function $\pi_M : M \to \mathbb{C}$. Choose a base point z_* and base path c_*. Let $c_1, \dots, c_r$ be a basis of vanishing paths, and $V_{c_1}, \dots, V_{c_r} \subset M_{z_*}$ the associated vanishing cycles. Make them into objects $V^\flat_{c_i}$ of $\mathcal{F}(M_{z_*})$; any choice of grading is allowed, and the *Spin* structure is unique except for $n = 2$, where one needs to take the *nontrivial* one. Let

$$\mathcal{F}^\to = \mathcal{F}^\to(V^\flat_{c_1}, \dots, V^\flat_{c_r}) \tag{9.3}$$

be the directed A_∞-subcategory of $\mathcal{F}(M_{z_*})$ consisting of these objects. We associate to each embedded vanishing path c, subject to the conditions $im(c) \cap im(c_*) = \{z_*\}$ and $\mathbb{R}^{\geq 0} c'(0) \neq \mathbb{R}^{\geq 0} c'_*(0)$, an object D_c of $D^b(\mathcal{F}^\to)$, unique up to isomorphism and shifts, according to the following rules:

- to c_i belongs the object $D_{c_i} = V^\flat_{c_i}$ of $\mathcal{F}^\to$ itself;
- an isotopy of c within the class of such paths does not affect D_c;
- if we have vanishing paths c, c', c'' as in Figure 2, then $D_{c'} \cong T_{D_c}(D_{c''})$ and $D_{c''} \cong T^\vee_{D_c}(D_{c'})$.

From this and the elements of the theory of mutations [55], one sees that all the D_c are exceptional objects, which means that $Hom^*(D_c, D_c)$ is one-dimensional in degree zero, and zero in all other degrees.

Next, we associate objects S_d of $D^b(\mathcal{F}^\to)$ to matching paths which are disjoint from c_*. For this, take paths c', c'', d as in Figure 3. Suppose that the vanishing cycles associated to c', c'' are isotopic, so that d is a matching paths. The objects associated to c', c'' will satisfy

$$Hom^k_{D^b(\mathcal{F}^\to)}(D_{c'}, D_{c''}) = \begin{cases} \mathbb{C} & k = l, l+n-1, \\ 0 & \text{otherwise} \end{cases} \tag{9.4}$$

for some $l \in \mathbb{Z}$. After a shift, we may assume that $l = 0$, and then the rule is:

- S_d is the mapping cone $\{D_{c'} \to D_{c''}\}$ over some nontrivial morphism.

An easy computation shows that $Hom^*(S_d, S_d)$ is one-dimensional in degrees 0 and n, and zero otherwise. This resemblance to the cohomology of an n-sphere is not accidental:

THEOREM 9.7. *Let $d_1, \dots, d_m$ be matching paths which are disjoint from our base path c_*. Assign objects $S_{d_1}, \dots, S_{d_m}$ in $D^b(\mathcal{F}^\to)$ as explained above. Then the full A_∞-subcategory of $Tw\mathcal{F}^\to$ consisting of these objects is quasi-isomorphic to the full subcategory of the Fukay category of the total space, $\mathcal{F}(M)$, whose objects are the matching cycles $\Sigma_{d_1}, \dots, \Sigma_{d_m}$, with some choice of gradings.* □



Figure 2.



Figure 3.

This way of stating the result leaves several things implicit. Since one can get any vanishing path by starting with $c_1, \dots, c_r$ and applying the moves shown in Figure 3, our rules determine the objects D_c. However, it is not a priori clear that the assignment $c \mapsto D_c$ is unambiguously defined, since one can get the same path by different sequences of moves. Similarly, (9.4) and the well-definedness of S_d are not obvious. The fact that all of this works is a nontrivial part of the statement. For proofs see [61].

(9e) The usefulness of Theorem 9.7 is particularly obvious for $n = 2$, when the vanishing cycles are curves on a surface, since then $\mathcal{F}^{\rightarrow}$ can be computed in a purely combinatorial way, up to quasi-isomorphism (versions of this fact have been discovered independently in various contexts, see e.g. [63, 50, 10]). We will now outline how to do this computation, referring to the papers just quoted and to [61] for further details and justification.

Exactness: For simplicity, we use a compact piece $N \subset M_{z_*}$ which is a surface with boundary, capturing all of the topology of M_{z_*} (the inclusion is a homotopy equivalence). Draw simple closed curves ν_k on N which are isotopic to V_{c_k} and in general position (any two intersect transversally, and there are no triple intersections). One should then verify that these curves are exact Lagrangian submanifolds for some choice of one-form θ_N with $d\theta_N > 0$. For this, consider the decomposition of N given by the graph $\Gamma = \bigcup_i \nu_i$. Suppose that one can associate to each component $Z \subset N \setminus \Gamma$ which is disjoint from ∂N a positive weight $w(Z) > 0$, with the following property: if $\sum_k m_k \bar{Z}_k$, $m_k \in \mathbb{Z}$, is a two-chain whose boundary is a linear combination of the ν_k, then

$$\sum_k m_k w(Z_k) = 0. \tag{9.5}$$

One needs to check this only for a finite number of chains, which form a basis for the homology relations between the ν_k. If this works out, one can find a symplectic form ω_N with $\int_Z \omega_N = w(Z)$, and that can be written as $\omega_N = d\theta_N$ with $\int_{\nu_k} \theta_N = 0$ for all k.

Remark 9.8. *This "painting by numbers" step is rather tedious, so we will mention a way of reducing the amount of work. Relations in homology between the vanishing cycles correspond to elements of $H_2(M)$ (because $H_2(M_{z_*}) = 0$, and M is obtained topologically by attaching two-cells to the vanishing cycles). It is sufficient to check* (9.5) *for relations corresponding to elements in $im(\pi_2(M) \to H_2(M))$. Equivalently, these are relations obtained from maps $\Sigma \to M_{z_*}$ where Σ is a genus zero surface with boundary, each boundary component getting mapped to some ν_i. This weaker condition does not guarantee that the vanishing cycles can be made exact, but it allows one to define the directed Fukaya category, and moving the ν_i within that class does not affect the quasi-isomorphism type of the category. Hence, after a (possibly non-Hamiltonian) isotopy, one sees that the directed Fukaya category is the same as for the exact choice of vanishing cycles. The appearance of $\pi_2(M)$ should remind the reader of the assumption $\omega|\pi_2(M) = 0$ used frequently in symplectic topology, and indeed it addresses the same issue (getting a bound on the energy of pseudo-holomorphic polygons).*

Gradings: Recall that the holomorphic one-form $\eta_{M_{z_*}}$ defines a phase function $\alpha_{M_{z_*}}$. One can associate to this an unoriented foliation of M_{z_*}, consisting of those tangent lines Λ for which $\alpha_{M_{z_*}}(\Lambda) = 1$. We draw some unoriented foliation $\mathcal{F}_N$ on N which is homotopic to the one determined by $\alpha_{M_{z_*}}$. To equip the ν_k with gradings, one has to choose a point $b_k \in \nu_k$ which should not be an intersection point with any other ν_l, and a homotopy between $\mathcal{F}_N$ and $T\nu_k$ at that point (within the $\mathbb{RP}^1$ of lines in TN_{b_k}). This determines the Maslov indices $I(x)$ of all intersection points $x \in \nu_i \cap \nu_j$, $i < j$, and also orientations of the ν_i. Note that $(-1)^{I(x)+1}$ is the local intersection number $(\nu_i \cdot \nu_j)_x$.

Spin structures: For every x as before with $(\nu_i \cdot \nu_j)_x = +1$, we choose a nonzero tangent vector ξ_x to ν_j at x. Of course, one can take these to be the vectors pointing along the orientation, but that is not strictly speaking necessary. Next, recall that we have to equip the ν_i with nontrivial *Spin* structures, or what is the same, with nontrivial double covers. Having already chosen points b_k, the simplest way to do this is to take the double cover which is trivial on $\nu_k \setminus b_k$, and whose two sheets get exchanged when passing over that point.

Counting polygons: We now introduce a directed A_∞-category $\mathcal{C}^{\to}$ with objects $\nu_1, \dots, \nu_r$. Because of directedness, it is sufficient to define the $hom_{\mathcal{B}}(\nu_i, \nu_j)$ for $i < j$, and the composition maps between those spaces. Set as usual

$$hom^k_{\mathcal{C}^{\to}}(\nu_i, \nu_j) = \bigoplus_{I(x)=k} \mathbb{C}_x,$$

but where now $\mathbb{C}_x$ is identified with $\mathbb{C}$ in a fixed way. Given $i_0 < \dots < i_d$ in $\{1, \dots, r\}$ and intersection points $x_0 \in \nu_{i_0} \cap \nu_{i_d}$, $x_1 \in \nu_{i_0} \cap \nu_{i_1}, \dots, x_d \in \nu_{i_{d-1}} \cap \nu_{i_d}$, one considers oriented immersions $u : \Delta \to N$, where Δ is some model $(d+1)$-gon in the plane with numbered sides and corners. Under u, the sides should go to the ν_{i_k}, and the corners to the x_k; moreover, the immersion should be a local diffeomorphism up to the corners, so that the image has locally an angle $< \pi$. One considers two such u to be equivalent if they differ by a diffeomorphism of Δ. Each equivalence class contributes to $\mu^k_{\mathcal{C}^{\to}}$ by

$$\pm 1 : \mathbb{C}_{x_d} \otimes \dots \otimes \mathbb{C}_{x_1} \to \mathbb{C}_{x_0},$$

and the sign is the product of the following factors: (i) for each $k \geq 1$ such that $(\nu_{i_{k-1}} \cdot \nu_{i_k})_{x_k} = 1$, we get a -1 iff the tangent vector ξ_{x_k} points away from $u(\Delta)$. (ii) We get another -1 if $(\nu_{i_0} \cap \nu_{i_d})_{x_0} = 1$ and the tangent vector ξ_{x_0} points towards $u(\Delta)$. (iii) Every time that $u|\partial\Delta$ passes over a point b_k, this adds a further -1.

Lemma 9.9. *$\mathcal{C}^{\to}$ is quasi-isomorphic to the directed subcategory $\mathcal{F}^{\to}$ from (9.3), at least up to a shift in the gradings of the $V^\flat_{c_i}$.* □

Starting from this, Theorem 9.7 allows one to determine the full A_∞-subcategory of $\mathcal{F}(M)$ consisting of any finite number of matching cycles, and in particular the Floer cohomology of any two such cycles. Note that matching cycles are Lagrangian two-spheres in a four-manifold, so their Floer cohomology is not amenable to computation in any direct way.

(9f) We will need a minor addition to the previously explained computational method, which takes into account the presence of a homomorphism $\rho : \pi_1(M) \to \Gamma$ into a finitely generated abelian group Γ. Restriction yields a homomorphism from the fundamental group of the fibre M_{z_*}, and all vanishing cycles V_{c_i} bound balls in M, hence the composition $\pi_1(V_{c_i}) \to \pi_1(M_{z_*}) \to \Gamma$ is zero. As discussed in Section 8b, after choosing lifts $V_{c_i,\Gamma}$ to the associated Γ-cover $M_{z_*,\Gamma}$, one has a Γ^*-action on (9.3). We can use the same rules as before to associate to each vanishing path c an object D_c of the equivariant derived category $D^b_{\Gamma^*}(\mathcal{A})$, see Remark 4.1. For c', c'', d are as in Figure 3, the objects $D_{c'}$, the objects $D_{c''}$ will now satisfy

$$Hom^k_{D^b(\mathcal{A})}(D_{c'}, D_{c''}) \cong \begin{cases} W & k = l, l+n-1, \\ 0 & \text{otherwise,} \end{cases}$$

where W is some one-dimensional Γ^*-representation. After shifting $D_{c'}$ and tensoring it with W^*, one may assume that $l = 0$ and W is trivial. Then the cone $S_d = \{D_{c'} \to D_{c''}\}$ over a nontrivial degree zero morphism is again an object of the equivariant derived category. The equivariant version of Theorem 9.7 says that the full A_∞-subcategory of $Tw_{\Gamma^*}\mathcal{F}^{\to}$ with objects $S_{d_1}, \dots, S_{d_m}$ is Γ^*-equivariantly quasi-isomorphic to the full subcategory of $\mathcal{F}(M)$ with objects $\Sigma_{d_1}, \dots, \Sigma_{d_m}$, where we assume that gradings and lifts to M_Γ have been chosen in some appropriate way for these matching cycles. For the proof, which is not particularly difficult, there are two essentially equivalent viewpoints: one either goes through the relevant material in [61], making sure that everything can be made compatible with the Γ^*-actions; or one works directly with the covering M_Γ and its holomorphic function $M_\Gamma \to M \to \mathbb{C}$.

10. The algebras Q_4 and Q_{64}

Using split-generators, the categories which occur in homological mirror symmetry can be described in terms of A_∞-structures on finite-dimensional algebras. This section introduces the algebra relevant for our computation, as well as a simpler version which serves as a stepping-stone. The main point is to use the general theory from Section 3 to partially classify A_∞-structures on those algebras. We end by applying the results to coherent sheaves on the mirror of the quartic surface: this finishes work on the algebro-geometric side of mirror symmetry, and it also helps to clarify the amount of things still to be proved on the symplectic side.

(10a) We begin by recalling a general "doubling" construction. Let $C^{\to}$ be a directed ($\mathbb{C}$-linear graded) category, with objects $(X_1, \dots, X_m)$. The *trivial extension category* of degree $d > 0$ is a category C with the same objects, and it satisfies:

- The directed subcategory of C is the given $C^{\to}$.
- C is a Frobenius category of degree d, in the sense that there are linear maps $\int_{X_j} : Hom^d_C(X_j, X_j) \to \mathbb{C}$, such that

$$\langle a, b\rangle = \int_{X_j} ab \tag{10.1}$$

  for $a \in Hom^*_C(X_k, X_j)$, $b \in Hom^*_C(X_j, X_k)$ is a nondegenerate pairing. These pairings are graded symmetric, $\langle a, b\rangle = (-1)^{deg(a)deg(b)}\langle b, a\rangle$.
- One can split $Hom^*_C(X_j, X_k) = Hom^*_{C^{\to}}(X_j, X_k) \oplus J_{jk}$ in such a way that the product of two morphisms belonging to the J spaces is zero (in view of the previous requirements, we have $J_{jk} = 0$ for $j < k$, $J_{jk} = Hom^*_C(X_j, X_k)$ for $j > k$; and since it is an isotropic subspace, J_{jj} must be one-dimensional and concentrated in degree d).

These properties characterize C up to isomorphism. Alternatively, one can define it explicitly by

$$Hom_C(X_j, X_k) = Hom_{C^{\to}}(X_j, X_k) \oplus Hom_{C^{\to}}(X_k, X_j)^{\vee}[-d] \tag{10.2}$$

and $(a, a^{\vee})(b, b^{\vee}) = (ab, a^{\vee}(b\,\cdot) + (-1)^{deg(a)(deg(b)+deg(b^{\vee}))} b^{\vee}(\cdot\, a))$.

The example relevant to us as is follows: Let V be a four-dimensional complex vector space. Consider the (linear graded) directed category $C_4^{\to}$ having four objects $X_1, \dots, X_4$ and morphisms

$$Hom_{C_4^{\to}}(X_j, X_k) = \begin{cases} \Lambda^{k-j} V & j \leq k, \\ 0 & \text{otherwise}, \end{cases}$$

with the obvious grading and (wedge product) composition. We define C_4 to be the trivial extension category of $C_4^{\to}$ in degree $d = 2$. Its total morphism algebra, linear over $R_4 \cong \mathbb{C}^4$, will be denoted by Q_4. Once one has fixed a choice of volume element in $\Lambda^4 V^{\vee}$, the duals in the general formula (10.2) can be removed, so that then

$$Hom_{C_4}(X_j, X_k) = \begin{cases} \Lambda^{k-j} V & j < k, \\ \Lambda^0 V \oplus (\Lambda^4 V)[2], & j = k, \\ (\Lambda^{k-j+4} V)[2] & j > k. \end{cases} \tag{10.3}$$

From a different viewpoint, take the exterior algebra ΛV, and consider the action of the center of $SL(V)$, $\Gamma_4 = \{I, iI, -I, -iI\}$, on it. For explicitness, pick the generator $\gamma = iI$. From our general discussion of semidirect products, we know that $\Lambda V \rtimes \Gamma_4$ is an algebra over $\mathbb{C}\Gamma_4 \cong R_4$, with a basis of idempotents given by $u_j = \frac{1}{4}\chi_j = \frac{1}{4}(e + i^{-j}\gamma + i^{-2j}\gamma^2 + i^{-3j}\gamma^3)$ for $j = 1, \dots, 4$. Concretely,

$$u_k(\Lambda V \rtimes \Gamma_4)u_j = \begin{cases} \Lambda^{k-j} V \otimes \mathbb{C}u_j & j < k, \\ (\Lambda^0 V \oplus \Lambda^4 V) \otimes \mathbb{C}u_j & j = k, \\ \Lambda^{k-j+4} V \otimes \mathbb{C}u_j & j > k. \end{cases} \tag{10.4}$$

So $\Lambda V \rtimes \Gamma_4$ is the total morphism algebra of a category with four objects. If we view it in this way, the directed subcategory is isomorphic to $C_4^{\to}$. Moreover, after choosing a volume element on V, the resulting integration maps make $\Lambda V \rtimes \Gamma_4$ into a Frobenius category of degree 4. One can split the morphism spaces into those of the directed subcategory and a complementary summand J, consisting of those $\Lambda^j V \otimes \mathbb{C}u_k$ with $j + k > 4$. This satisfies the properties stated above, so we conclude that $\Lambda V \rtimes \Gamma_4$ is the trivial extension category of $C_4^{\to}$ in degree 4. Hence there is an isomorphism $\Lambda V \rtimes \Gamma_4 \cong Q_4$ of algebras, which is R_4-linear but only $\mathbb{Z}/2$-graded.

Later on, we will also need some understanding of the automorphisms of Q_4, as a graded algebra over R_4. To get the trivial thing out of the way, the group $I = (\mathbb{C}^*)^4$ acts by (elementary) inner automorphisms ρ_I, with $\rho_I(r_1, \dots, r_4)$ being multiplication with r_k/r_j on $u_k Q_4 u_j$. Next, the canonical action of $G =$

$GL(V)$ on $C_4^{\to}$ (leaving the objects fixed, and being the obvious representation on $\Lambda^{k-j}V$) extends to an action ρ_G on C_4, determined by asking that

$$\int_{X_j} \rho_G(A)(a) = det(A) \int_{X_j} a. \tag{10.5}$$

With respect to the isomorphism introduced above, this is the same as the G-action on $\Lambda V \rtimes \Gamma_4$ induced by the natural action on V. It is a general feature of semidirect products that $\Gamma_4 \subset G$ acts by inner automorphisms, more precisely

$$\rho_G(iI) = \rho_I(1, i, i^2, i^3).$$

This is the only overlap between ρ_I and ρ_G, which together make up the whole automorphism group. To see this, note that any automorphism of Q_4 determines linear automorphisms of the spaces $u_{j+1}Q_4u_j \cong V$ for all j (in this case, the index is cyclic, so we are thinking of $u_1Q_4u_4$ as well). Moreover, the product of two elements for successive j is zero iff they are linearly dependent in V, and this shows that the actions for various j are by linear maps that differ only by multiplicative constants. By composing the given automorphism with an inner one, these constants can be made equal to 1, and then the remaining automorphism is easily seen to lie in the image of ρ_G. To summarize, we have

$$\begin{aligned} Inn(Q_4/R_4) &\cong (\mathbb{C}^*)^4/\mathbb{C}^*, \\ Out(Q_4/R_4) &\cong GL(V)/\Gamma_4, \\ Aut(Q_4/R_4) &\cong (\mathbb{C}^*)^4/\mathbb{C}^* \times_{\mathbb{Z}/4} GL(V). \end{aligned}$$

At this point, fix a maximal torus $H \subset SL(V)$, and let $T = H/\Gamma_4$ be its image in $PSL(V)$. Choose a splitting $\Sigma : T \to H$ of the projection, and denote by ρ_T the T-action obtained by Σ from ρ_G (this should not be confused with ρ_I). Let Γ_{64} be the subgroup of those $A \in H$ such that $A^4 = I$, and $\Gamma_{16} = \Gamma_{64}/\Gamma_4$ its image in T. We use the restriction of ρ_T to that finite group to form the semidirect product

$$Q_{64} \overset{\text{def}}{=} Q_4 \rtimes \Gamma_{16}.$$

This is an algebra linear over $\mathbb{C}\Gamma_4 \otimes \mathbb{C}\Gamma_{16} = \mathbb{C}\Gamma_{64}$, so one can also view it as a category C_{64} with 64 objects. The actions $\rho_T|\Gamma_{16}$ obtained from any two choices of s differ by homomorphisms $\Gamma_{16} \to Inn(Q_4/R_4)$, which can always be lifted to I. This implies that Q_{64} is independent of Σ up to ($\mathbb{C}\Gamma_{64}$-linear) isomorphism, see (4.1). Moreover, by combining our previous discussion with (4.2) one finds that there is a $\mathbb{Z}/2$-graded isomorphism

$$Q_{64} \cong (\Lambda V \rtimes \Gamma_4) \rtimes \Gamma_{16} \cong \Lambda V \rtimes \Gamma_{64}.$$

(10b) We want to use the interpretation of Q_4 as a twisted exterior algebra to compute its Hochschild cohomology, and that requires paying careful attention to the different gradings. It is convenient to temporarily introduce a bigrading indexed by $\frac{1}{2}\mathbb{Z}\times\mathbb{Z}/2$ on $A = \Lambda V \rtimes \Gamma_4$, with elements of $\Lambda^k V \otimes \mathbb{C}\Gamma_4$ given bidegree $(k/2, [k])$. One defines the Hochschild cohomology of such bigraded algebras by using the $\mathbb{Z}/2$-grading for parities and hence the signs in the Hochschild differential, and the $\frac{1}{2}\mathbb{Z}$-grading to determine the degree of Hochschild cochains, hence the t-part of $HH^{s+t}(A,A)^t$. It is straightforward to adapt the proof of Proposition 4.2 to the new gradings, so we have

$$HH^{s+t}(A,A)^t \cong \bigoplus_{\gamma \in \Gamma_4} \Big(S^s(V^{\gamma,\vee}) \otimes \Lambda^{s+2t-codim(V^\gamma)}(V^\gamma) \Big)^{\Gamma_4}. \tag{10.6}$$

Having done that, we modify the bigrading by adding $((k-j)/2, 0)$ to the bidegree of each piece $(1 \otimes u_k)A(1 \otimes u_j)$, and denote the outcome by B. If we view our algebras as categories, this corresponds to a fractional shift or translation of the j-th object downwards by $j/2$. As a consequence, restriction to the quasi-isomorphic subcomplex of R_4-linear cochains shows that the Hochschild cohomologies of A and B are the same. On the other hand, a direct check using (10.4) shows that B is nontrivial only in bidegrees $(k,[k])$ with $k \in \mathbb{Z}$, so that we can consider it as an ordinary graded algebra. As such, it becomes isomorphic to Q_4, see (10.3); which shows that (10.6) also computes the Hochschild cohomology of that algebra. In particular, the $\gamma \neq e$ summands only contribute to $(s,t) = (0,2)$.

Let $\mathcal{Q}_4$ be an R_4-linear and ρ_T-equivariant A_∞-algebra whose underlying vector space is Q_4, which satisfies (3.1), and which is nontrivial in the sense of not being gauge equivalent to the trivial A_∞-structure. As an aside, note that because the elementary inner automorphisms of Q_4 are also automorphisms of $\mathcal{Q}_4$, ρ_T-equivariance is actually the same as $\rho_G|H$-equivariance, hence independent of Σ. We have no need of proving the existence of such an A_∞-algebra abstractly, since it will be constructed geometrically later on from coherent sheaves or as a Fukaya category. The uniqueness statement is:

LEMMA 10.1. *$\mathcal{Q}_4$ is unique up to (R_4-linear, T-equivariant) A_∞-isomorphism.*

PROOF. We will use the equivariant version of the deformation theory from Section 3, which was discussed extensively in Section 4 in the case of finite groups. We need it for T, but that is not a problem since the T-action on all relevant Hochschild cochain spaces splits them into direct products of subspaces with given weights, which is all we needed. Recall that the HKR isomorphism is G-equivariant. From this and (10.6), one gets

$$\big(HH^2(Q_4,Q_4)^{2-d}\big)^T = \big(S^d V^\vee \otimes \Lambda^{4-d} V\big)^H = \begin{cases} \mathbb{C} & d = 4, \\ 0 & \text{for all other } d > 2. \end{cases}$$

More explicitly, choose an identification $V = \mathbb{C}^4$ making the maximal torus H standard. Denote the standard basis of V by $v_0, \dots, v_3$, and the dual basis by $y_0, \dots, y_3$. Then the only nontrivial summand consists of multiples of the H-invariant polynomial $y_0 y_1 y_2 y_3$. Since $u_k Q_4 u_j$ is nonzero only in degrees $\equiv k-j$ mod 2, any odd degree R_4-bimodule map $Q_4 \otimes_{R_4} \cdots \otimes_{R_4} Q_4 \longrightarrow Q_4$ is necessarily zero. That applies to all the odd order $\mu^d_{\mathcal{Q}_4}$ and in particular to $d = 3$, which shows that $\mathcal{Q}_4$ has a well-defined order 4 deformation class

$$m^4_{\mathcal{Q}_4} \in \big(HH^2(Q_4, Q_4)^{-2}\big)^T.$$

This is nonzero because otherwise, since all the higher obstruction groups vanish, $\mathcal{Q}_4$ would be gauge equivalent to the trivial A_∞-structure, which is contrary to assumption. Using the equivariant version of Lemma 3.2, we find that versality applies: any other A_∞-structure with the same properties as $\mathcal{Q}_4$ is gauge equivalent to $\epsilon^*\mathcal{Q}_4$ for some $\epsilon \in \mathbb{C}^*$, and in particular isomorphic to $\mathcal{Q}_4$. □

Remark 10.2. *One can formulate the previous Lemma as a more general recognition principle. Suppose that we have an A_∞-algebra $\mathcal{A}$ linear over R_4 and carrying a T-action, whose cohomology algebra is equivariantly isomorphic to Q_4. Choose λ, π and h as in Remark 3.1 (and R_4-linearly), so that we can run the explicit form of the Homological Perturbation Lemma to produce an A_∞-structure $\mathcal{B}$ on $H(\mathcal{A})$. This will satisfy $\mu^3_{\mathcal{B}} = 0$ for the same reasons as in the proof above. Take classes $a_4 \in u_1H(\mathcal{A})u_4$, $a_3 \in u_4H(\mathcal{A})u_3$, $a_2 \in u_3H(\mathcal{A})u_2$, $a_1 \in u_2H(\mathcal{A})u_1$, such that the product of any two successive ones is zero. Plug the a_k into the expression* (3.4) *for $\mu^4_{\mathcal{B}}$; if the outcome is nonzero, then $\mathcal{A}$ is quasi-isomorphic to $\mathcal{Q}_4$. To see this, note that by the explicit formula* (4.10) *for the HKR isomorphism, or rather a slight variation of it for semidirect products, $\mu^4_{\mathcal{B}}(a_4, a_3, a_2, a_1) = p(v, v, v, v) \otimes u_1$, where $p \in S^4V^\vee$ is the polynomial corresponding to the obstruction class $m^4_{\mathcal{B}} \in HH^2(Q_4, Q_4)^{-2}$, and $v \in V$ reflects the choice of the a_k. As soon as one knows that $m^4_{\mathcal{B}}$ is nonzero, Lemma 10.1 does the rest.*

(10c) Since $\mathcal{Q}_4$ is ρ_T-invariant, we can form the semidirect product A_∞-algebra $\mathcal{Q}_{64} = \mathcal{Q}_4 \rtimes \Gamma_{16}$, whose underlying algebra we denote by Q_{64}.

Lemma 10.3. *The truncated Hochschild cohomology of $\mathcal{Q}_{64}$ satisfies*

$$HH^1(\mathcal{Q}_{64}, \mathcal{Q}_{64})^{\leq 0} \cong \mathbb{C}^3, \quad HH^2(\mathcal{Q}_{64}, \mathcal{Q}_{64})^{\leq 0} \cong \mathbb{C}^7. \tag{10.7}$$

Proof. We use the spectral sequence (3.9). The same trick of passing through an appropriate bigrading which we used to compute the Hochschild cohomology of Q_4 also works for Q_{64}, yielding

$$\begin{aligned} &HH^{s+t}(Q_{64}, Q_{64})^t \cong \\ &\cong \bigoplus_{\gamma \in \Gamma_{64}} \Big(S^s(V^{\gamma,\vee}) \otimes \Lambda^{s+2t-codim(V^\gamma)}(V^\gamma) \otimes \Lambda^{codim(V^\gamma)}(V/V^\gamma)\Big)^{\Gamma_{64}}. \end{aligned} \tag{10.8}$$

In particular, our spectral sequence satisfies $E_2^{s,t} = 0$ for $s \leq 0$ or $s + 2t < 0$, which ensures its convergence. Case-by-case study of the conjugacy classes in Γ_{64} shows that the summands with nontrivial γ contribute only to $E_2^{s,t}$ with $s+t \geq 3$. Since the differentials in the spectral sequence preserve each summand, we can concentrate on $\gamma = e$. Here is the relevant part of the E_2 term ($S^j\Lambda^k$ is shorthand for $(S^jV^\vee \otimes \Lambda^kV)^{\Gamma_{64}}$, and the crossed out summands vanish because the Γ_4-fixed point sets are already trivial):

	$s=0$	$s=1$	$s=2$	$s=3$	$s=4$	$s=5$	$s=6$
$t=1$	0	$\cancel{S^1\Lambda^3}$	$\cancel{S^2\Lambda^4}$	0	0	0	0
$t=0$	0	$S^1\Lambda^1$	$S^2\Lambda^2$	$S^3\Lambda^3$	$S^4\Lambda^4$	0	0
$t=-1$	0	0	$\cancel{S^2\Lambda^0}$	$\cancel{S^3\Lambda^1}$	$\cancel{S^4\Lambda^2}$	$\cancel{S^5\Lambda^3}$	$\cancel{S^6\Lambda^4}$
$t=-2$	0	0	0	0	$S^4\Lambda^0$	$S^5\Lambda^1$	$S^6\Lambda^2$
$t=-3$	0	0	0	0	0	0	$\cancel{S^6\Lambda^0}$

The first nonzero differential is δ_3, which is the Schouten bracket with the order four deformation class. We identify $V = \mathbb{C}^4$ as before, so that the deformation class is $y_0y_1y_2y_3$ (up to a multiple, which we may safely ignore since it can be absorbed into a rescaling of V). In total degree $s+t=1$, we have the Γ_{64}-invariant part of $V^\vee \otimes V$ which is spanned by elements $y_k \otimes v_k$, and these satisfy

$$\delta_3^{1,0}(y_k \otimes v_k) = y_0y_1y_2y_3 \tag{10.9}$$

for all k. Note that $\frac{1}{2}\sum_k y_k \otimes v_k$ is what we called the Euler element in Section 3, and so the fact that its image under δ_3 is twice the deformation class is an instance of the general property (3.8). For $s+t=2$ we have firstly the Γ_{64}-invariant part of $S^2V^\vee \otimes \Lambda^2V$, generated by the six elements $y_jy_k \otimes v_j \wedge v_k$ which satisfy

$$\delta_3^{2,0}(y_jy_k \otimes v_j \wedge v_k) = (y_0y_1y_2y_3)y_k \otimes v_k - (y_0y_1y_2y_3)y_j \otimes v_j;$$

and secondly the invariant part of S^4V^*, whose basis consists of the y_k^4 together with $y_0y_1y_2y_3$, the latter being annihilated by being in the image of (10.9). The structure of the E_2 term shows that the differentials of order $d > 3$ all vanish, so our computation is complete. To summarize, explicit generators of E_2 which represent the Hochschild cohomology groups in (10.7) are (with some slight redundancy): for HH^1, $y_0 \otimes v_0 - y_1 \otimes v_1$ and its three cyclic permutations (the sum of all four being zero). And for HH^2, $y_0y_1 \otimes v_0 \wedge v_1 + y_12y_2 \otimes v_1 \wedge v_2 + y_2y_0 \otimes v_2 \wedge v_0$ and its cyclic permutations (the sum being again zero), together with the monomials y_k^4, $k = 0, \dots, 3$. $\square$

Consider $U_4 \in Aut(Q_4/R_4)$ of the form $U_4 = \rho_I(r)\rho_G(A)$, where $Av_k = \lambda_kv_{k+1}$, $Av_3 = \lambda_4v_0$ with $\lambda_0\lambda_1\lambda_2\lambda_3 = 1$. Slightly more intrinsically, this class of automorphisms can be characterized by the following properties:

- U_4 acts on the second summand in $u_jQ_4u_j \cong \mathbb{C} \oplus (\Lambda^4V)[2]$ by -1;
- Let $\xi_H : H \to H$ be the shift $diag(t_0, \dots, t_3) \mapsto diag(t_1, t_2, t_3, t_0)$, and ξ_T the induced automorphism of T. Then there is a homomorphism $\iota : \Gamma_{16} \to I$, such that

$$\rho_T(\xi_T(\gamma)) \circ U_4 = \rho_I(\iota(\gamma)) \circ U_4 \circ \rho_T(\gamma). \tag{10.10}$$

> ι is necessary here because the splitting Σ breaks some natural symmetries: essentially, it measures the difference between $\xi_H(\Sigma(\gamma))$ and $\Sigma(\xi_T(\gamma))$.

This is the situation from (4.4), so one can use ι to lift U_4 to an automorphism $U_{64} = U_4 \rtimes (\xi|\Gamma_{16})$ as in (4.5). Suppose that $\mathcal{U}_{64}$ is an A_∞-isomorphism from $\mathcal{Q}_{64}$ to itself whose underlying algebra automorphism is U_{64}, and let $\mathcal{M}_{64}$ be its graph, which is an A_∞-bimodule over $\mathcal{Q}_{64}$. Consider a one-parameter deformation $\mathcal{Q}_{64,q}$ of $\mathcal{Q}_{64}$ which is nontrivial at first order in q, and has the property that $\mathcal{U}_{64}$ can be simultaneously deformed to an A_∞-homomorphism $\mathcal{U}_{64,q}$ from $\mathcal{Q}_{64,q}$ to itself. The existence of these structures is again not an issue, but uniqueness is:

Lemma 10.4. *Up to the pullback action of $End(\Lambda_{\mathbb{N}})^\times$ combined with equivalence of one-parameter deformations, $\mathcal{Q}_{64,q}$ is unique.*

Proof. We use coupled deformation theory for the pair $(\mathcal{Q}_{64}, \mathcal{M}_{64})$. The same parity issues as in Lemma 10.1 show that μ^1 remains zero during the deformation to $\mathcal{Q}_{64,q}$, and that also holds for the differential $\mu^{0,0}$ of the A_∞-bimodule. Here is the relevant piece of the long exact sequence (3.18), with the information from Lemma 3.6 (or rather its analogue for truncated Hochschild cohomology) already built in:

$$\begin{aligned} HH^1(\mathcal{Q}_{64}, \mathcal{Q}_{64})^{\leq 0} &\xrightarrow{1-\mathcal{U}_{64}^*} HH^1(\mathcal{Q}_{64}, \mathcal{Q}_{64})^{\leq 0} \longrightarrow \\ &\longrightarrow HH^2(\mathcal{Q}_{64}\&\mathcal{M}_{64}, \mathcal{Q}_{64}\&\mathcal{M}_{64})^{\leq 0} \longrightarrow \\ &\longrightarrow HH^2(\mathcal{Q}_{64}, \mathcal{Q}_{64})^{\leq 0} \xrightarrow{1-\mathcal{U}_{64}^*} HH^2(\mathcal{Q}_{64}, \mathcal{Q}_{64})^{\leq 0}. \end{aligned} \tag{10.11}$$

At the end of the proof of the previous Lemma, we wrote down four generators which give an H-equivariant isomorphism $HH^1(\mathcal{Q}_{64}, \mathcal{Q}_{64})^{\leq 0} \cong \mathbb{C}^4/\mathbb{C}(1,1,1,1)$; since the coordinates are permuted cyclically by U_{64}, the first arrow in (10.11) is an isomorphism. HH^2 requires slightly more care, since two different pieces of the E_∞ term contribute to it. What the spectral sequence shows is that there is a short exact sequence with an endomorphism,

$$\begin{tikzcd} 0 \arrow[r] & \mathbb{C}^4 \arrow[r] \arrow[d] & HH^2(\mathcal{Q}_{64}, \mathcal{Q}_{64})^{\leq 0} \arrow[r] \arrow[d, "\mathcal{U}_{64}^*"] & \mathbb{C}^4/\mathbb{C}(1,1,1,1) \arrow[r] \arrow[d] & 0 \\ 0 \arrow[r] & \mathbb{C}^4 \arrow[r] & HH^2(\mathcal{Q}_{64}, \mathcal{Q}_{64})^{\leq 0} \arrow[r] & \mathbb{C}^4/\mathbb{C}(1,1,1,1) \arrow[r] & 0 \end{tikzcd}$$

where both the left and right $\downarrow$ are cyclic permutations of the coordinates. This implies that the last arrow in (10.11) has one-dimensional kernel, hence the coupled truncated Hochschild cohomology group is one-dimensional. On the other hand, since $\mathcal{Q}_{64,q}$ is nontrivial at first order, the primary deformation class of any pair $(\mathcal{Q}_{64,q}, \mathcal{M}_{64,q})$ in that group must be nonzero; at which point Lemma 3.8 takes over. □

It is maybe not entirely clear from the statement, but the proof of the Lemma above shows that we were talking about uniqueness of $\mathcal{Q}_{64,q}$ with respect to a fixed choice of U_4, ι and $\mathcal{U}_{64}$. The next results removes the apparent dependence of those data:

LEMMA 10.5. *Up to isomorphism of A_∞-algebras linear over $\Lambda_{\mathbb{N}}$, combined with the action of $End(\Lambda_{\mathbb{N}})^\times$, $\mathcal{Q}_{64,q}$ is independent of all choices made in its construction.*

PROOF. It is convenient to take on the choices in reverse order. (1) Concerning the A_∞-bimodule structure $\mathcal{M}_{64}$ on the fixed bimodule $M_{64} = Graph(U_{64})$, we know from (10.8) and classical homological algebra that

$$Ext^k(M_{64}, M_{64}[1-k]) \cong HH^1(Q_{64}, Q_{64})^{1-k} = 0$$

for all $k \geq 2$, and then Lemma 3.7 implies that $\mathcal{M}_{64}$ is unique up to isomorphism. (2) Any two choices of ι differ by a $\chi \in \Gamma_{16}^*$, and that defines an automorphism of $\mathcal{Q}_{64}$. Since all the Hochschild cohomology in (10.7) lies in the Γ_{16}^*-invariant subspace, one can use the standard quasi-isomorphism theorem of abstract deformation theory [22, Theorem 2.4] to transform a given $\mathcal{Q}_{64,q}$ into an equivalent deformation which is Γ_{16}^*-equivariant. Having done that, it is easy to see that if $\mathcal{U}_{64}$ deforms compatibly to an $\mathcal{U}_{64,q}$ for one choice of ι, then it does for all the others. (3) Changing U_4 by an inner automorphism does not affect its graph (up to bimodule isomorphism), so we may assume that $U_4 = \rho_G(A)$. The remaining freedom is the choice of the λ_k, but one can relate different choices by an automorphism of Q_4 which rescales the different coordinates of V in a suitable way. □

(10d) Take the Beilinson basis of the derived category of coherent sheaves on $P = \mathbb{P}(V)$, which consists of $F_k = \Omega^{4-k}(4-k)[4-k]$ for $k = 1, \dots, 4$. The Ω^k are naturally $PGL(V)$-equivariant sheaves, and if we give $\mathcal{O}(-1)$ the $GL(V)$-action obtained by embedding it into $\mathcal{O} \otimes V$, the $\Omega^{4-k}(4-k) = \Omega^{4-k} \otimes \mathcal{O}(-1)^{\otimes k-4}$ become $GL(V)$-equivariant. A straightforward computation, which is the basis of the argument in [7], shows that

LEMMA 10.6. *The linear graded category with objects $F_1, \dots, F_4$, and where the morphisms are homomorphisms of all degrees in the derived category, is $GL(V)$-equivariantly isomorphic to $C_4^{\rightarrow}$.* □

Let $\iota_0 : Y_0 \hookrightarrow P$ be the unique H-invariant quartic hypersurface, in appropriate coordinates $Y_0 = \{y_0y_1y_2y_3 = 0\}$. Write $E_{0,k} = \iota_0^* F_k$. Repeating an argument from [62], we observe that

$$\begin{aligned} Hom_{Y_0}^*(E_{0,j}, E_{0,k}) &= Hom_P^*(F_j, (\iota_0)_*\iota_0^* F_k) = Hom_P^*(F_j, \mathcal{O}_{Y_0} \overset{\mathbf{L}}{\otimes} F_k) \\ &= Hom_P^*(F_j, \{\mathcal{K}_P \otimes F_k \rightarrow \mathcal{O}_P \otimes F_k\}) \end{aligned}$$

sits in a long exact sequence which one can write, using Serre duality on P, as

$$\cdots \to Hom^d_P(F_j, F_k) \xrightarrow{\iota_0^*} Hom^d_{Y_0}(E_{0,j}, E_{0,k}) \xrightarrow{(\iota_0^*)^\vee} Hom^{2-d}_P(F_k, F_j)^\vee \to \cdots$$

Since the F_k form an exceptional collection, we have unique splittings

$$Hom^*_{Y_0}(E_{0,j}, E_{0,k}) = Hom^*_P(F_j, F_k) \oplus Hom^*_P(F_k, F_j)^\vee[-2]. \tag{10.12}$$

Y_0 has trivial dualizing sheaf, so Serre duality makes $D^bCoh(Y_0)$ into a Frobenius category of degree $d = 2$. If we denote by J_{jk} the second summand in (10.12), then the composition of two morphisms in J spaces is zero for dimension reasons (these morphisms are Ext^2's of locally free sheaves on Y_0). We have now checked off the properties of a trivial extension category. Moreover, the induced $GL(V)$-action on $Hom^*_{Y_0}(E_{0,j}, E_{0,k})$ satisfies (10.5), so

Lemma 10.7. *The linear graded category with objects $E_{0,k}$, and where the morphisms are maps of arbitrary degree in $D^bCoh(Y_0)$, is $GL(V)$-equivariantly isomorphic to C_4.* □

Denote by S_4 the total morphism algebra of that subcategory, so $S_4 \cong Q_4$. This is the cohomology algebra of a dg algebra $\mathcal{S}_4$, canonical up to quasi-isomorphism, which can be defined as total dg algebra of the same objects in the Čech dg category $\check{\mathcal{S}}(Y_0)$. The next step comes from [13]:

Lemma 10.8. *$\mathcal{S}_4$ is not formal, meaning not quasi-isomorphic to its cohomology.*

Proof. The skyscraper sheaf at any point $x = \mathbb{C}v \in P$ admits a locally free Koszul resolution

$$\{\Omega^3(3) \longrightarrow \Omega^2(2) \longrightarrow \Omega^1(1) \longrightarrow \mathcal{O}\} \cong \mathcal{O}_x$$

where each differential corresponds to $v \in V$ under the isomorphism from Lemma 10.6. After restricting, it follows that there is a spectral sequence converging to $Hom^*_{Y_0}(\Omega^3(3)[3], L\iota_0^*\mathcal{O}_x)$ whose E_1 term is

	$s=0$	$s=1$	$s=2$	$s=3$	$s=4$
$t=2$	$\mathbb{C}$	0	0	0	0
$t=1$	0	0	0	0	0
$t=0$	$\mathbb{C}$	V	$\Lambda^2 V$	$\Lambda^3 V$	0

The differential $\delta_1 : E_1^{s,t} \to E_1^{s+1,t}$ is wedge product with v in the bottom row, which leaves just two nonzero terms $E_2^{0,2} \cong \mathbb{C}$, $E_2^{3,0} \cong \mathbb{C}$. If $x \notin Y_0$ then $L\iota_0^*\mathcal{O}_x = 0$, so these two terms must kill each other through the only remaining differential $\delta_3^{0,2}$. On the other hand, if $x \in Y_0$ then $L\iota_0^*\mathcal{O}_x = \mathcal{O}_x \oplus \mathcal{O}_x[1]$ has nonzero sections, so the spectral sequence degenerates at E_2.

The connection with the question we are interested in comes about as follows. If $\mathcal{S}_4$ was formal, the triangulated subcategory of $D^bCoh(Y_0)$ generated by the

$E_{0,k}$ could be modelled by twisted complexes defined on the category C_4 with its trivial A_∞-structure. We know that $L\iota_0^*\mathcal{O}_x$ is a twisted complex of the form $(E_{0,1} \oplus E_{0,2} \oplus E_{0,3} \oplus E_{0,4}, \delta)$, and the behaviour of first differential in the spectral sequence means that δ must be (up to irrelevant scalar multiples)

$$\delta = \begin{pmatrix} 0 & & & \\ v & 0 & & \\ * & v & 0 & \\ * & * & v & 0 \end{pmatrix},$$

But for degree reasons the terms $*$ are necessarily equal to zero. Having identified the object, we can check that in $D^b(C_4)$ the analogue of the spectral sequence above degenerates at E_2 for any nonzero v, which is a contradiction. □

By choosing the Čech covering to be H-invariant, one can arrange that $\mathcal{S}_4$ has a natural H-action. Take the same splitting $\Sigma : T \to H$ as before, and consider the induced T-action. The T-equivariant version of the Homological Perturbation Theorem (which applies because the action on $hom_{\mathcal{S}_4}$ is well-behaved), together with Lemma 10.1, show that $\mathcal{S}_4$ is A_∞-quasi-isomorphic to $\mathcal{Q}_4$, and moreover that quasi-isomorphism is R_4-linear and T-equivariant. As a consequence, the semidirect product $\mathcal{S}_{64} = \mathcal{S}_4 \rtimes \Gamma_{16}$ defined using the restriction of the T-action to Γ_{16} is quasi-isomorphic to $\mathcal{Q}_{64}$. On the geometric side, consider each $E_{0,k}$ as a Γ_{16}-equivariant sheaf in all 16 possible ways (mutually twisted by characters). This gives a total of 64 objects in the derived category $D^bCoh_{\Gamma_{16}}(Y_0)$, and Remark 5.2 tells us that $\mathcal{S}_{64}$ is the dg algebra underlying their total morphism algebra. The flat formal deformation $Y_q = \{p_q(v) = 0\}$ of Y_0 given by (1.1) still admits a $\Lambda_\mathbb{N}$-linear Γ_{16}-action. Since the $E_{0,k}$ are restricted from projective space, they have obvious equivariant extensions $E_{q,k}$ to locally free coherent sheaves over Y_q. By the discussion in Section 5b, or rather its equivariant analogue, this gives rise to a Γ_{16}-equivariant one-parameter deformation $\mathcal{S}_{64,q}$ of $\mathcal{S}_{64}$, which satisfies

$$H^*(\mathcal{S}_{64,q}) \cong H^*(\mathcal{S}_{64}) \otimes_\mathbb{C} \Lambda_\mathbb{N}. \tag{10.13}$$

One can check (10.13) either by repeating the duality argument above for Y_q, or more simply by observing that the morphisms between any two $E_{0,k}$ in $D^bCoh(Y_0)$ are concentrated in a single degree, which precludes the possibility of any deformation of the differential affecting the cohomology.

Lemma 10.9. *$\mathcal{S}_{64,q}$ is nontrivial at first order in q.*

Proof. This is quite similar to the previous Lemma. Let x_q be a $\Lambda_\mathbb{N}$-point of $P \times_\mathbb{C} \Lambda_\mathbb{N}$, which is now given by a family v_q of nonzero vectors in V varying with the formal parameter q. We have the same spectral sequence as before for the derived restriction of its structure sheaf $\mathcal{O}_{x_q}$ to $\iota_q : Y_q \hookrightarrow P \times_\mathbb{C} \Lambda_\mathbb{N}$, and therefore $Hom^*_{Y_q}(\Omega^3(3)[3], L\iota_q^*\mathcal{O}_{x_q})$ is the cohomology of

$$\Lambda_{\mathbb{N}} \xrightarrow{q\mapsto p_q(v_q)} \Lambda_{\mathbb{N}}.$$

$L\iota_q^*\mathcal{O}_{x_q}$ is not an equivariant sheaf, but one can remedy this by summing over the Γ_{16}-orbit of x_q. By taking x_q in such a way that $p_q(v_q)$ has precisely order $O(q)$, one obtains examples of twisted complexes over $\mathcal{S}_{64,q}$ (considered here as an A_∞-category with 64 objects) such that the map (3.22) is not an isomorphism. Now apply Lemma 3.12 to $Tw\,\mathcal{S}_{64,q}$. □

The automorphism $A \in GL(V)$, $Av_k = v_{k+1}$, $Av_3 = Av_0$ preserves Y_0 and the objects $E_{0,k}$, hence defines an automorphism of $\mathcal{S}_4$, unique up to elementary inner automorphisms. Since $det(A) = -1$, this automorphism acts by -1 on the space of sections of the canonical sheaf of Y_0, hence by Serre duality on $Hom^2(E_{0,k}, E_{0,k})$ for any k. Moreover, it skew-commutes with the T-action on Y_0 in the obvious sense, so it can be identified on the level of cohomology with one of the automorphisms U_4 of Q_4 introduced before. Because of the twisted equivariance, one can define an induced automorphism of $\mathcal{S}_{64}$, and since A survives the deformation to Y_q, this can be deformed to an automorphism of $\mathcal{S}_{64,q}$. Together with the previous Lemma, this allows us to apply Lemma 10.4 which shows that $\mathcal{S}_{64,q}$ is quasi-isomorphic to $\mathcal{Q}_{64,q}$, at least in the sense in which the latter object is unique, which means up to reparametrization by $\psi \in End(\Lambda_{\mathbb{N}})^\times$.

Let $Y_q^* = Y_q \times_{\Lambda_{\mathbb{N}}} \Lambda_{\mathbb{Q}}$ be the general fibre of our formal scheme, and $E^*_{q,k}$ the restrictions of the $E_{q,k}$ to it. Let Z_q^* be the minimal resolution of the quotient Y_q^*/Γ_{16}. We need two more basic facts:

Lemma 10.10. *The 64 equivariant versions of the $E^*_{q,k}$ are split-generators for $D^bCoh_{\Gamma_{16}}(Y_q^*)$.*

Proof. Beilinson's resolution of the diagonal [7] shows that the $F^*_{q,k}$ are generators for the derived category of $P \times_{\mathbb{C}} \Lambda_{\mathbb{Q}}$. His argument carries over to the equivariant case, which means that the 64 objects obtained by making the $F^*_{q,k}$ equivariant in all possible ways generate the derived category of Γ_{16}-equivariant sheaves. The equivariant analogue of Lemma 5.4 shows that the restrictions of these objects to Y_q^* split-generate its derived category of equivariant sheaves. □

Lemma 10.11. *$D^bCoh(Z_q^*) \cong D^bCoh_{\Gamma_{16}}(Y_q^*)$, as triangulated categories linear over $\Lambda_{\mathbb{Q}}$.*

This is a theorem of Kapranov and Vasserot [33], valid for any finite group action on a smooth $K3$ surface which preserves the holomorphic two-form. We now have the situation where we know that

$$D^bCoh_{\Gamma_{16}}(Y_q^*) \cong D^\pi(\mathcal{S}(Y_q^*))$$

is split-generated by certain objects, and where we know that the dg subcategory of $\mathcal{S}_{\Gamma_{16}}(Y_q^*)$ consisting of those objects, denoted by $\mathcal{S}_{64,q}$ above, is quasi-isomorphic to $\mathcal{Q}_{64,q}$. Lemma 2.5 then says that

Corollary 10.12. *There is some* $\psi \in End(\Lambda_{\mathbb{N}})^\times$, *such that* $D^bCoh(Z_q^*) \cong D^\pi(\psi^*\mathcal{Q}_{64,q} \otimes_{\Lambda_{\mathbb{N}}} \Lambda_{\mathbb{Q}})$. □

11. Counting polygons

We begin by showing that for a quartic surface in $\mathbb{C}\mathrm{P}^3$ whose part at infinity is the intersection with the "simplex" $x_0x_1x_2x_3 = 0$, the deformation from affine to relative Fukaya category is nontrivial. This is based on Corollary 8.12, but the construction of the relevant Lagrangian torus involves geometric ideas related to the SYZ conjecture [53], in particular the appearance of amoebas in that context [52, 43]. Having done that, the next step is to apply Corollary 9.6 to prove that the 64 vanishing cycles of the Fermat pencil of quartics are split-generators for the derived Fukaya category. This reduces us to studying the full A_∞-subcategory formed by these cycles, and finally, the dimensional induction machinery from Section 9e allows us to compute that subcategory combinatorially.

(11a) Let $X = \mathbb{C}\mathrm{P}^3$ with $o_X = \mathcal{K}_X^{-1} = \mathcal{O}(4)$, and $\sigma_{X,\infty} = x_0x_1x_2x_3$. Choose some section $\sigma_{X,0}$ which with $\sigma_{X,\infty}$ generates a quasi-Lefschetz pencil, and such that X_0 is regular.

Lemma 11.1. *$\mathcal{F}(X_0, X_{0,\infty})$, as a one-parameter deformation of A_∞-categories, is nontrivial at first order in q.*

The validity of the statement is independent of the particular choice of $\sigma_{X,0}$. Any two different choices $\sigma_{X,0}^0$, $\sigma_{X,0}^1$ can be joined by a smooth family $\sigma_{X,0}^t$, of sections of o_X, such that for all t,

$$X_0^t = (\sigma_{X,0}^t)^{-1}(0)$$

is a smooth quartic and intersects each stratum of X_∞ transversally. For the same reasons as in the proof of Proposition 7.22, there is a family of symplectic isomorphisms $\phi^t : X_0^0 \to X_0^t$ which maps $X_0^0 \cap X_\infty$ to $X_0^t \cap X_\infty$. Since quartic surfaces are simply-connected, one sees from Remark 8.5 that ϕ^1 defines an equivalence of relative Fukaya categories.

It remains to prove Lemma 11.1 in a single case. Let $C = \{x_3 = 0\} \cong \mathbb{C}\mathrm{P}^2$ be one of the irreducible components of X_∞. The intersection with the other components is $\Delta = C \cap C' \cong \{x_0x_1x_2 = 0\} \subset \mathbb{C}\mathrm{P}^2$. By construction, $\mathbb{C}\mathrm{P}^2$ carries the Fubini-Study symplectic form ω_C, rescaled in such a way that a line has area 4. We also have the meromorphic complex volume form η_C with poles along Δ. Take (this will not quite be our final word) a section of the form

$$\sigma_{X,0} = x_0^4 + x_1^4 + \tfrac{1}{4}x_2^4 + 2x_0x_1x_2^2 + (\text{monomials containing } x_3). \tag{11.1}$$

$X_0 \cap C$ is the smooth quartic curve $Q = \{x_0^4 + x_1^4 + \frac{1}{4}x_2^4 + 2x_0x_1x_2^2 = 0\} \subset \mathbb{C}\mathrm{P}^2$, which intersects all the strata of Δ transversally. Hence, by suitably choosing the terms not spelled out in (11.1), one can achieve that $\sigma_{X,0}$ generates a quasi-Lefschetz pencil. Consider the family of Lagrangian torus orbits

$$T_\lambda = \{\lambda_2|x_0|^2 = \lambda_0|x_2|^2,\ \lambda_2|x_1|^2 = \lambda_1|x_2|^2\} \subset \mathbb{C}\mathrm{P}^2 \setminus \Delta,$$

for $\lambda = (\lambda_0, \lambda_1, \lambda_2)$ with $\lambda_k > 0$ and $\sum_k \lambda_k = 1$. We take

$$L_0 = T_{\frac{1}{4},\frac{1}{4},\frac{1}{2}},$$

which by an elementary estimate is disjoint from Q. The next question is how far this satisfies Assumptions 8.15. Each T_λ is graded with respect to η_C, and $O(4)|T_\lambda$ is the trivial flat line bundle iff $\lambda_k \in \frac{1}{4}\mathbb{Z}$ for all k, which is the case for L_0. Look at the holomorphic discs $w : (D, \partial D) \to (C, L_0)$ given by

$$w(z) = \begin{cases} [\zeta : z : \sqrt{2}], \\ [z : \zeta : \sqrt{2}] \end{cases} \quad \text{for some } \zeta \in \mathbb{C} \text{ with } \zeta^4 = -1. \tag{11.2}$$

They satisfy $\int w^*\omega_C = w \cdot Q = 1$, and their boundary circles generate $H_1(L_0)$. By Lemma 6.1, this implies that L_0 is exact with respect to θ_C.

Lemma 11.2. *Every holomorphic disc $w : (D, \partial D) \longrightarrow (\mathbb{C}\mathrm{P}^2, T_\lambda)$ is regular.*

Proof. Take the Euler sequence on $\mathbb{C}\mathrm{P}^2$,

$$0 \longrightarrow \mathcal{O} \longrightarrow \mathcal{O}(1) \otimes H^0(\mathcal{O}(1))^\vee \longrightarrow T\mathbb{C}\mathrm{P}^2 \longrightarrow 0. \tag{11.3}$$

Choose a basis for $H^0(\mathcal{O}(1))^\vee$ dual to the three coordinate functions x_0, x_1, x_2. Then the first map is just given by these functions, and the second map takes $i(x_k \otimes x_k^\vee)$ to the vector field which infinitesimally rotates the k-th coordinate. It follows that over T_λ, we have a sequence of real subbundles of (11.3),

$$0 \longrightarrow i\mathbb{R} \longrightarrow i\mathbb{R}x_0 \oplus i\mathbb{R}x_1 \oplus i\mathbb{R}x_2 \longrightarrow T(T_\lambda) \longrightarrow 0.$$

After pulling this back to the disc via w and introducing the associated Cauchy-Riemann operators, one finds that there is a surjection

$$\mathrm{coker}(D_0 \oplus D_1 \oplus D_2) \twoheadrightarrow \mathrm{coker}\, D_{w,\mathbb{C}\mathrm{P}^2},$$

where D_k is the $\bar\partial$-operator on $w^*\mathcal{O}(1)$ with totally real boundary conditions $w^*(i\mathbb{R}x_k)$. The Maslov number of $(w^*\mathcal{O}(1), w^*(i\mathbb{R}x_k))$ is twice the intersection number of w with the line $\{x_k = 0\}$, hence nonnegative, which means that D_k has trivial cokernel. □

Lemma 11.3. *Up to the* $PSL(2,\mathbb{R})$*-action on* D*, any holomorphic disc* $w : (D, \partial D) \rightarrow (\mathbb{C}\mathrm{P}^2, T_\lambda)$ *with Maslov number 2 is either of the form*

$$w(z) = [\lambda_0 : \zeta\lambda_1 : \lambda_2 z] \quad or \quad w(z) = [\lambda_0 : \zeta\lambda_1 : \lambda_2 z^{-1}]$$

for some $|\zeta| \in S^1$*, or else the same with the roles of the coordinates exchanged.*

Proof. The virtual dimension of the moduli space of unparametrized discs is equal to one. By Lemma 11.2, this moduli space is indeed a smooth one-dimensional manifold. Since on the other hand, $T^2 \subset PU(3)$ acts on it, we find that there must be a sub-circle of T^2 which leaves w invariant as an unparametrized disc. In other words, after applying an initial $PSL(2,\mathbb{R})$-action one can assume that there are weights $d_0, d_1, d_2 \in \mathbb{Z}^3$, not all of them equal, such that

$$w(e^{it}z) = \mathrm{diag}(e^{itd_0}, e^{itd_1}, e^{itd_2})\, w(z).$$

By holomorphicity $w(z) = \mathrm{diag}(z^{d_0}, z^{d_1}, z^{d_2})\, w(1)$. However, at $z = 0$ our disc may intersect Δ only with multiplicity one, and that implies $d = (0, 0, \pm 1)$ or else the same with the coordinates permuted. □

Lemma 11.4. *Up to the action of* S^1 *on* D *by rotations, there are precisely eight holomorphic discs* $w : (D, \partial D) \rightarrow (\mathbb{C}\mathrm{P}^2, L_0)$ *with* $w \cdot \Delta = w \cdot Q = 1$ *and* $w(0) \in \Delta \cap Q$*. Moreover, for each of them* $[w|\partial D] \in H_1(L_0)$ *is nontrivial, and the restricted linearized operator* $D\bar{\partial}^{res}_{w,\mathbb{C}\mathrm{P}^2}$ *as in (8.18) has one-dimensional cokernel.*

Proof. Because L_0 is graded with respect to η_C, $w \cdot \Delta = 1$ implies that the Maslov number is 2, so we can apply Lemma 11.3. Out of the possible six choices of weights, only $d = (1, 0, 0)$ or $(0, 1, 0)$ lead to discs with $\int w^*\omega_C = w \cdot Q = 1$. The condition that $w(0) \in \Delta \cap Q$ further restricts us to the eight discs from (11.2). The statement about $[w|\partial D]$ is obvious, and that about the cokernel is a straightforward computation. □

Now perturb the equation defining Q slightly, in such a way that out of the eight points of $Q \cap \{x_0 x_1 = 0\}$, one remains fixed while the others move in a generic way. Then if we replace Q by its perturbed version Q' in Lemma 11.4, only one of the 8 discs persists, which we call w_0. Take a corresponding slight perturbation $\sigma'_{X,0}$ of $\sigma_{X,0}$ which still generates a quasi-Lefschetz pencil. Then Assumptions 8.15 are satisfied: we already verified those concerning L_0 and w_0, and as for those excluding other pseudo-holomorphic maps, part (i) comes out of Lemma 11.4; (ii) follows from Lemma 11.2 because a disc w which does not intersect Δ would have zero Maslov number, hence negative virtual dimension; and (iii) is obvious, since the ω_C-area of any $v : S^2 \rightarrow \mathbb{C}\mathrm{P}^2$ is a multiple of 4. An appeal to Corollary 8.21 completes the proof of Lemma 11.1.

(11b) As before take $X = \mathbb{C}\mathrm{P}^3$ with $\sigma_{X,\infty}(x) = x_0 x_1 x_2 x_3$, so that the complement of X_∞ can be written as a quotient by the action of $\Gamma_4^* = \{\pm I, \pm iI\} \subset$

$SL_4(\mathbb{C})$,

$$M = \{x \in \mathbb{C}^4 \;:\; x_0x_1x_2x_3 = 1\}/\Gamma_4^*. \tag{11.4}$$

We now specify X_0 to be the Fermat quartic $\sigma_{X,0} = \frac{1}{4}(x_0^4 + x_1^4 + x_2^4 + x_3^4)$. X_0 intersects the strata of X_∞ transversally, and $\pi_M = \sigma_{X,0}/\sigma_{X,\infty}$ has 64 nondegenerate critical points, coming in groups of 16 which lie in the same fibre (this is not a problem, see Remark 6.4). We take $z_* = 0$ as our base point, and use the same embedded vanishing path for all critical points which lie in a given fibre. This yields a collection of 64 vanishing cycles in $M_0 = X_0 \setminus X_{0,\infty}$, divided into four groups of 16 mutually disjoint ones. We make them into branes by choosing gradings and, for the projective and relative versions of that notion, the other additional data. Notation: $\mathcal{F}_{64}$ will be the full A_∞-subcategory of $\mathcal{F}(M_0)$ whose objects are these vanishing cycles; and $\mathcal{F}_{64,q}$, $\mathcal{F}_{64,q}^*$ the corresponding A_∞-subcategories of $\mathcal{F}(X_0)$, $\mathcal{F}(X_0, X_{0,\infty})$. From Propositions 8.6 and 8.7 we know that $\mathcal{F}_{64,q}$ is a one-parameter deformation of $\mathcal{F}_{64}$, and $\mathcal{F}_{64,q}^* \cong \mathcal{F}_{64,q} \otimes_{\Lambda_\mathbb{N}} \Lambda_\mathbb{Q}$.

Lemma 11.5. $D^\pi\mathcal{F}(X_0) \cong D^\pi(\mathcal{F}_{64,q}^*)$.

Proof. Corollary 9.6 shows that the vanishing cycles are split-generators for $D^\pi\mathcal{F}(X_0)$, and Lemma 2.4 yields the desired conclusion. □

Lemma 11.6. *$\mathcal{F}_{64,q}$, seen as a one-parameter deformation of $\mathcal{F}_{64}$, is nontrivial at first order.*

Proof. We know from Lemma 11.1 that nontriviality is true for the whole of $\mathcal{F}(X_0, X_{0,\infty})$. A closer look at the argument, and in particular at the proof of Corollary 8.12, reveals that we have in fact established the assumptions of Lemma 3.12, which shows that the deformation remains nontrivial when restricted to any subcategory which has enough objects to split-generate the whole. □

To simplify the geometry of our pencil, consider the action of the group $\Gamma_{16}^* \subset PSL_4(\mathbb{C})$ of diagonal matrices A satisfying $A^4 = I$ on X. This action is free on M; it can be lifted to o_X and preserves the Fubini-Study connection, as well as $\sigma_{X,0}, \sigma_{X,\infty}$; and it is compatible with the isomorphism $o_X \cong \mathcal{K}_X^{-1}$. Hence, on the quotient $\bar{M} = M/\Gamma_{16}^*$ we have an induced symplectic form, a one-form primitive for it, a holomorphic volume form, and a function $\pi_{\bar{M}} : \bar{M} \to \mathbb{C}$. Explicitly,

$$\bar{M} = \{u_0u_1u_2u_3 = 1\}, \quad \pi_{\bar{M}}(u) = \tfrac{1}{4}(u_0 + u_1 + u_2 + u_3)$$

where the coordinates are related to the previous ones by $u_k = x_k^4$. Since Γ_{16}^* permutes the critical points in any given fibre transitively, $\pi_{\bar{M}}$ has one nondegenerate critical point in each each of the four singular fibres. In other words, the 64 vanishing cycles in M_0 are all possible lifts of the corresponding 4 vanishing cycles in the quotient $\bar{M}_0$. We need some basic information about the topology of this space:

$$\begin{aligned} &\pi_1(\bar{M}_0) = \{h \in \mathbb{Z}^4 \ : \ h_0 + \dots + h_3 = 0\}, \\ &im(\pi_2(\bar{M}_0) \otimes \mathbb{Q} \rightarrow H_2(\bar{M}_0; \mathbb{Q})) \cong \mathbb{Q}^3 \end{aligned} \tag{11.5}$$

where the first group maps isomorphically to $\pi_1(\bar{M})$, with the generators h_k becoming loops around the coordinates axes $u_k = 0$ at infinity; and the second is generated by the 4 vanishing cycles. Both things follow from elementary Morse and Lefschetz theory, since $\bar{M} \simeq T^3$ is obtained by attaching four 3-cells to $\bar{M}_0$, which is itself homotopy equivalent to a 2-complex. As explained in Section 8b, this means that we have an action of $\pi_1(\bar{M}_0)^*$ on a subcategory of $\mathcal{F}(\bar{M}_0)$ which consists of branes with a certain condition on their fundamental groups, and in particular contains our vanishing cycles since they are spheres. Recall that in Section 10a we considered the group $T = H/\Gamma_4 \subset PSL(V)$. If one chooses a basis of V making the maximal torus $H \subset SL(V)$ standard, T can be written as $(\mathbb{C}^*)^4/\mathbb{C}^*$ (the quotient of the group of all diagonal matrices by the multiples of the identity) and hence can be identified with $\pi_1(\bar{M}_0)^*$. The following is our main computational result, and will be proved later:

Proposition 11.7. *For a suitable choice of the lifts to M_0 which define the T-action, full A_∞-subcategory $\mathcal{F}_4 \subset \mathcal{F}(\bar{M}_0)$ whose objects are the 4 vanishing cycles of $\pi_{\bar{M}}$ is T-equivariantly quasi-isomorphic to $\mathcal{Q}_4$.*

The covering $M_0 \rightarrow \bar{M}_0$ provides an epimorphism $\pi_1(\bar{M}_0) \rightarrow \Gamma_{16}^*$, and the dual of that is the inclusion of the subgroup $\Gamma_{16} \hookrightarrow T$ considered in Section 10a. Combining the statement above with Lemma 8.2, one finds that

$$\mathcal{F}_{64} \cong \mathcal{F}_4 \rtimes \Gamma_{16} \cong \mathcal{Q}_4 \rtimes \Gamma_{16} \cong \mathcal{Q}_{64}.$$

Since each morphism space $u_i Q_{64} u_j$ is concentrated in a single degree, the one-parameter deformation $\mathcal{F}_{64,q}$ necessarily satisfies $H^*(\mathcal{F}_{64,q}) \cong H^*(\mathcal{F}_{64}) \otimes_{\mathbb{C}} \Lambda_{\mathbb{N}}$, so we can use the Perturbation Lemma to find a quasi-isomorphic deformation of $\mathcal{Q}_{64}$. To complete the argument, consider the symplectic automorphism $\bar{\phi}$ of $\bar{M}$ given by $\bar{\phi}(u_0, u_1, u_2, u_3) = (u_1, u_2, u_3, u_0)$. This acts fibrewise with respect to $\pi_{\bar{M}}$, hence takes each vanishing cycle to itself, but it reverses their orientation since it acts with determinant -1 on the tangent space to each critical point. If we equip $\bar{\phi}_0 = \bar{\phi}|\bar{M}_0$ with a suitable "odd" grading $\tilde{\alpha}_{\bar{\phi}_0}$, it will map each vanishing cycle to itself as a graded Lagrangian submanifold. Since it preserves the complex structure, it is no problem to make $\bar{\phi}_0$ act on the Fukaya category, yielding an A_∞-homomorphism $W_4 : \mathcal{F}_4 \rightarrow \mathcal{F}_4$ which acts as -1 on $HF^2(V, V)$ for each of the four vanishing cycles. This also skew-commutes with the T-action up to an automorphism ξ of T, which is simply the dual of the action of $\bar{\phi}$ on $\pi_1(\bar{M})$, hence which permutes the coordinates cyclically. Under the isomorphism $H(\mathcal{F}_4) \cong Q_4$, we can therefore identify $H(W_4)$ with one of the automorphisms U_4 considered in Section 10c. $\bar{\phi}_0$ lifts to an automorphism of M_0, $\phi_0(x_0, \dots, x_3) = (x_1, x_2, x_3, x_0)$, and this extends to the compactification X_0. This means that $W_{64} = W_4 \rtimes (\xi|\Gamma_{16})$ which can be identified with the action of ϕ_0 on $\mathcal{F}_{64}$, admits a deformation $W_{64,q}$ to an A_∞-homomorphism from $\mathcal{F}_{64,q}$ to itself. Together with Lemma 11.6, this means that we have verified all the conditions of Lemma 10.4, and therefore can conclude that $\mathcal{F}_{64,q}$ is quasi-isomorphic to $\mathcal{Q}_{64,q}$ up to some change of parameter in $End(\Lambda_\mathbb{N})^\times$. Using Lemma 11.5, we obtain:

Corollary 11.8. *There is a* $\psi \in End(\Lambda_\mathbb{N})^\times$ *such that*

$$D^\pi \mathcal{F}(X_0) \cong D^\pi(\mathcal{F}^*_{64,q}) \cong \psi^* D^\pi(\mathcal{Q}_{64,q} \otimes_{\Lambda_\mathbb{N}} \Lambda_\mathbb{Q}). \quad \square$$

This is the mirror statement to Corollary 10.12, and by comparing the two, Theorem 1.2 follows.

(11c) It remains to explain Proposition 11.7. Following the strategy from Section 6f, the first step in understanding the geometry of the vanishing cycles is to represent them as matching cycles, which involves the choice of a generic auxiliary section. Let us momentarily work "upstairs" in M. An ansatz for the section is

$$\sigma'_{X,0} = \tfrac{1}{4}(\xi_0 x_0^4 + \xi_1 x_1^4 + \xi_2 x_2^4 + \xi_3 x_3^4) \tag{11.6}$$

with pairwise distinct constants $\xi_k \in \mathbb{C}^*$. This always satisfies the first of the genericity assumptions, namely $(\sigma'_{X,0})^{-1}(0)$ intersects each stratum of $X_{0,\infty}$ transversally. The critical point set of the associated map (6.10) is always smooth,

$$Crit(b_M) = \Big\{ x_k^4 = (s + \xi_k t)^{-1}, \text{ where } \prod_k (s + t\xi_k) = 1 \Big\}.$$

The parameter value $t = 0$ corresponds precisely to the 64 critical points of π_M. At those points, $b_M : Crit(b_M) \to \mathbb{C}^2$ is an immersion, and its second component $\sigma'_{X,0}/\sigma_{X,\infty} : Crit(b_M) \to \mathbb{C}$ a simple branched cover, iff the coefficients ξ_k satisfy

$$\sum_k \xi_k^2 \neq \tfrac{1}{4}\Big(\sum_k \xi_k\Big)^2. \tag{11.7}$$

Next, $b_M|Crit(b_M)$ cannot be a generic embedding because it is Γ^*_{16}-invariant, but after passing to the quotient we do get a generic embedding $b_{\bar{M}} : Crit(b_{\bar{M}}) \to \mathbb{C}^2$, for elementary reasons: the image $\bar{C}$ of this map is an irreducible algebraic curve in $\mathbb{C}^2$ having four distinct points at infinity, and since $Crit(b_{\bar{M}})$ has Euler characteristic -2, the map cannot be a multiple branch cover. The upshot is that the dimensional induction technique from Section 9d can be applied "downstairs", meaning to the vanishing cycles of $\pi_{\bar{M}}$.

Let's get down to concrete numbers. Our choice of weights is $\xi_0 = 5/4$, $\xi_1 = i$, $\xi_2 = -1$, $\xi_3 = -i$, which satisfies (11.7). The branch curve (determined by computer using elimination theory) is

$$
\begin{aligned}
\bar{C} = \{&128000z^{12} - 76800z^{11}w + 92160z^{10}w^2 - 108544z^9w^3 - 258048z^8w^4 \\
&+ 86016z^7w^5 - 112640z^6w^6 + 141312z^5w^7 + 211968z^4w^8 - 25600z^3w^9 \\
&+ 36864z^2w^{10} - 49152zw^{11} - 65536w^{12} - 391307z^8 + 180632z^7w \\
&+ 1940z^6w^2 - 503112z^5w^3 - 2102866z^4w^4 + 457512z^3w^5 - 122764z^2w^6 \\
&- 102904zw^7 - 251483w^8 + 399430z^4 - 106312z^3w - 155028z^2w^2 \\
&- 53752zw^3 - 320858w^4 - 136161 = 0\}.
\end{aligned}
$$

Figure 4 shows the z-plane $\mathbb{C}$ with the 4 points of $Critv(\pi_{\bar{M}}) = \{\pm 1, \pm i\}$, as well the 40 points of $Fakev(\pi_{\bar{M}})$. Luckily, none of the latter lie on $[-1;1]$ or $i[-1;1]$, so we can take the basis of embedded critical paths which are straight lines $b_k(t) = i^{2-k}t$, $k = 1, \dots, 4$. Denote by $V_{b_k} \subset \bar{M}_0$ the resulting vanishing cycles. We now turn to the function

$$
q_{\bar{M}_0} = \tfrac{1}{4} \sum_k \xi_k u_k : \bar{M}_0 = \pi_{\bar{M}}^{-1}(0) \longrightarrow \mathbb{C}, \tag{11.8}
$$

which is also a quasi-Lefschetz fibration (or more accurately, the Γ_{16}^*-quotient of such a fibration on M_0). Figure 5 shows its basis $\mathbb{C}$ with the 12 critical values, which are also the points of $\bar{C} \cap \{z = 0\}$, and the matching paths d_k obtained from the braid monodromy of $\bar{C}$ over b_k. The associated matching cycles are our vanishing cycles: $V_{b_k} \simeq \Sigma_{d_k}$.

Taking 0 as a base point, choose a base path c_* and a basis of vanishing paths $c_1, \dots, c_{12}$ for (11.8) as indicated in Figure 6. This gives rise to vanishing cycles $V_{c_k} \subset \bar{M}_{0,0} = q_{\bar{M}_0}^{-1}(0)$, which are simple closed loops on a four-pointed genus three Riemann surface.

Remark 11.9. *If one took $\xi_k = i^k$ instead, condition* (11.7) *would be violated but the structure of $q_{\bar{M}_0}$ would be much simpler: it would have 4 degenerate critical points of type A_3. The actually chosen ξ_k are a small perturbation of these values, in the sense that one can still recognize the groups of 3 critical points obtained by "morsifying" the degenerate singularities. This allows one to choose the vanishing paths c_k in such a way as to minimize the intersections between the vanishing cycles in each group.*

Let $\mathcal{F}_{\overrightarrow{12}}$ be the directed A_∞-subcategory of the Fukaya category of $\bar{M}_{0,0}$ whose objects are the V_{c_k}. From the restriction of the universal cover of $\bar{M}_0$, we get an action of $\pi_1(\bar{M}_0)^* \cong T$ on this A_∞-category. The equivariant version of Theorem 9.7 shows that $\mathcal{F}_4$ is equivariantly quasi-isomorphic to the full A_∞-subcategory of $Tw_T(\mathcal{F}_{\overrightarrow{12}})$ whose objects are the following four equivariant twisted complexes, obtained by applying the rules from Section 9d to the paths from Figures 5, 6:

$$\begin{aligned} S_1 &= \bigl\{T'_{V_{c_7}}T'_{V_{c_5}}(V_{c_3}) \longrightarrow T_{V_{c_{10}}}T_{V_{c_{11}}}(V_{c_{12}})\bigr\}, \\ S_2 &= \bigl\{T_{V_{c_1}}T_{V_{c_2}}(V_{c_3}) \longrightarrow T'_{V_{c_{10}}}T'_{V_{c_8}}(V_{c_6})\bigr\}, \\ S_3 &= \bigl\{T_{V_{c_1}}T_{V_{c_4}}T_{V_{c_5}}(V_{c_6}) \longrightarrow T'_{V_{c_{11}}}(V_{c_9})\bigr\}, \\ S_4 &= \bigl\{T_{V_{c_2}}T_{V_{c_4}}T_{V_{c_7}}T_{V_{c_8}}(V_{c_9}) \longrightarrow V_{c_{12}}\bigr\} \end{aligned} \tag{11.9}$$

where $\{\cdots \to \cdots\}$ is the cone over the lowest degree nonzero morphism, which is unique up to a $\mathbb{C}^*$ multiple. Actually, that morphism may not have degree zero, or it may not be T-invariant, so one needs to change one of the two objects forming the cone by a translation and tensoring with a suitable T-character. In that sense, the formulae given (11.9) are not entirely precise, but the appropriate corrections can not yet be determined at this stage in the computation.

The next step is to determine $\mathcal{F}_{\overrightarrow{12}}$, or rather a directed A_∞-category $\mathcal{C}_{\overrightarrow{12}}$ quasi-isomorphic to it, using the combinatorial description of directed Fukaya categories on Riemann surfaces. Following the procedure described in Section 9e, we take a compact piece $N \subset \bar{M}_{0,0}$ which is a surface with boundary and a deformation retract. Figure 7 shows a decomposition of N into eight polygons (I)–(VIII), with the solid lines the actual ∂N, and the dashed lines to be glued to the boundaries of other polygons as indicated. $\bar{M}_{0,0}$ inherits the structure of an affine Calabi-Yau from $\bar{M}$, and Figure 8 shows an unoriented foliation on N lying in the same homotopy class as the distinguished trivialization of the squared canonical bundle. Moreover, restriction

$$\mathbb{Z}^4/\mathbb{Z}(1,1,1,1) \cong H^1(\bar{M}) \longrightarrow H^1(\bar{M}_{0,0})$$

gives four canonical elements of $H^1(N)$ whose sum is zero; the Poincaré duals of the first three elements are represented by the one-cycles x, y, z in $(N, \partial N)$ shown in Figure 9. Figures 10–17 show simple closed curves ν_k isotopic to the vanishing cycles V_{c_k}, drawn on each of the eight polygons which make up N.

Remark 11.10. *These pictures were arrived at by an application of braid monodromy (in one dimension lower than before), which means that we introduce another auxiliary holomorphic function on $\bar{M}_0$, whose restriction to any generic fibre of $q_{\bar{M}_0}$ represents that surface as a branched cover with only simple branch points. The function we use is simply a projection*

$$r_{\bar{M}_0}(u) = u_2 \in \mathbb{C}^*,$$

making $\bar{M}_{0,0}$ into a fourfold branched cover of $\mathbb{C}^*$ with eight simple branch points. Cutting the base of that into suitable pieces gives rise to our decomposition of N into polygons. Elimination theory is used to compute the branch curve of

$$(q_{\bar{M}_0}, r_{\bar{M}_0}) : \bar{M}_{0,0} \longrightarrow \mathbb{C} \times \mathbb{C}^*,$$

and from that one obtains the vanishing cycles $V_{c_1}, \dots, V_{c_{12}}$. We have omitted the details of this step, since it is essentially topological, and completely parallel to the procedure which led to the paths $b_1, \dots, b_4$ in Figure 5.

When drawing the ν_k, there is some freedom in their relative position with respect to each other. The only constraint, which comes from exactness, says that we should be able to assign positive areas to the connected components of $N \setminus \bigcup_k \nu_k$ such that the two-chains which represent relations between the ν_k have signed area zero. In fact, as explained in Remark 9.8, not all relations need to be considered; in our case (11.5) shows that there are three essential ones. These can be read off from (11.9) and the intersection numbers of the ν_k, and are

$$\begin{aligned} &\nu_5 + \nu_8 + \nu_{10} - \nu_1 - \nu_2 - \nu_3 \sim 0, \\ &\nu_3 + \nu_5 + \nu_7 - \nu_{10} - \nu_{11} - \nu_{12} \sim 0, \\ &\nu_8 + \nu_2 + \nu_4 - \nu_7 - \nu_8 - \nu_0 \sim 0. \end{aligned} \tag{11.10}$$

After having drawn the relevant two-chains explicitly in Figures 10–17, one realizes that for each of them, there are pieces of $N \setminus \bigcup_k \nu_k$ which appear in it with either sign, and which do not appear in the other two-chains. This means that the configuration of curves ν_k can indeed be used as basis for the computation.

Choose gradings and *Spin* structures for the ν_k, lifts to the abelian covering of N induced from $\pi_1(N) \cong \pi_1(\bar{M}_{0,0}) \rightarrow \pi_1(\bar{M}_0)$, and identifications $\mathbb{C}_x \cong \mathbb{C}$ for each intersection point of $\nu_k \cap \nu_l$ with $k < l$. One can then list all intersection points with their (Maslov index) degrees as well as their (T-action) weights. This is done in Tables 1 and 2, where the notation is that

$$m^d x^a y^b z^c \tag{11.11}$$

is an intersection point with Maslov index d, and on which $[x, y, z, 1] \in T$ acts as $x^a y^b z^c$, respectively. By inspecting the table, one sees that different intersection points of the same $\nu_k \cap \nu_l$ are always distinguished by (a, b, c, d). We may therefore denote the generator of $hom_{\mathcal{C}_{\overrightarrow{12}}}(\nu_k, \nu_l)$ given by an intersection point (11.11) by

$$\nu_k \cap \nu_l : m^d x^a y^b z^c.$$

The next step is to count all immersed holomorphic polygons which have sides on the ν_k (in the correct order), with the appropriate signs. The outcome of that are the composition maps μ^d of $\mathcal{C}_{\overrightarrow{12}}$. It turns out that only μ^2, μ^3 are nonzero; their coefficients are listed in Tables 3 and 4.

Remark 11.11. *The T-action is actually a big bonus in this computation, since the fact that the μ^d have to be homogeneous means that a lot of their coefficients have to vanish. For instance, the fact that $\mu^d = 0$ for $d > 3$ follows by inspection of the list of T-weights of the intersection points, so one does not need to verify geometrically that there are no immersed pentagons, hexagons, and so forth.*

At this point, we can write down the objects S_k from (11.9), or rather the corresponding objects C_k in $Tw_T(\mathcal{C}_{\overrightarrow{12}})$, more explicitly. They are

$$C_1 = \nu_3[1]\langle x\rangle \oplus \nu_5\langle y\rangle \oplus \nu_7\langle yz\rangle \oplus \nu_{10}\langle x^{-1}y\rangle \oplus \nu_{11}[-1]\langle y\rangle \oplus \nu_{12}[-1],$$

$$\delta_1 = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ \nu_3 \cap \nu_5 : x^{-1}y & 0 & 0 & 0 & 0 & 0 \\ 0 & \nu_5 \cap \nu_7 : mz & 0 & 0 & 0 & 0 \\ 0 & \nu_5 \cap \nu_{10} : mx^{-1} & 0 & 0 & 0 & 0 \\ 0 & 0 & \nu_7 \cap \nu_{11} : z^{-1} & \nu_{10} \cap \nu_{11} : x & 0 & 0 \\ 0 & 0 & 0 & 0 & \nu_{11} \cap \nu_{12} : my^{-1} & 0 \end{pmatrix},$$

$$C_2 = \nu_1[1]\langle x^{-1}y\rangle \oplus \nu_2\langle y\rangle \oplus \nu_3 \oplus \nu_6\langle z^{-1}\rangle \oplus \nu_8[-1]\langle z^{-1}\rangle \oplus \nu_{10}[-1]\langle x^{-1}\rangle,$$

$$\delta_2 = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ \nu_1 \cap \nu_2 : x & 0 & 0 & 0 & 0 & 0 \\ 0 & \nu_2 \cap \nu_3 : my^{-1} & 0 & 0 & 0 & 0 \\ 0 & \nu_2 \cap \nu_6 : my^{-1}z^{-1} & 0 & 0 & 0 & 0 \\ 0 & 0 & -(\nu_3 \cap \nu_8 : z^{-1}) & \nu_6 \cap \nu_8 : 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & \nu_8 \cap \nu_{10} : mx^{-1}z & 0 \end{pmatrix},$$

$$\begin{aligned} C_3 = {} & \nu_1\langle x^{-1}\rangle \oplus \nu_4\langle x^{-1}y\rangle \oplus \nu_5[-1]\langle x^{-1}y\rangle \oplus \nu_6[-1]\langle x^{-1}z^{-1}\rangle \\ & \oplus \nu_9[-1]\langle y^{-1}z^{-1}\rangle \oplus \nu_{11}[-2]\langle z^{-1}\rangle, \end{aligned}$$

$$\delta_3 = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ \nu_1 \cap \nu_5 : y & \nu_4 \cap \nu_5 : 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & \nu_5 \cap \nu_6 : my^{-1}z^{-1} & 0 & 0 & 0 \\ 0 & 0 & \nu_5 \cap \nu_9 : mxy^{-2}z^{-1} & 0 & 0 & 0 \\ 0 & 0 & 0 & \nu_6 \cap \nu_{11} : x & \nu_9 \cap \nu_{11} : y & 0 \end{pmatrix},$$

$$\begin{aligned} C_4 = {} & \nu_2[1]\langle z^{-1}\rangle \oplus \nu_4[1]\langle x^{-1}\rangle \oplus \nu_7[1]\langle x^{-1}y\rangle \oplus \nu_8\langle x^{-1}z^{-1}\rangle \\ & \oplus \nu_9\langle x^{-1}y^{-1}z^{-1}\rangle \oplus \nu_{12}\langle y^{-1}z^{-1}\rangle, \end{aligned}$$

$$\delta_4 = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ \nu_2 \cap \nu_4 : mx^{-1}z & 0 & 0 & 0 & 0 & 0 \\ \nu_2 \cap \nu_7 : mx^{-1}yz & 0 & 0 & 0 & 0 & 0 \\ 0 & \nu_4 \cap \nu_8 : z^{-1} & \nu_7 \cap \nu_8 : y^{-1}z^{-1} & 0 & 0 & 0 \\ 0 & 0 & 0 & \nu_8 \cap \nu_9 : my^{-1} & 0 & 0 \\ 0 & 0 & 0 & \nu_8 \cap \nu_{12} : mxy^{-1} & 0 & 0 \end{pmatrix}.$$

where $\nu_k[d]\langle x^a y^b z^c\rangle$ means the object ν_k shifted down by d, and tensored with the one-dimensional representation of T on which $[x, y, z, 1]$ acts as $x^{-a}y^{-b}z^{-c}$.

It is now easy to compute explicitly the full A_∞-subcategory of $Tw_T\mathcal{C}_{\overrightarrow{12}}$ consisting of these four equivariant twisted complexes C_k. More precisely, one verifies (by computer) first that the cohomology of this subcategory is T-equivariantly isomorphic to Q_4 and then, following the indications in Remark 10.2, that a suitable fourth order Massey product is nonzero, which implies that our full subcategory is T-equivariantly quasi-isomorphic to $\mathcal{Q}_4$. By Theorem 9.7, or rather the equivariant version of it explained in Section 9f, we have a T-equivariant quasi-isomorphism of our subcategory with $\mathcal{F}_4$, and this completes the proof of Proposition 11.7.



FIGURE 4. The actual (•) and fake (∘) critical points of $\pi_{\bar{M}}$, together with our choice of vanishing paths $b_1, \dots, b_4$. The image has been distorted for better legibility.



FIGURE 5. The four matching paths obtained from the vanishing paths in Figure 4 by braid monodromy. These were drawn using a computer to watch how the points of $\bar{C} \cap \{z = b_k(t)\}$ move in $\mathbb{C}$ as one goes along the vanishing paths. The picture would be $\mathbb{Z}/4$-symmetric if we hadn't moved the paths by an isotopy, so as to avoid the future base path (the dashed line).



FIGURE 6. The base path and the 12 vanishing paths $c_1, \dots, c_{12}$ (ordered as they must be, clockwise with respect to the derivatives at the base point).


VII
II
I
III
VIII
VIII
III
II
IV
I
I
IV
III
V
II
II
V
IV
VI
III
III
VI
V
VII
IV
IV
VII
VI
VIII
V
V
VIII
VII
I
VI
VI
I
VIII
II
VII

Figure 7.



Figure 8.



Figure 9.



Figure 10.



Figure 11.



Figure 12.



Figure 13.



FIGURE 14.



FIGURE 15.



Figure 16.



Figure 17.

	1	2	3	4	5	6
1						
2	x					
3		my^{-1}				
4	1 x	$mx^{-1}z$				
5	x 1 y	$x^{-1}y$ 1	$x^{-1}y$	1		
6		$my^{-1}z^{-1}$ $mx^{-1}z^{-1}$ $mx^{-1}y^{-1}$	$x^{-1}yz^{-1}$ x^{-1}		$my^{-1}z^{-1}$	
7	myz x	$mx^{-1}z$ $mx^{-1}yz$ mz		1 x	mz	
8	xz^{-1} $xy^{-1}z^{-1}$ z^{-1}	z^{-1} $mx^{-1}y^{-1}$	x^{-1} z^{-1} $x^{-1}yz^{-1}$	$xy^{-1}z^{-1}$ $y^{-1}z^{-1}$ z^{-1}	z^{-1} $xy^{-1}z^{-1}$	1
9		$my^{-1}z^{-1}$ $mx^{-1}y^{-1}$ my^{-2}	x^{-1} $my^{-1}z^{-1}$		$mxy^{-2}z^{-1}$ $my^{-1}z^{-1}$ my^{-2}	1 $y^{-1}z$
10	m mx^{-1}	mx^{-2} mx^{-1} $mx^{-1}y^{-1}$		mx^{-1} $y^{-1}z^{-1}$	$mx^{-1}y^{-1}$ mx^{-1} my^{-1}	
11	xz^{-1}	my^{-1} mx^{-1}	$x^{-1}y$ 1 yz^{-1}	xz^{-1} $xy^{-1}z^{-1}$ z^{-1}	xz^{-1} my^{-1}	z x y
12		my^{-1}	my^{-1} mz^{-1}		$mxy^{-1}z^{-1}$ my^{-1} mxy^{-2}	z mxy^{-1}

TABLE 1. The points of $\nu_k \cap \nu_l$ for $1 \leq k \leq 6$ and $l > k$, with their Maslov indices and T-action weights.

	7	8	9	10	11	12
7						
8	$y^{-1}z^{-1}$					
9		my^{-1}				
10	$x^{-1}y^{-1}z^{-1}$ $y^{-1}z^{-1}$	$mx^{-1}z$				
11	$xy^{-1}z^{-1}$ $y^{-1}z^{-1}$ z^{-1}	y x	y	x		
12		mxy^{-1} m $my^{-1}z$	y z		my^{-1}	

TABLE 2. The points of $\nu_k \cap \nu_l$ for $7 \le k \le 12$ and $l > k$, with their Maslov indices and T-action weights.

a_1	a_2	b	$\pm$
$\nu_1 \cap \nu_2 : x$	$\nu_2 \cap \nu_5 : 1$	$\nu_1 \cap \nu_5 : x$	$+$
$\nu_1 \cap \nu_2 : x$	$\nu_2 \cap \nu_5 : x^{-1}y$	$\nu_1 \cap \nu_5 : y$	$-$
$\nu_1 \cap \nu_4 : x$	$\nu_4 \cap \nu_5 : 1$	$\nu_1 \cap \nu_5 : x$	$+$
$\nu_1 \cap \nu_4 : 1$	$\nu_4 \cap \nu_5 : 1$	$\nu_1 \cap \nu_5 : 1$	$-$
$\nu_2 \cap \nu_3 : my^{-1}$	$\nu_3 \cap \nu_6 : x^{-1}yz^{-1}$	$\nu_2 \cap \nu_6 : mx^{-1}z^{-1}$	$-$
$\nu_2 \cap \nu_3 : my^{-1}$	$\nu_3 \cap \nu_6 : x^{-1}$	$\nu_2 \cap \nu_6 : mx^{-1}y^{-1}$	$-$
$\nu_2 \cap \nu_5 : x^{-1}y$	$\nu_5 \cap \nu_6 : my^{-1}z^{-1}$	$\nu_2 \cap \nu_6 : mx^{-1}z^{-1}$	$-$
$\nu_2 \cap \nu_5 : 1$	$\nu_5 \cap \nu_6 : my^{-1}z^{-1}$	$\nu_2 \cap \nu_6 : my^{-1}z^{-1}$	$+$
$\nu_4 \cap \nu_5 : 1$	$\nu_5 \cap \nu_8 : xy^{-1}z^{-1}$	$\nu_4 \cap \nu_8 : xy^{-1}z^{-1}$	$+$
$\nu_4 \cap \nu_5 : 1$	$\nu_5 \cap \nu_8 : z^{-1}$	$\nu_4 \cap \nu_8 : z^{-1}$	$-$
$\nu_4 \cap \nu_7 : x$	$\nu_7 \cap \nu_8 : y^{-1}z^{-1}$	$\nu_4 \cap \nu_8 : xy^{-1}z^{-1}$	$+$
$\nu_4 \cap \nu_7 : 1$	$\nu_7 \cap \nu_8 : y^{-1}z^{-1}$	$\nu_4 \cap \nu_8 : y^{-1}z^{-1}$	$-$
$\nu_5 \cap \nu_6 : my^{-1}z^{-1}$	$\nu_6 \cap \nu_9 : 1$	$\nu_5 \cap \nu_9 : my^{-1}z^{-1}$	$-$
$\nu_5 \cap \nu_6 : my^{-1}z^{-1}$	$\nu_6 \cap \nu_9 : y^{-1}z$	$\nu_5 \cap \nu_9 : my^{-2}$	$-$
$\nu_5 \cap \nu_8 : z^{-1}$	$\nu_8 \cap \nu_9 : my^{-1}$	$\nu_5 \cap \nu_9 : my^{-1}z^{-1}$	$-$
$\nu_5 \cap \nu_8 : xy^{-1}z^{-1}$	$\nu_8 \cap \nu_9 : my^{-1}$	$\nu_5 \cap \nu_9 : mxy^{-2}z^{-1}$	$+$
$\nu_7 \cap \nu_8 : y^{-1}z^{-1}$	$\nu_8 \cap \nu_{11} : x$	$\nu_7 \cap \nu_{11} : xy^{-1}z^{-1}$	$+$
$\nu_7 \cap \nu_8 : y^{-1}z^{-1}$	$\nu_8 \cap \nu_{11} : y$	$\nu_7 \cap \nu_{11} : z^{-1}$	$-$
$\nu_7 \cap \nu_{10} : y^{-1}z^{-1}$	$\nu_{10} \cap \nu_{11} : x$	$\nu_7 \cap \nu_{11} : xy^{-1}z^{-1}$	$+$
$\nu_7 \cap \nu_{10} : x^{-1}y^{-1}z^{-1}$	$\nu_{10} \cap \nu_{11} : x$	$\nu_7 \cap \nu_{11} : y^{-1}z^{-1}$	$-$
$\nu_8 \cap \nu_9 : my^{-1}$	$\nu_9 \cap \nu_{12} : y$	$\nu_8 \cap \nu_{12} : m$	$-$
$\nu_8 \cap \nu_9 : my^{-1}$	$\nu_9 \cap \nu_{12} : z$	$\nu_8 \cap \nu_{12} : my^{-1}z$	$-$
$\nu_8 \cap \nu_{11} : y$	$\nu_{11} \cap \nu_{12} : my^{-1}$	$\nu_8 \cap \nu_{12} : m$	$-$
$\nu_8 \cap \nu_{11} : x$	$\nu_{11} \cap \nu_{12} : my^{-1}$	$\nu_8 \cap \nu_{12} : mxy^{-1}$	$+$
$\nu_2 \cap \nu_4 : mx^{-1}z$	$\nu_4 \cap \nu_7 : 1$	$\nu_2 \cap \nu_7 : mx^{-1}z$	$-$
$\nu_2 \cap \nu_4 : mx^{-1}z$	$\nu_4 \cap \nu_7 : x$	$\nu_2 \cap \nu_7 : mz$	$+$
$\nu_2 \cap \nu_5 : 1$	$\nu_5 \cap \nu_7 : mz$	$\nu_2 \cap \nu_7 : mz$	$+$
$\nu_2 \cap \nu_5 : x^{-1}y$	$\nu_5 \cap \nu_7 : mz$	$\nu_2 \cap \nu_7 : mx^{-1}yz$	$-$
$\nu_3 \cap \nu_5 : x^{-1}y$	$\nu_5 \cap \nu_8 : xy^{-1}z^{-1}$	$\nu_3 \cap \nu_8 : z^{-1}$	$-$
$\nu_3 \cap \nu_5 : x^{-1}y$	$\nu_5 \cap \nu_8 : z^{-1}$	$\nu_3 \cap \nu_8 : x^{-1}yz^{-1}$	$-$

$\nu_3 \cap \nu_6 : x^{-1}yz^{-1}$	$\nu_6 \cap \nu_8 : 1$	$\nu_3 \cap \nu_8 : x^{-1}yz^{-1}$	$-$
$\nu_3 \cap \nu_6 : x^{-1}$	$\nu_6 \cap \nu_8 : 1$	$\nu_3 \cap \nu_8 : x^{-1}$	$+$
$\nu_1 \cap \nu_2 : x$	$\nu_2 \cap \nu_7 : mx^{-1}yz$	$\nu_1 \cap \nu_7 : myz$	$+$
$\nu_1 \cap \nu_4 : x$	$\nu_4 \cap \nu_7 : 1$	$\nu_1 \cap \nu_7 : x$	$-$
$\nu_1 \cap \nu_4 : 1$	$\nu_4 \cap \nu_7 : x$	$\nu_1 \cap \nu_7 : x$	$+$
$\nu_1 \cap \nu_5 : y$	$\nu_5 \cap \nu_7 : mz$	$\nu_1 \cap \nu_7 : myz$	$+$
$\nu_2 \cap \nu_3 : my^{-1}$	$\nu_3 \cap \nu_8 : x^{-1}$	$\nu_2 \cap \nu_8 : mx^{-1}y^{-1}$	$+$
$\nu_2 \cap \nu_4 : mx^{-1}z$	$\nu_4 \cap \nu_8 : y^{-1}z^{-1}$	$\nu_2 \cap \nu_8 : mx^{-1}y^{-1}$	$+$
$\nu_2 \cap \nu_5 : x^{-1}y$	$\nu_5 \cap \nu_8 : xy^{-1}z^{-1}$	$\nu_2 \cap \nu_8 : z^{-1}$	$+$
$\nu_2 \cap \nu_5 : 1$	$\nu_5 \cap \nu_8 : z^{-1}$	$\nu_2 \cap \nu_8 : z^{-1}$	$-$
$\nu_2 \cap \nu_6 : mx^{-1}y^{-1}$	$\nu_6 \cap \nu_8 : 1$	$\nu_2 \cap \nu_8 : mx^{-1}y^{-1}$	$+$
$\nu_2 \cap \nu_7 : mx^{-1}z$	$\nu_7 \cap \nu_8 : y^{-1}z^{-1}$	$\nu_2 \cap \nu_8 : mx^{-1}y^{-1}$	$-$
$\nu_3 \cap \nu_5 : x^{-1}y$	$\nu_5 \cap \nu_9 : mxy^{-2}z^{-1}$	$\nu_3 \cap \nu_9 : my^{-1}z^{-1}$	$+$
$\nu_3 \cap \nu_6 : x^{-1}$	$\nu_6 \cap \nu_9 : 1$	$\nu_3 \cap \nu_9 : x^{-1}$	$-$
$\nu_3 \cap \nu_6 : x^{-1}yz^{-1}$	$\nu_6 \cap \nu_9 : y^{-1}z$	$\nu_3 \cap \nu_9 : x^{-1}$	$+$
$\nu_3 \cap \nu_8 : z^{-1}$	$\nu_8 \cap \nu_9 : my^{-1}$	$\nu_3 \cap \nu_9 : my^{-1}z^{-1}$	$-$
$\nu_1 \cap \nu_2 : x$	$\nu_2 \cap \nu_8 : z^{-1}$	$\nu_1 \cap \nu_8 : xz^{-1}$	$-$
$\nu_1 \cap \nu_4 : x$	$\nu_4 \cap \nu_8 : z^{-1}$	$\nu_1 \cap \nu_8 : xz^{-1}$	$-$
$\nu_1 \cap \nu_4 : x$	$\nu_4 \cap \nu_8 : y^{-1}z^{-1}$	$\nu_1 \cap \nu_8 : xy^{-1}z^{-1}$	$+$
$\nu_1 \cap \nu_4 : 1$	$\nu_4 \cap \nu_8 : z^{-1}$	$\nu_1 \cap \nu_8 : z^{-1}$	$+$
$\nu_1 \cap \nu_4 : 1$	$\nu_4 \cap \nu_8 : xy^{-1}z^{-1}$	$\nu_1 \cap \nu_8 : xy^{-1}z^{-1}$	$+$
$\nu_1 \cap \nu_5 : x$	$\nu_5 \cap \nu_8 : z^{-1}$	$\nu_1 \cap \nu_8 : xz^{-1}$	$+$
$\nu_1 \cap \nu_5 : y$	$\nu_5 \cap \nu_8 : xy^{-1}z^{-1}$	$\nu_1 \cap \nu_8 : xz^{-1}$	$+$
$\nu_1 \cap \nu_5 : 1$	$\nu_5 \cap \nu_8 : z^{-1}$	$\nu_1 \cap \nu_8 : z^{-1}$	$+$
$\nu_1 \cap \nu_5 : 1$	$\nu_5 \cap \nu_8 : xy^{-1}z^{-1}$	$\nu_1 \cap \nu_8 : xy^{-1}z^{-1}$	$-$
$\nu_1 \cap \nu_7 : x$	$\nu_7 \cap \nu_8 : y^{-1}z^{-1}$	$\nu_1 \cap \nu_8 : xy^{-1}z^{-1}$	$+$
$\nu_2 \cap \nu_3 : my^{-1}$	$\nu_3 \cap \nu_9 : x^{-1}$	$\nu_2 \cap \nu_9 : mx^{-1}y^{-1}$	$+$
$\nu_2 \cap \nu_5 : 1$	$\nu_5 \cap \nu_9 : my^{-2}$	$\nu_2 \cap \nu_9 : my^{-2}$	$+$
$\nu_2 \cap \nu_5 : 1$	$\nu_5 \cap \nu_9 : my^{-1}z^{-1}$	$\nu_2 \cap \nu_9 : my^{-1}z^{-1}$	$+$
$\nu_2 \cap \nu_5 : x^{-1}y$	$\nu_5 \cap \nu_9 : mxy^{-2}z^{-1}$	$\nu_2 \cap \nu_9 : my^{-1}z^{-1}$	$+$
$\nu_2 \cap \nu_5 : x^{-1}y$	$\nu_5 \cap \nu_9 : my^{-2}$	$\nu_2 \cap \nu_9 : mx^{-1}y^{-1}$	$+$
$\nu_2 \cap \nu_6 : my^{-1}z^{-1}$	$\nu_6 \cap \nu_9 : 1$	$\nu_2 \cap \nu_9 : my^{-1}z^{-1}$	$-$
$\nu_2 \cap \nu_6 : mx^{-1}z^{-1}$	$\nu_6 \cap \nu_9 : y^{-1}z$	$\nu_2 \cap \nu_9 : mx^{-1}y^{-1}$	$+$
$\nu_2 \cap \nu_6 : mx^{-1}y^{-1}$	$\nu_6 \cap \nu_9 : 1$	$\nu_2 \cap \nu_9 : mx^{-1}y^{-1}$	$-$
$\nu_2 \cap \nu_6 : my^{-1}z^{-1}$	$\nu_6 \cap \nu_9 : y^{-1}z$	$\nu_2 \cap \nu_9 : my^{-2}$	$-$
$\nu_2 \cap \nu_8 : z^{-1}$	$\nu_8 \cap \nu_9 : my^{-1}$	$\nu_2 \cap \nu_9 : my^{-1}z^{-1}$	$+$
$\nu_5 \cap \nu_7 : mz$	$\nu_7 \cap \nu_{10} : x^{-1}y^{-1}z^{-1}$	$\nu_5 \cap \nu_{10} : mx^{-1}y^{-1}$	$-$
$\nu_5 \cap \nu_7 : mz$	$\nu_7 \cap \nu_{10} : y^{-1}z^{-1}$	$\nu_5 \cap \nu_{10} : my^{-1}$	$+$
$\nu_5 \cap \nu_8 : xy^{-1}z^{-1}$	$\nu_8 \cap \nu_{10} : mx^{-1}z$	$\nu_5 \cap \nu_{10} : my^{-1}$	$+$
$\nu_5 \cap \nu_8 : z^{-1}$	$\nu_8 \cap \nu_{10} : mx^{-1}z$	$\nu_5 \cap \nu_{10} : mx^{-1}$	$-$
$\nu_6 \cap \nu_8 : 1$	$\nu_8 \cap \nu_{11} : x$	$\nu_6 \cap \nu_{11} : x$	$-$
$\nu_6 \cap \nu_8 : 1$	$\nu_8 \cap \nu_{11} : y$	$\nu_6 \cap \nu_{11} : y$	$-$
$\nu_6 \cap \nu_9 : 1$	$\nu_9 \cap \nu_{11} : y$	$\nu_6 \cap \nu_{11} : y$	$-$
$\nu_6 \cap \nu_9 : y^{-1}z$	$\nu_9 \cap \nu_{11} : y$	$\nu_6 \cap \nu_{11} : z$	$+$
$\nu_4 \cap \nu_5 : 1$	$\nu_5 \cap \nu_{10} : mx^{-1}$	$\nu_4 \cap \nu_{10} : mx^{-1}$	$+$
$\nu_4 \cap \nu_7 : x$	$\nu_7 \cap \nu_{10} : x^{-1}y^{-1}z^{-1}$	$\nu_4 \cap \nu_{10} : y^{-1}z^{-1}$	$-$
$\nu_4 \cap \nu_7 : 1$	$\nu_7 \cap \nu_{10} : y^{-1}z^{-1}$	$\nu_4 \cap \nu_{10} : y^{-1}z^{-1}$	$+$
$\nu_4 \cap \nu_8 : z^{-1}$	$\nu_8 \cap \nu_{10} : mx^{-1}z$	$\nu_4 \cap \nu_{10} : mx^{-1}$	$+$
$\nu_5 \cap \nu_6 : my^{-1}z^{-1}$	$\nu_6 \cap \nu_{11} : z$	$\nu_5 \cap \nu_{11} : my^{-1}$	$+$
$\nu_5 \cap \nu_7 : mz$	$\nu_7 \cap \nu_{11} : y^{-1}z^{-1}$	$\nu_5 \cap \nu_{11} : my^{-1}$	$+$
$\nu_5 \cap \nu_8 : z^{-1}$	$\nu_8 \cap \nu_{11} : x$	$\nu_5 \cap \nu_{11} : xz^{-1}$	$+$
$\nu_5 \cap \nu_8 : xy^{-1}z^{-1}$	$\nu_8 \cap \nu_{11} : y$	$\nu_5 \cap \nu_{11} : xz^{-1}$	$-$
$\nu_5 \cap \nu_9 : my^{-2}$	$\nu_9 \cap \nu_{11} : y$	$\nu_5 \cap \nu_{11} : my^{-1}$	$+$

$\nu_5 \cap \nu_{10} : mx^{-1}y^{-1}$	$\nu_{10} \cap \nu_{11} : x$	$\nu_5 \cap \nu_{11} : my^{-1}$	$-$
$\nu_6 \cap \nu_8 : 1$	$\nu_8 \cap \nu_{12} : mxy^{-1}$	$\nu_6 \cap \nu_{12} : mxy^{-1}$	$+$
$\nu_6 \cap \nu_9 : y^{-1}z$	$\nu_9 \cap \nu_{12} : y$	$\nu_6 \cap \nu_{12} : z$	$-$
$\nu_6 \cap \nu_9 : 1$	$\nu_9 \cap \nu_{12} : z$	$\nu_6 \cap \nu_{12} : z$	$+$
$\nu_6 \cap \nu_{11} : x$	$\nu_{11} \cap \nu_{12} : my^{-1}$	$\nu_6 \cap \nu_{12} : mxy^{-1}$	$-$
$\nu_4 \cap \nu_5 : 1$	$\nu_5 \cap \nu_{11} : xz^{-1}$	$\nu_4 \cap \nu_{11} : xz^{-1}$	$-$
$\nu_4 \cap \nu_7 : x$	$\nu_7 \cap \nu_{11} : z^{-1}$	$\nu_4 \cap \nu_{11} : xz^{-1}$	$-$
$\nu_4 \cap \nu_7 : x$	$\nu_7 \cap \nu_{11} : y^{-1}z^{-1}$	$\nu_4 \cap \nu_{11} : xy^{-1}z^{-1}$	$+$
$\nu_4 \cap \nu_7 : 1$	$\nu_7 \cap \nu_{11} : z^{-1}$	$\nu_4 \cap \nu_{11} : z^{-1}$	$+$
$\nu_4 \cap \nu_7 : 1$	$\nu_7 \cap \nu_{11} : xy^{-1}z^{-1}$	$\nu_4 \cap \nu_{11} : xy^{-1}z^{-1}$	$+$
$\nu_4 \cap \nu_8 : xy^{-1}z^{-1}$	$\nu_8 \cap \nu_{11} : y$	$\nu_4 \cap \nu_{11} : xz^{-1}$	$+$
$\nu_4 \cap \nu_8 : z^{-1}$	$\nu_8 \cap \nu_{11} : x$	$\nu_4 \cap \nu_{11} : xz^{-1}$	$+$
$\nu_4 \cap \nu_8 : y^{-1}z^{-1}$	$\nu_8 \cap \nu_{11} : y$	$\nu_4 \cap \nu_{11} : z^{-1}$	$+$
$\nu_4 \cap \nu_8 : y^{-1}z^{-1}$	$\nu_8 \cap \nu_{11} : x$	$\nu_4 \cap \nu_{11} : xy^{-1}z^{-1}$	$-$
$\nu_4 \cap \nu_{10} : y^{-1}z^{-1}$	$\nu_{10} \cap \nu_{11} : x$	$\nu_4 \cap \nu_{11} : xy^{-1}z^{-1}$	$+$
$\nu_5 \cap \nu_6 : my^{-1}z^{-1}$	$\nu_6 \cap \nu_{12} : z$	$\nu_5 \cap \nu_{12} : my^{-1}$	$+$
$\nu_5 \cap \nu_8 : xy^{-1}z^{-1}$	$\nu_8 \cap \nu_{12} : my^{-1}z$	$\nu_5 \cap \nu_{12} : mxy^{-2}$	$+$
$\nu_5 \cap \nu_8 : xy^{-1}z^{-1}$	$\nu_8 \cap \nu_{12} : m$	$\nu_5 \cap \nu_{12} : mxy^{-1}z^{-1}$	$+$
$\nu_5 \cap \nu_8 : z^{-1}$	$\nu_8 \cap \nu_{12} : mxy^{-1}$	$\nu_5 \cap \nu_{12} : mxy^{-1}z^{-1}$	$+$
$\nu_5 \cap \nu_8 : z^{-1}$	$\nu_8 \cap \nu_{12} : my^{-1}z$	$\nu_5 \cap \nu_{12} : my^{-1}$	$+$
$\nu_5 \cap \nu_9 : mxy^{-2}z^{-1}$	$\nu_9 \cap \nu_{12} : y$	$\nu_5 \cap \nu_{12} : mxy^{-1}z^{-1}$	$-$
$\nu_5 \cap \nu_9 : my^{-1}z^{-1}$	$\nu_9 \cap \nu_{12} : z$	$\nu_5 \cap \nu_{12} : my^{-1}$	$+$
$\nu_5 \cap \nu_9 : my^{-2}$	$\nu_9 \cap \nu_{12} : y$	$\nu_5 \cap \nu_{12} : my^{-1}$	$-$
$\nu_5 \cap \nu_9 : mxy^{-2}z^{-1}$	$\nu_9 \cap \nu_{12} : z$	$\nu_5 \cap \nu_{12} : mxy^{-2}$	$-$
$\nu_5 \cap \nu_{11} : xz^{-1}$	$\nu_{11} \cap \nu_{12} : my^{-1}$	$\nu_5 \cap \nu_{12} : mxy^{-1}z^{-1}$	$+$
$\nu_2 \cap \nu_4 : mx^{-1}z$	$\nu_4 \cap \nu_{10} : y^{-1}z^{-1}$	$\nu_2 \cap \nu_{10} : mx^{-1}y^{-1}$	$-$
$\nu_2 \cap \nu_5 : x^{-1}y$	$\nu_5 \cap \nu_{10} : my^{-1}$	$\nu_2 \cap \nu_{10} : mx^{-1}$	$+$
$\nu_2 \cap \nu_5 : x^{-1}y$	$\nu_5 \cap \nu_{10} : mx^{-1}y^{-1}$	$\nu_2 \cap \nu_{10} : mx^{-2}$	$-$
$\nu_2 \cap \nu_5 : 1$	$\nu_5 \cap \nu_{10} : mx^{-1}y^{-1}$	$\nu_2 \cap \nu_{10} : mx^{-1}y^{-1}$	$+$
$\nu_2 \cap \nu_5 : 1$	$\nu_5 \cap \nu_{10} : mx^{-1}$	$\nu_2 \cap \nu_{10} : mx^{-1}$	$+$
$\nu_2 \cap \nu_7 : mx^{-1}yz$	$\nu_7 \cap \nu_{10} : y^{-1}z^{-1}$	$\nu_2 \cap \nu_{10} : mx^{-1}$	$-$
$\nu_2 \cap \nu_7 : mx^{-1}yz$	$\nu_7 \cap \nu_{10} : x^{-1}y^{-1}z^{-1}$	$\nu_2 \cap \nu_{10} : mx^{-2}$	$-$
$\nu_2 \cap \nu_7 : mz$	$\nu_7 \cap \nu_{10} : x^{-1}y^{-1}z^{-1}$	$\nu_2 \cap \nu_{10} : mx^{-1}y^{-1}$	$-$
$\nu_2 \cap \nu_7 : mx^{-1}z$	$\nu_7 \cap \nu_{10} : y^{-1}z^{-1}$	$\nu_2 \cap \nu_{10} : mx^{-1}y^{-1}$	$-$
$\nu_2 \cap \nu_8 : z^{-1}$	$\nu_8 \cap \nu_{10} : mx^{-1}z$	$\nu_2 \cap \nu_{10} : mx^{-1}$	$+$
$\nu_3 \cap \nu_5 : x^{-1}y$	$\nu_5 \cap \nu_{11} : xz^{-1}$	$\nu_3 \cap \nu_{11} : yz^{-1}$	$-$
$\nu_3 \cap \nu_6 : x^{-1}yz^{-1}$	$\nu_6 \cap \nu_{11} : z$	$\nu_3 \cap \nu_{11} : x^{-1}y$	$+$
$\nu_3 \cap \nu_6 : x^{-1}yz^{-1}$	$\nu_6 \cap \nu_{11} : x$	$\nu_3 \cap \nu_{11} : yz^{-1}$	$+$
$\nu_3 \cap \nu_6 : x^{-1}$	$\nu_6 \cap \nu_{11} : y$	$\nu_3 \cap \nu_{11} : x^{-1}y$	$+$
$\nu_3 \cap \nu_6 : x^{-1}$	$\nu_6 \cap \nu_{11} : x$	$\nu_3 \cap \nu_{11} : 1$	$-$
$\nu_3 \cap \nu_8 : x^{-1}yz^{-1}$	$\nu_8 \cap \nu_{11} : x$	$\nu_3 \cap \nu_{11} : yz^{-1}$	$+$
$\nu_3 \cap \nu_8 : x^{-1}$	$\nu_8 \cap \nu_{11} : x$	$\nu_3 \cap \nu_{11} : 1$	$+$
$\nu_3 \cap \nu_8 : z^{-1}$	$\nu_8 \cap \nu_{11} : y$	$\nu_3 \cap \nu_{11} : yz^{-1}$	$-$
$\nu_3 \cap \nu_8 : x^{-1}$	$\nu_8 \cap \nu_{11} : y$	$\nu_3 \cap \nu_{11} : x^{-1}y$	$-$
$\nu_3 \cap \nu_9 : x^{-1}$	$\nu_9 \cap \nu_{11} : y$	$\nu_3 \cap \nu_{11} : x^{-1}y$	$+$
$\nu_1 \cap \nu_2 : x$	$\nu_2 \cap \nu_{10} : mx^{-1}$	$\nu_1 \cap \nu_{10} : m$	$+$
$\nu_1 \cap \nu_2 : x$	$\nu_2 \cap \nu_{10} : mx^{-2}$	$\nu_1 \cap \nu_{10} : mx^{-1}$	$+$
$\nu_1 \cap \nu_4 : x$	$\nu_4 \cap \nu_{10} : mx^{-1}$	$\nu_1 \cap \nu_{10} : m$	$+$
$\nu_1 \cap \nu_4 : 1$	$\nu_4 \cap \nu_{10} : mx^{-1}$	$\nu_1 \cap \nu_{10} : mx^{-1}$	$+$
$\nu_1 \cap \nu_5 : y$	$\nu_5 \cap \nu_{10} : mx^{-1}y^{-1}$	$\nu_1 \cap \nu_{10} : mx^{-1}$	$+$
$\nu_1 \cap \nu_5 : 1$	$\nu_5 \cap \nu_{10} : mx^{-1}$	$\nu_1 \cap \nu_{10} : mx^{-1}$	$-$
$\nu_1 \cap \nu_5 : y$	$\nu_5 \cap \nu_{10} : my^{-1}$	$\nu_1 \cap \nu_{10} : m$	$-$
$\nu_1 \cap \nu_5 : x$	$\nu_5 \cap \nu_{10} : mx^{-1}$	$\nu_1 \cap \nu_{10} : m$	$+$

$\nu_1 \cap \nu_7 : myz$	$\nu_7 \cap \nu_{10} : x^{-1}y^{-1}z^{-1}$	$\nu_1 \cap \nu_{10} : mx^{-1}$	$-$
$\nu_1 \cap \nu_7 : myz$	$\nu_7 \cap \nu_{10} : y^{-1}z^{-1}$	$\nu_1 \cap \nu_{10} : m$	$-$
$\nu_1 \cap \nu_8 : xz^{-1}$	$\nu_8 \cap \nu_{10} : mx^{-1}z$	$\nu_1 \cap \nu_{10} : m$	$-$
$\nu_1 \cap \nu_8 : z^{-1}$	$\nu_8 \cap \nu_{10} : mx^{-1}z$	$\nu_1 \cap \nu_{10} : mx^{-1}$	$+$
$\nu_2 \cap \nu_3 : my^{-1}$	$\nu_3 \cap \nu_{11} : x^{-1}y$	$\nu_2 \cap \nu_{11} : mx^{-1}$	$-$
$\nu_2 \cap \nu_3 : my^{-1}$	$\nu_3 \cap \nu_{11} : 1$	$\nu_2 \cap \nu_{11} : my^{-1}$	$+$
$\nu_2 \cap \nu_4 : mx^{-1}z$	$\nu_4 \cap \nu_{11} : z^{-1}$	$\nu_2 \cap \nu_{11} : mx^{-1}$	$+$
$\nu_2 \cap \nu_4 : mx^{-1}z$	$\nu_4 \cap \nu_{11} : xy^{-1}z^{-1}$	$\nu_2 \cap \nu_{11} : my^{-1}$	$-$
$\nu_2 \cap \nu_5 : 1$	$\nu_5 \cap \nu_{11} : my^{-1}$	$\nu_2 \cap \nu_{11} : my^{-1}$	$+$
$\nu_2 \cap \nu_5 : x^{-1}y$	$\nu_5 \cap \nu_{11} : my^{-1}$	$\nu_2 \cap \nu_{11} : mx^{-1}$	$+$
$\nu_2 \cap \nu_6 : my^{-1}z^{-1}$	$\nu_6 \cap \nu_{11} : z$	$\nu_2 \cap \nu_{11} : my^{-1}$	$+$
$\nu_2 \cap \nu_6 : mx^{-1}y^{-1}$	$\nu_6 \cap \nu_{11} : x$	$\nu_2 \cap \nu_{11} : my^{-1}$	$-$
$\nu_2 \cap \nu_6 : mx^{-1}z^{-1}$	$\nu_6 \cap \nu_{11} : z$	$\nu_2 \cap \nu_{11} : mx^{-1}$	$-$
$\nu_2 \cap \nu_6 : mx^{-1}y^{-1}$	$\nu_6 \cap \nu_{11} : y$	$\nu_2 \cap \nu_{11} : mx^{-1}$	$-$
$\nu_2 \cap \nu_7 : mx^{-1}z$	$\nu_7 \cap \nu_{11} : xy^{-1}z^{-1}$	$\nu_2 \cap \nu_{11} : my^{-1}$	$-$
$\nu_2 \cap \nu_7 : mz$	$\nu_7 \cap \nu_{11} : y^{-1}z^{-1}$	$\nu_2 \cap \nu_{11} : my^{-1}$	$+$
$\nu_2 \cap \nu_7 : mx^{-1}yz$	$\nu_7 \cap \nu_{11} : y^{-1}z^{-1}$	$\nu_2 \cap \nu_{11} : mx^{-1}$	$-$
$\nu_2 \cap \nu_7 : mx^{-1}z$	$\nu_7 \cap \nu_{11} : z^{-1}$	$\nu_2 \cap \nu_{11} : mx^{-1}$	$+$
$\nu_2 \cap \nu_8 : mx^{-1}y^{-1}$	$\nu_8 \cap \nu_{11} : y$	$\nu_2 \cap \nu_{11} : mx^{-1}$	$-$
$\nu_2 \cap \nu_8 : mx^{-1}y^{-1}$	$\nu_8 \cap \nu_{11} : x$	$\nu_2 \cap \nu_{11} : my^{-1}$	$-$
$\nu_2 \cap \nu_9 : my^{-2}$	$\nu_9 \cap \nu_{11} : y$	$\nu_2 \cap \nu_{11} : my^{-1}$	$+$
$\nu_2 \cap \nu_9 : mx^{-1}y^{-1}$	$\nu_9 \cap \nu_{11} : y$	$\nu_2 \cap \nu_{11} : mx^{-1}$	$+$
$\nu_2 \cap \nu_{10} : mx^{-1}y^{-1}$	$\nu_{10} \cap \nu_{11} : x$	$\nu_2 \cap \nu_{11} : my^{-1}$	$-$
$\nu_2 \cap \nu_{10} : mx^{-2}$	$\nu_{10} \cap \nu_{11} : x$	$\nu_2 \cap \nu_{11} : mx^{-1}$	$+$
$\nu_3 \cap \nu_5 : x^{-1}y$	$\nu_5 \cap \nu_{12} : mxy^{-1}z^{-1}$	$\nu_3 \cap \nu_{12} : mz^{-1}$	$+$
$\nu_3 \cap \nu_5 : x^{-1}y$	$\nu_5 \cap \nu_{12} : mxy^{-2}$	$\nu_3 \cap \nu_{12} : my^{-1}$	$-$
$\nu_3 \cap \nu_6 : x^{-1}$	$\nu_6 \cap \nu_{12} : mxy^{-1}$	$\nu_3 \cap \nu_{12} : my^{-1}$	$-$
$\nu_3 \cap \nu_6 : x^{-1}yz^{-1}$	$\nu_6 \cap \nu_{12} : mxy^{-1}$	$\nu_3 \cap \nu_{12} : mz^{-1}$	$+$
$\nu_3 \cap \nu_8 : z^{-1}$	$\nu_8 \cap \nu_{12} : my^{-1}z$	$\nu_3 \cap \nu_{12} : my^{-1}$	$+$
$\nu_3 \cap \nu_8 : x^{-1}$	$\nu_8 \cap \nu_{12} : mxy^{-1}$	$\nu_3 \cap \nu_{12} : my^{-1}$	$-$
$\nu_3 \cap \nu_8 : z^{-1}$	$\nu_8 \cap \nu_{12} : m$	$\nu_3 \cap \nu_{12} : mz^{-1}$	$-$
$\nu_3 \cap \nu_8 : x^{-1}yz^{-1}$	$\nu_8 \cap \nu_{12} : mxy^{-1}$	$\nu_3 \cap \nu_{12} : mz^{-1}$	$-$
$\nu_3 \cap \nu_9 : my^{-1}z^{-1}$	$\nu_9 \cap \nu_{12} : y$	$\nu_3 \cap \nu_{12} : mz^{-1}$	$-$
$\nu_3 \cap \nu_9 : my^{-1}z^{-1}$	$\nu_9 \cap \nu_{12} : z$	$\nu_3 \cap \nu_{12} : my^{-1}$	$+$
$\nu_3 \cap \nu_{11} : yz^{-1}$	$\nu_{11} \cap \nu_{12} : my^{-1}$	$\nu_3 \cap \nu_{12} : mz^{-1}$	$-$
$\nu_3 \cap \nu_{11} : 1$	$\nu_{11} \cap \nu_{12} : my^{-1}$	$\nu_3 \cap \nu_{12} : my^{-1}$	$-$
$\nu_1 \cap \nu_4 : 1$	$\nu_4 \cap \nu_{11} : xz^{-1}$	$\nu_1 \cap \nu_{11} : xz^{-1}$	$+$
$\nu_1 \cap \nu_4 : x$	$\nu_4 \cap \nu_{11} : z^{-1}$	$\nu_1 \cap \nu_{11} : xz^{-1}$	$+$
$\nu_1 \cap \nu_5 : 1$	$\nu_5 \cap \nu_{11} : xz^{-1}$	$\nu_1 \cap \nu_{11} : xz^{-1}$	$+$
$\nu_1 \cap \nu_7 : x$	$\nu_7 \cap \nu_{11} : z^{-1}$	$\nu_1 \cap \nu_{11} : xz^{-1}$	$-$
$\nu_1 \cap \nu_8 : z^{-1}$	$\nu_8 \cap \nu_{11} : x$	$\nu_1 \cap \nu_{11} : xz^{-1}$	$+$
$\nu_1 \cap \nu_8 : xy^{-1}z^{-1}$	$\nu_8 \cap \nu_{11} : y$	$\nu_1 \cap \nu_{11} : xz^{-1}$	$+$
$\nu_2 \cap \nu_5 : x^{-1}y$	$\nu_5 \cap \nu_{12} : mxy^{-2}$	$\nu_2 \cap \nu_{12} : my^{-1}$	$+$
$\nu_2 \cap \nu_5 : 1$	$\nu_5 \cap \nu_{12} : my^{-1}$	$\nu_2 \cap \nu_{12} : my^{-1}$	$-$
$\nu_2 \cap \nu_6 : my^{-1}z^{-1}$	$\nu_6 \cap \nu_{12} : z$	$\nu_2 \cap \nu_{12} : my^{-1}$	$-$
$\nu_2 \cap \nu_8 : z^{-1}$	$\nu_8 \cap \nu_{12} : my^{-1}z$	$\nu_2 \cap \nu_{12} : my^{-1}$	$+$
$\nu_2 \cap \nu_9 : my^{-2}$	$\nu_9 \cap \nu_{12} : y$	$\nu_2 \cap \nu_{12} : my^{-1}$	$+$
$\nu_2 \cap \nu_9 : my^{-1}z^{-1}$	$\nu_9 \cap \nu_{12} : z$	$\nu_2 \cap \nu_{12} : my^{-1}$	$-$

TABLE 3. Products μ^2 in the directed A_∞-category $\mathcal{C}_{\overrightarrow{12}}$. The notation means that $\mu^2(a_2, a_1) = \pm b$.

a_1	a_2	a_3	b	$\pm$
$\nu_1 \cap \nu_2 : x$	$\nu_2 \cap \nu_3 : my^{-1}$	$\nu_3 \cap \nu_5 : x^{-1}y$	$\nu_1 \cap \nu_5 : 1$	$+$
$\nu_2 \cap \nu_4 : mx^{-1}z$	$\nu_4 \cap \nu_5 : 1$	$\nu_5 \cap \nu_6 : my^{-1}z^{-1}$	$\nu_2 \cap \nu_6 : mx^{-1}y^{-1}$	$-$
$\nu_4 \cap \nu_5 : 1$	$\nu_5 \cap \nu_6 : my^{-1}z^{-1}$	$\nu_6 \cap \nu_8 : 1$	$\nu_4 \cap \nu_8 : y^{-1}z^{-1}$	$+$
$\nu_5 \cap \nu_7 : mz$	$\nu_7 \cap \nu_8 : y^{-1}z^{-1}$	$\nu_8 \cap \nu_9 : my^{-1}$	$\nu_5 \cap \nu_9 : my^{-2}$	$-$
$\nu_7 \cap \nu_8 : y^{-1}z^{-1}$	$\nu_8 \cap \nu_9 : my^{-1}$	$\nu_9 \cap \nu_{11} : y$	$\nu_7 \cap \nu_{11} : y^{-1}z^{-1}$	$+$
$\nu_8 \cap \nu_{10} : mx^{-1}z$	$\nu_{10} \cap \nu_{11} : x$	$\nu_{11} \cap \nu_{12} : my^{-1}$	$\nu_8 \cap \nu_{12} : my^{-1}z$	$-$
$\nu_1 \cap \nu_2 : x$	$\nu_2 \cap \nu_3 : my^{-1}$	$\nu_3 \cap \nu_8 : x^{-1}yz^{-1}$	$\nu_1 \cap \nu_8 : z^{-1}$	$-$
$\nu_1 \cap \nu_2 : x$	$\nu_2 \cap \nu_3 : my^{-1}$	$\nu_3 \cap \nu_8 : z^{-1}$	$\nu_1 \cap \nu_8 : xy^{-1}z^{-1}$	$+$
$\nu_2 \cap \nu_3 : my^{-1}$	$\nu_3 \cap \nu_5 : x^{-1}y$	$\nu_5 \cap \nu_7 : mz$	$\nu_2 \cap \nu_7 : mx^{-1}z$	$+$
$\nu_2 \cap \nu_7 : mz$	$\nu_7 \cap \nu_8 : y^{-1}z^{-1}$	$\nu_8 \cap \nu_9 : my^{-1}$	$\nu_2 \cap \nu_9 : my^{-2}$	$-$
$\nu_2 \cap \nu_7 : mx^{-1}yz$	$\nu_7 \cap \nu_8 : y^{-1}z^{-1}$	$\nu_8 \cap \nu_9 : my^{-1}$	$\nu_2 \cap \nu_9 : mx^{-1}y^{-1}$	$+$
$\nu_1 \cap \nu_2 : x$	$\nu_2 \cap \nu_6 : my^{-1}z^{-1}$	$\nu_6 \cap \nu_8 : 1$	$\nu_1 \cap \nu_8 : xy^{-1}z^{-1}$	$+$
$\nu_1 \cap \nu_2 : x$	$\nu_2 \cap \nu_6 : mx^{-1}z^{-1}$	$\nu_6 \cap \nu_8 : 1$	$\nu_1 \cap \nu_8 : z^{-1}$	$+$
$\nu_1 \cap \nu_5 : x$	$\nu_5 \cap \nu_6 : my^{-1}z^{-1}$	$\nu_6 \cap \nu_8 : 1$	$\nu_1 \cap \nu_8 : xy^{-1}z^{-1}$	$+$
$\nu_1 \cap \nu_5 : y$	$\nu_5 \cap \nu_6 : my^{-1}z^{-1}$	$\nu_6 \cap \nu_8 : 1$	$\nu_1 \cap \nu_8 : z^{-1}$	$+$
$\nu_2 \cap \nu_4 : mx^{-1}z$	$\nu_4 \cap \nu_5 : 1$	$\nu_5 \cap \nu_9 : my^{-1}z^{-1}$	$\nu_2 \cap \nu_9 : mx^{-1}y^{-1}$	$-$
$\nu_2 \cap \nu_4 : mx^{-1}z$	$\nu_4 \cap \nu_5 : 1$	$\nu_5 \cap \nu_9 : mxy^{-2}z^{-1}$	$\nu_2 \cap \nu_9 : my^{-2}$	$+$
$\nu_2 \cap \nu_4 : mx^{-1}z$	$\nu_4 \cap \nu_8 : xy^{-1}z^{-1}$	$\nu_8 \cap \nu_9 : my^{-1}$	$\nu_2 \cap \nu_9 : my^{-2}$	$+$
$\nu_2 \cap \nu_4 : mx^{-1}z$	$\nu_4 \cap \nu_8 : z^{-1}$	$\nu_8 \cap \nu_9 : my^{-1}$	$\nu_2 \cap \nu_9 : mx^{-1}y^{-1}$	$-$
$\nu_3 \cap \nu_5 : x^{-1}y$	$\nu_5 \cap \nu_7 : mz$	$\nu_7 \cap \nu_8 : y^{-1}z^{-1}$	$\nu_3 \cap \nu_8 : x^{-1}$	$+$
$\nu_4 \cap \nu_5 : 1$	$\nu_5 \cap \nu_6 : my^{-1}z^{-1}$	$\nu_6 \cap \nu_{11} : y$	$\nu_4 \cap \nu_{11} : z^{-1}$	$-$
$\nu_4 \cap \nu_5 : 1$	$\nu_5 \cap \nu_6 : my^{-1}z^{-1}$	$\nu_6 \cap \nu_{11} : x$	$\nu_4 \cap \nu_{11} : xy^{-1}z^{-1}$	$+$
$\nu_5 \cap \nu_6 : my^{-1}z^{-1}$	$\nu_6 \cap \nu_8 : 1$	$\nu_8 \cap \nu_{10} : mx^{-1}z$	$\nu_5 \cap \nu_{10} : mx^{-1}y^{-1}$	$+$
$\nu_5 \cap \nu_{10} : my^{-1}$	$\nu_{10} \cap \nu_{11} : x$	$\nu_{11} \cap \nu_{12} : my^{-1}$	$\nu_5 \cap \nu_{12} : mxy^{-2}$	$-$
$\nu_5 \cap \nu_{10} : mx^{-1}$	$\nu_{10} \cap \nu_{11} : x$	$\nu_{11} \cap \nu_{12} : my^{-1}$	$\nu_5 \cap \nu_{12} : my^{-1}$	$+$
$\nu_4 \cap \nu_5 : 1$	$\nu_5 \cap \nu_9 : mxy^{-2}z^{-1}$	$\nu_9 \cap \nu_{11} : y$	$\nu_4 \cap \nu_{11} : xy^{-1}z^{-1}$	$+$
$\nu_4 \cap \nu_5 : 1$	$\nu_5 \cap \nu_9 : my^{-1}z^{-1}$	$\nu_9 \cap \nu_{11} : y$	$\nu_4 \cap \nu_{11} : z^{-1}$	$+$
$\nu_4 \cap \nu_8 : xy^{-1}z^{-1}$	$\nu_8 \cap \nu_9 : my^{-1}$	$\nu_9 \cap \nu_{11} : y$	$\nu_4 \cap \nu_{11} : xy^{-1}z^{-1}$	$+$
$\nu_4 \cap \nu_8 : z^{-1}$	$\nu_8 \cap \nu_9 : my^{-1}$	$\nu_9 \cap \nu_{11} : y$	$\nu_4 \cap \nu_{11} : z^{-1}$	$+$
$\nu_5 \cap \nu_7 : mz$	$\nu_7 \cap \nu_8 : y^{-1}z^{-1}$	$\nu_8 \cap \nu_{12} : m$	$\nu_5 \cap \nu_{12} : my^{-1}$	$-$
$\nu_5 \cap \nu_7 : mz$	$\nu_7 \cap \nu_8 : y^{-1}z^{-1}$	$\nu_8 \cap \nu_{12} : mxy^{-1}$	$\nu_5 \cap \nu_{12} : mxy^{-2}$	$+$
$\nu_5 \cap \nu_7 : mz$	$\nu_7 \cap \nu_{11} : xy^{-1}z^{-1}$	$\nu_{11} \cap \nu_{12} : my^{-1}$	$\nu_5 \cap \nu_{12} : mxy^{-2}$	$+$
$\nu_5 \cap \nu_7 : mz$	$\nu_7 \cap \nu_{11} : z^{-1}$	$\nu_{11} \cap \nu_{12} : my^{-1}$	$\nu_5 \cap \nu_{12} : my^{-1}$	$-$
$\nu_6 \cap \nu_8 : 1$	$\nu_8 \cap \nu_{10} : mx^{-1}z$	$\nu_{10} \cap \nu_{11} : x$	$\nu_6 \cap \nu_{11} : z$	$+$
$\nu_1 \cap \nu_2 : x$	$\nu_2 \cap \nu_3 : my^{-1}$	$\nu_3 \cap \nu_{11} : yz^{-1}$	$\nu_1 \cap \nu_{11} : xz^{-1}$	$-$
$\nu_2 \cap \nu_{10} : mx^{-1}$	$\nu_{10} \cap \nu_{11} : x$	$\nu_{11} \cap \nu_{12} : my^{-1}$	$\nu_2 \cap \nu_{12} : my^{-1}$	$-$
$\nu_1 \cap \nu_2 : x$	$\nu_2 \cap \nu_6 : mx^{-1}z^{-1}$	$\nu_6 \cap \nu_{11} : x$	$\nu_1 \cap \nu_{11} : xz^{-1}$	$-$
$\nu_1 \cap \nu_2 : x$	$\nu_2 \cap \nu_6 : my^{-1}z^{-1}$	$\nu_6 \cap \nu_{11} : y$	$\nu_1 \cap \nu_{11} : xz^{-1}$	$-$
$\nu_1 \cap \nu_2 : x$	$\nu_2 \cap \nu_9 : my^{-1}z^{-1}$	$\nu_9 \cap \nu_{11} : y$	$\nu_1 \cap \nu_{11} : xz^{-1}$	$+$
$\nu_1 \cap \nu_5 : y$	$\nu_5 \cap \nu_6 : my^{-1}z^{-1}$	$\nu_6 \cap \nu_{11} : x$	$\nu_1 \cap \nu_{11} : xz^{-1}$	$-$
$\nu_1 \cap \nu_5 : x$	$\nu_5 \cap \nu_6 : my^{-1}z^{-1}$	$\nu_6 \cap \nu_{11} : y$	$\nu_1 \cap \nu_{11} : xz^{-1}$	$-$
$\nu_2 \cap \nu_3 : my^{-1}$	$\nu_3 \cap \nu_5 : x^{-1}y$	$\nu_5 \cap \nu_{10} : mx^{-1}$	$\nu_2 \cap \nu_{10} : mx^{-2}$	$-$
$\nu_2 \cap \nu_3 : my^{-1}$	$\nu_3 \cap \nu_5 : x^{-1}y$	$\nu_5 \cap \nu_{10} : my^{-1}$	$\nu_2 \cap \nu_{10} : mx^{-1}y^{-1}$	$-$
$\nu_2 \cap \nu_3 : my^{-1}$	$\nu_3 \cap \nu_8 : z^{-1}$	$\nu_8 \cap \nu_{10} : mx^{-1}z$	$\nu_2 \cap \nu_{10} : mx^{-1}y^{-1}$	$+$
$\nu_2 \cap \nu_3 : my^{-1}$	$\nu_3 \cap \nu_8 : x^{-1}yz^{-1}$	$\nu_8 \cap \nu_{10} : mx^{-1}z$	$\nu_2 \cap \nu_{10} : mx^{-2}$	$-$
$\nu_2 \cap \nu_6 : my^{-1}z^{-1}$	$\nu_6 \cap \nu_8 : 1$	$\nu_8 \cap \nu_{10} : mx^{-1}z$	$\nu_2 \cap \nu_{10} : mx^{-1}y^{-1}$	$+$
$\nu_2 \cap \nu_6 : mx^{-1}z^{-1}$	$\nu_6 \cap \nu_8 : 1$	$\nu_8 \cap \nu_{10} : mx^{-1}z$	$\nu_2 \cap \nu_{10} : mx^{-2}$	$+$
$\nu_2 \cap \nu_4 : mx^{-1}z$	$\nu_4 \cap \nu_8 : xy^{-1}z^{-1}$	$\nu_8 \cap \nu_{12} : m$	$\nu_2 \cap \nu_{12} : my^{-1}$	$-$
$\nu_2 \cap \nu_4 : mx^{-1}z$	$\nu_4 \cap \nu_8 : z^{-1}$	$\nu_8 \cap \nu_{12} : mxy^{-1}$	$\nu_2 \cap \nu_{12} : my^{-1}$	$+$
$\nu_3 \cap \nu_5 : x^{-1}y$	$\nu_5 \cap \nu_7 : mz$	$\nu_7 \cap \nu_{11} : z^{-1}$	$\nu_3 \cap \nu_{11} : x^{-1}y$	$+$
$\nu_3 \cap \nu_5 : x^{-1}y$	$\nu_5 \cap \nu_7 : mz$	$\nu_7 \cap \nu_{11} : xy^{-1}z^{-1}$	$\nu_3 \cap \nu_{11} : 1$	$+$
$\nu_1 \cap \nu_8 : xz^{-1}$	$\nu_8 \cap \nu_9 : my^{-1}$	$\nu_9 \cap \nu_{11} : y$	$\nu_1 \cap \nu_{11} : xz^{-1}$	$-$

$\nu_3 \cap \nu_5 : x^{-1}y$	$\nu_5 \cap \nu_{10} : mx^{-1}$	$\nu_{10} \cap \nu_{11} : x$	$\nu_3 \cap \nu_{11} : x^{-1}y$	$-$
$\nu_3 \cap \nu_5 : x^{-1}y$	$\nu_5 \cap \nu_{10} : my^{-1}$	$\nu_{10} \cap \nu_{11} : x$	$\nu_3 \cap \nu_{11} : 1$	$-$
$\nu_3 \cap \nu_8 : x^{-1}yz^{-1}$	$\nu_8 \cap \nu_{10} : mx^{-1}z$	$\nu_{10} \cap \nu_{11} : x$	$\nu_3 \cap \nu_{11} : x^{-1}y$	$-$
$\nu_3 \cap \nu_8 : z^{-1}$	$\nu_8 \cap \nu_{10} : mx^{-1}z$	$\nu_{10} \cap \nu_{11} : x$	$\nu_3 \cap \nu_{11} : 1$	$+$
$\nu_1 \cap \nu_5 : x$	$\nu_5 \cap \nu_9 : my^{-1}z^{-1}$	$\nu_9 \cap \nu_{11} : y$	$\nu_1 \cap \nu_{11} : xz^{-1}$	$+$
$\nu_1 \cap \nu_5 : y$	$\nu_5 \cap \nu_9 : mxy^{-2}z^{-1}$	$\nu_9 \cap \nu_{11} : y$	$\nu_1 \cap \nu_{11} : xz^{-1}$	$-$
$\nu_2 \cap \nu_4 : mx^{-1}z$	$\nu_4 \cap \nu_5 : 1$	$\nu_5 \cap \nu_{12} : mxy^{-1}z^{-1}$	$\nu_2 \cap \nu_{12} : my^{-1}$	$-$
$\nu_2 \cap \nu_4 : mx^{-1}z$	$\nu_4 \cap \nu_{11} : xz^{-1}$	$\nu_{11} \cap \nu_{12} : my^{-1}$	$\nu_2 \cap \nu_{12} : my^{-1}$	$+$
$\nu_2 \cap \nu_7 : mx^{-1}yz$	$\nu_7 \cap \nu_8 : y^{-1}z^{-1}$	$\nu_8 \cap \nu_{12} : mxy^{-1}$	$\nu_2 \cap \nu_{12} : my^{-1}$	$-$
$\nu_2 \cap \nu_7 : mz$	$\nu_7 \cap \nu_8 : y^{-1}z^{-1}$	$\nu_8 \cap \nu_{12} : m$	$\nu_2 \cap \nu_{12} : my^{-1}$	$+$
$\nu_2 \cap \nu_7 : mz$	$\nu_7 \cap \nu_{11} : z^{-1}$	$\nu_{11} \cap \nu_{12} : my^{-1}$	$\nu_2 \cap \nu_{12} : my^{-1}$	$+$
$\nu_2 \cap \nu_7 : mx^{-1}yz$	$\nu_7 \cap \nu_{11} : xy^{-1}z^{-1}$	$\nu_{11} \cap \nu_{12} : my^{-1}$	$\nu_2 \cap \nu_{12} : my^{-1}$	$-$

TABLE 4. Products μ^3 in $\mathcal{C}_{\overrightarrow{12}}$, written in the same way as in the previous table.

REFERENCES

[1] V. I. Arnol'd, *Some remarks on symplectic monodromy of Milnor fibrations*, The Floer Memorial Volume (H. Hofer, C. Taubes, A. Weinstein, and E. Zehnder, eds.), Progress in Mathematics, vol. 133, Birkhäuser, 1995, pp. 99–104.

[2] P. S. Aspinwall, *Some navigation rules for D-brane monodromy*, J. Math. Phys. **42** (2001), 5534–5552.

[3] D. Auroux, *Symplectic 4-manifolds as branched coverings of* $\mathbb{C}\mathrm{P}^2$., Invent. Math. **139** (2000), 551–602.

[4] ——, *Symplectic maps to projective spaces and symplectic invariants*, Turkish J. Math. **25** (2001), 1–42.

[5] D. Auroux, S. Donaldson, L. Katzarkov, and M. Yotov, *Fundamental groups of complements of plane curves and symplectic invariants*, Preprint math.GT/0203183.

[6] P. Balmer and M. Schlichting, *Idempotent completion of triangulated categories*, J. Algebra **236** (2001), 819–834.

[7] A. A. Beilinson, *Coherent sheaves on* $\mathbb{P}^n$ *and problems of linear algebra*, Funct. Anal. Appl. **12** (1978), 214–216.

[8] R. Blattner and S. Montgomery, *A duality theorem for Hopf module algebras*, J. Algebra **95** (1985), 153–172.

[9] A. Bondal and M. Van den Bergh, *Generators and representability of functors in commutative and noncommutative geometry*, Preprint math.AG/0204218.

[10] Yu. Chekanov, *Differential algebras of Legendrian links*, Invent. Math. **150** (2002), 441–483.

[11] S. K. Donaldson, *Lefschetz pencils on symplectic manifolds*, J. Differential Geom. **53** (1999), 205–236.

[12] ——, *Polynomials, vanishing cycles, and Floer homology*, Mathematics: frontiers and perspectives, Amer. Math. Soc., 2000, pp. 55–64.

[13] M. Douglas, S. Govindarajan, T. Jayaraman, and A. Tomasiello, *D-branes on Calabi-Yau manifolds and superpotentials*, Preprint hep-th/0203173.

[14] K. Fukaya, *Asymptotic analysis, multivalued Morse theory, and mirror symmetry*, Preprint 2002.

[15] ——, *Morse homotopy, A_∞-categories, and Floer homologies*, Proceedings of GARC workshop on Geometry and Topology (H. J. Kim, ed.), Seoul National University, 1993.

[16] ——, *Deformation theory, homological algebra, and mirror symmetry*, Geometry and physics of branes (Como, 2001), IOP, 2003, pp. 121–209.

[17] ———, *Galois symmetry on Floer cohomology*, Turkish J. Math. **27** (2003), 11–32.

[18] K. Fukaya, Y.-G. Oh, H. Ohta, and K. Ono, *Lagrangian intersection Floer theory - anomaly and obstruction*, Preprint, 2000.

[19] Kenji Fukaya, *Mirror symmetry of abelian varieties and multi-theta functions*, J. Algebraic Geom. **11** (2002), 393–512.

[20] S. Gelfand and Yu. Manin, *Methods of homological algebra*, Springer, 1996.

[21] E. Getzler and J. D. S. Jones, *A_∞-algebras and the cyclic bar complex*, Illinois J. Math. **34** (1990), 256–283.

[22] W. Goldman and J. Millson, *The deformation theory of the fundamental group of compact Kähler manifolds*, IHES Publ. Math. **67** (1988), 43–96.

[23] R. Gompf and A. Stipsicz, *4-manifolds and Kirby calculus*, Amer. Math. Soc., 1999.

[24] M. Gross, *Examples of special Lagrangian fibrations*, Symplectic geometry and mirror symmetry (Seoul), World Sci. Publishing, 2001, pp. 81–109.

[25] J. Guccione and J. Guccione, *Hochschild cohomology of triangular matrix algebras*, Preprint math.KT/0104068.

[26] V. K. A. M. Gugenheim, L. A. Lambe, and J. D. Stasheff, *Algebraic aspects of Chen's twisting cochain*, Illinois J. Math. **34** (1990), 485–502.

[27] S. Halperin and J. Stasheff, *Obstructions to homotopy equivalences*, Advances in Math. **32** (1979), 233–279.

[28] R. Hartshorne, *Algebraic geometry*, Springer, 1977.

[29] G. Hochschild, B. Kostant, and A. Rosenberg, *Differential forms on regular affine algebras*, Trans. Amer. Math. Soc. **102** (1962), 383–408.

[30] H. Hofer and D. Salamon, *Floer homology and Novikov rings*, The Floer memorial volume (H. Hofer, C. Taubes, A. Weinstein, and E. Zehnder, eds.), Progress in Mathematics, vol. 133, Birkhäuser, 1995, pp. 483–524.

[31] K. Hori, A. Iqbal, and C. Vafa, *D-branes and mirror symmetry*, Preprint hep-th/0005247.

[32] T. V. Kadeishvili, *The structure of the A_∞-algebra, and the Hochschild and Harrison cohomologies*, Trudy Tbiliss. Mat. Inst. Razmadze Akad. Nauk Gruzin. SSR **91** (1988), 19–27 (Russian).

[33] M. Kapranov and E. Vasserot, *Kleinian singularities, derived categories and Hall algebras*, Math. Ann. **316** (2000), 565–576.

[34] B. Keller, *Introduction to A-infinity algebras and modules*, Homology Homotopy Appl. (electronic) **3** (2001), 1–35.

[35] M. Kontsevich, *Lectures at ENS Paris, Spring 1998*, set of notes taken by J. Bellaiche, J.-F. Dat, I. Marin, G. Racinet and H. Randriambololona.

[36] ———, *Homological algebra of mirror symmetry*, Proceedings of the International Congress of Mathematicians (Zürich, 1994), Birkhäuser, 1995, pp. 120–139.

[37] M. Kontsevich and Y. Soibelman, *Homological mirror symmetry and torus fibrations*, Symplectic geometry and mirror symmetry, World Scientific, 2001, pp. 203–263.

[38] D. Kwon and Y.-G. Oh, *Structure of the image of (pseudo)-holomorphic discs with totally real boundary conditions*, Comm. Anal. Geom. **8** (2000), 31–82.

[39] L. Lazzarini, *Existence of a somewhere injective pseudo-holomorphic disc*, Geom. Funct. Anal. **10** (2000), 829–862.

[40] K. Lefevre, *Sur les A_∞-catégories*, Ph.D. thesis, Université Paris 7, 2002.

[41] M. Levine, *Mixed motives*, Amer. Math. Soc., 1998.

[42] J.-L. Loday, *Cyclic homology*, Springer, 2nd edition, 1997.

[43] G. Mikhalkin, *Amoebas of algebraic varieties*, Preprint math.AG/0108225.

[44] S. Montgomery, *Hopf algebras and their actions on rings*, CBMS Regional Conference Series, vol. 82, Amer. Math. Soc., 1992.

[45] A. Neeman, *Triangulated categories*, Annals of Math. Studies, vol. 148, Princeton Univ. Press, 2001.

[46] T. Nishinou, *Lagrangian submanifold associated with a degenerating family of stable bundles on t^4*, Preprint math.SG/0301324.

[47] Y.-G. Oh, *Floer cohomology, spectral sequences, and the Maslov class of Lagrangian embeddings*, Int. Math. Res. Notices (1996), 305–346.

[48] S. Piunikhin, D. Salamon, and M. Schwarz, *Symplectic Floer-Donaldson theory and quantum cohomology*, Contact and symplectic geometry (C. B. Thomas, ed.), Cambridge Univ. Press, 1996, pp. 171–200.

[49] A. Polishchuk, *Massey and Fukaya products on elliptic curves*, Adv. Theor. Math. Phys. **4** (2000), 1187–1207.

[50] A. Polishchuk and E. Zaslow, *Categorical mirror symmetry: the elliptic curve*, Adv. Theor. Math. Phys. **2** (1998), 443–470.

[51] J. Robbin and D. Salamon, *The Maslov index for paths*, Topology **32** (1993), 827–844.

[52] W.-D. Ruan, *Newton polygon and string diagram*, Preprint math.DG/0011012.

[53] ———, *Lagrangian torus fibration of quintic hypersurfaces. I. Fermat quintic case*, Winter School on Mirror Symmetry, Vector Bundles and Lagrangian Submanifolds (Cambridge, MA, 1999), AMS/IP Stud. Adv. Math., vol. 23, Amer. Math. Soc., 2001, pp. 297–332.

[54] ———, *Lagrangian torus fibration of quintic Calabi-Yau hypersurfaces. II. Technical results on gradient flow construction*, J. Symplectic Geom. **1** (2002), 435–521.

[55] A. Rudakov et al., *Helices and vector bundles: Seminaire Rudakov*, LMS Lecture Note Series, vol. 148, Cambridge University Press, 1990.

[56] D. Salamon and E. Zehnder, *Morse theory for periodic solutions of Hamiltonian systems and the Maslov index*, Comm. Pure Appl. Math. **45** (1992), 1303–1360.

[57] P. Seidel, *Graded Lagrangian submanifolds*, Bull. Soc. Math. France **128** (2000), 103–146.

[58] ———, *Vanishing cycles and mutation*, Proceedings of the 3rd European Congress of Mathematics (Barcelona), vol. 2, Birkhäuser, 2001, pp. 65–85.

[59] ———, *Fukaya categories and deformations*, Proceedings of the International Congress of Mathematicians (Beijing), vol. 2, Higher Ed. Press, 2002, pp. 351–360.

[60] ———, *A long exact sequence for symplectic Floer cohomology*, Topology **42** (2003), 1003–1063.

[61] ———, *Fukaya categories and Picard-Lefschetz theory*, Manuscript in preparation, 2004?

[62] P. Seidel and R. Thomas, *Braid group actions on derived categories of coherent sheaves*, Duke Math. J. **108** (2001), 37–108.

[63] V. De Silva, *Products in the symplectic Floer homology of Lagrangian intersections*, Ph.D. thesis, Oxford University, 1998.

[64] N. Spaltenstein, *Resolutions of unbounded complexes*, Compositio Math. **65** (1988), 121–154.

[65] R. Thomas and S.-T. Yau, *Special Lagrangians, stable bundles and mean curvature flow.*, Comm. Anal. Geom. **10** (2002), 1075–1113.

[66] T. Tradler, *Infinity-inner-products on A-infinity-algebras*, Preprint math.AT/0108027.